

**TAGAMET- cimetidine tablet**  
**Medtech Products Inc.**

-----

**Tagamet**

***Drug Facts***

***Active ingredient (in each tablet)***

*Cimetidine 200mg*

***Purpose***

*Acid reducer*

***Uses***

- **relieves** heartburn associated with acid indigestion and sour stomach
- **prevents** heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

***Warnings***

**Allergy alert:** Do not use if you are allergic to cimetidine or other acid reducers.

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**Ask a doctor before use if you have**

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- kidney disease
- liver disease

**Ask a doctor or pharmacist before use if  
you are taking a prescription drug.**

Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if**

- you need to take this product for more than 14 days
- stomach pain continues
- your heartburn continues or worsens

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

***Directions***

- adults and children 12 years and over:
  - to **relieve** symptoms, swallow 1 tablet with a glass of water
  - to **prevent** symptoms, swallow 1 tablet with a glass of water **right before or any time up to 30 minutes before** eating food or drinking beverages that cause heartburn
  - do not take more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

***Other information***

store at 15-30°C (59-86°F)

***Inactive ingredients***

cellulose, corn starch, hypromellose, magnesium stearate, polyethylene glycol, polysorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

**Questions?**

Call toll-free **1-800-482-4394** weekdays

**PRINCIPAL DISPLAY PANEL**

Tagamet® HB 200  
Cimetidine Tablets 200 mg  
Acid Reducer  
70 tablets (70 doses)

TAKE ANYTIME YOU NEED IT:  
before meal • during meal • after meal

# Tagamet<sup>®</sup> HB 200

Cimetidine Tablets 200 mg  
Acid Reducer

**JUST ONE TABLET**  
RELIEVES and PREVENTS  
Heartburn and Acid Indigestion

Actual size

**70 TABLETS**  
(70 DOSES)

Keep cards for future reference.

Tagamet<sup>®</sup> HB 200 tablets are individually sealed in a blister pack. Do not use if blister or seal is open or torn.

**Tagamet<sup>®</sup> HB 200**  
Cimetidine Tablets 200 mg  
Acid Reducer

**Just ONE TABLET**  
RELIEVES and PREVENTS  
Heartburn and  
Acid Indigestion

### Drug Facts

| Active ingredient (in each tablet) | Purpose      |
|------------------------------------|--------------|
| Cimetidine 200 mg                  | Acid reducer |

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

**Warnings**

**Allergy alert:** Do not use if you are allergic to cimetidine or other acid reducers.

**Do not use** if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have**

- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
- kidney disease
- liver disease

**Ask a doctor or pharmacist before use if you are taking** a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if** you need to take this product for more than 14 days

- stomach pain continues
- your heartburn continues or worsens

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) immediately.

**Directions**

Adults and children 12 years and over:

- to relieve symptoms, swallow 1 tablet with a glass of water
- to prevent symptoms, swallow 1 tablet with a glass of water right before or any time up to 30 minutes before eating food or drinking beverages that cause heartburn
- do not take more than 2 tablets in 24 hours.

Children under 12 years: ask a doctor

**Other information** store at 15°-30°C (59°-86°F)

**Inactive ingredients** cellulose, corn starch, hypromellose, magnesium stearate, polyethylene glycol, polysorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

**Questions?** Call toll-free 1-800-482-4394 weekdays.

## PRINCIPAL DISPLAY PANEL

Tagamet<sup>®</sup> HB 200  
Cimetidine Tablets 200 mg  
Acid Reducer  
30 tablets (30 doses)

TAKE ANYTIME YOU NEED IT:  
before meal • during meal • after meal

# Tagamet<sup>®</sup> HB 200

Cimetidine Tablets 200 mg  
Acid Reducer

**JUST ONE TABLET**  
RELIEVES and PREVENTS  
Heartburn and Acid Indigestion

Actual size

**30 TABLETS**  
(30 DOSES)

**Tips for Managing Heartburn**

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

### Drug Facts

| Active ingredient (in each tablet) | Purpose      |
|------------------------------------|--------------|
| Cimetidine 200 mg                  | Acid reducer |

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

**Warnings**

**Allergy alert:** Do not use if you are allergic to cimetidine or other acid reducers.

**Do not use** if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have**

- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
- kidney disease
- liver disease

**Ask a doctor or pharmacist before use if you are taking** a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if** you need to take this product for more than 14 days

- stomach pain continues
- your heartburn continues or worsens

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

Adults and children 12 years and over:

- to relieve symptoms, swallow 1 tablet with a glass of water
- to prevent symptoms, swallow 1 tablet with a glass of water right before or any time up to 30 minutes before eating food or drinking beverages that cause heartburn
- do not take more than 2 tablets in 24 hours

Children under 12 years: ask a doctor

**Other information** store at 15°-30°C (59°-86°F)

**Inactive ingredients** cellulose, corn starch, hypromellose, magnesium stearate, polyethylene glycol, polysorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

**Questions?** Call toll-free 1-800-482-4394 weekdays.

# TAGAMET

cimetidine tablet

## Product Information

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:63029-222 |
| <b>Route of Administration</b> | ORAL           |                           |               |

## Active Ingredient/Active Moiety

| <b>Ingredient Name</b>  | <b>Basis of Strength</b> | <b>Strength</b> |
|---|--------------------------|-----------------|
| <b>CIMETIDINE</b> (UNII: 80061L1WGD) (CIMETIDINE - UNII:80061L1WGD) | CIMETIDINE               | 200 mg          |

## Inactive Ingredients

| <b>Ingredient Name</b>  | <b>Strength</b> |
|---|-----------------|
| <b>STARCH, CORN</b> (UNII: O8232NY3SJ)                          |                 |
| <b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)                         |                 |
| <b>MAGNESIUM STEARATE</b> (UNII: 70097M6I3O)                    |                 |
| <b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)      |                 |
| <b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)                        |                 |
| <b>POVIDONE</b> (UNII: FZ989GH94E)                              |                 |
| <b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)                 |                 |
| <b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2) |                 |
| <b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)                      |                 |
| <b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)            |                 |
| <b>FD&amp;C BLUE NO. 1 ALUMINUM LAKE</b> (UNII: J9EQA3S2JM)     |                 |

## Product Characteristics

|                 |         |                     |             |
|-----------------|---------|---------------------|-------------|
| <b>Color</b>    | white   | <b>Score</b>        | no score    |
| <b>Shape</b>    | DIAMOND | <b>Size</b>         | 13mm        |
| <b>Flavor</b>   |         | <b>Imprint Code</b> | TAGAMET;200 |
| <b>Contains</b> |         |                     |             |

## Packaging

| <b>#</b> | <b>Item Code</b> | <b>Package Description</b>                              | <b>Marketing Start Date</b> | <b>Marketing End Date</b> |
|----------|------------------|---|-----------------------------|---------------------------|
| 1        | NDC:63029-222-02 | 1 in 1 BOX  | 06/01/2012                  |                           |
| 1        |                  | 6 in 1 BLISTER PACK; Type 0: Not a Combination Product  |                             |                           |
| 2        | NDC:63029-222-01 | 1 in 1 BOX  | 06/01/2012                  |                           |
| 2        |                  | 30 in 1 BLISTER PACK; Type 0: Not a Combination Product |                             |                           |
| 3        | NDC:63029-222-03 | 1 in 1 BOX  | 06/01/2012                  |                           |
| 3        |                  | 50 in 1 BLISTER PACK; Type 0: Not a Combination Product |                             |                           |

|    |                  |   |            |  |
|----|------------------|---|------------|--|
| 4  | NDC:63029-222-04 | 4 in 1 BOX  | 06/01/2019 |  |
| 4  |                  | 10 in 1 BLISTER PACK; Type 0: Not a Combination Product |            |  |
| 5  | NDC:63029-222-05 | 4 in 1 BOX  | 06/01/2019 |  |
| 5  |                  | 9 in 1 BLISTER PACK; Type 0: Not a Combination Product  |            |  |
| 6  | NDC:63029-222-70 | 1 in 1 CARTON   | 06/01/2012 |  |
| 6  |                  | 70 in 1 BOTTLE; Type 0: Not a Combination Product       |            |  |
| 7  | NDC:63029-222-06 | 7 in 1 BOX  | 06/01/2012 |  |
| 7  |                  | 10 in 1 BLISTER PACK; Type 0: Not a Combination Product |            |  |
| 8  | NDC:63029-222-09 | 1 in 1 CARTON   | 06/01/2012 |  |
| 8  |                  | 90 in 1 BOTTLE; Type 0: Not a Combination Product       |            |  |
| 9  | NDC:63029-222-11 | 1 in 1 CARTON   | 06/01/2012 |  |
| 9  |                  | 100 in 1 BOTTLE; Type 0: Not a Combination Product      |            |  |
| 10 | NDC:63029-222-12 | 2 in 1 BOX  | 06/01/2012 |  |
| 10 |                  | 6 in 1 BLISTER PACK; Type 0: Not a Combination Product  |            |  |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| NDA                | NDA020238                                | 06/01/2012           |                    |

## TAGAMET

cimetidine tablet

### Product Information

|                         |                |                    |               |
|-------------------------|----------------|--------------------|---------------|
| Product Type            | HUMAN OTC DRUG | Item Code (Source) | NDC:63029-223 |
| Route of Administration | ORAL           |                    |               |

### Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength | Strength |
|--|-------------------|----------|
| CIMETIDINE (UNII: 80061L1WGD) (CIMETIDINE - UNII:80061L1WGD) | CIMETIDINE        | 200 mg   |

### Inactive Ingredients

| Ingredient Name                 | Strength |
|---------------------------------|----------|
| STARCH, CORN (UNII: O8232NY3SJ) |          |

|   |  |
|---|--|
| <b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)                         |  |
| <b>MAGNESIUM STEARATE</b> (UNII: 70097M6I3O)                    |  |
| <b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)      |  |
| <b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)                        |  |
| <b>POVIDONE</b> (UNII: FZ989GH94E)                              |  |
| <b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)                 |  |
| <b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2) |  |
| <b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)                      |  |
| <b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)            |  |
| <b>FD&amp;C BLUE NO. 1 ALUMINUM LAKE</b> (UNII: J9EQA3S2JM)     |  |

## Product Characteristics

|                 |            |                     |             |
|-----------------|------------|---------------------|-------------|
| <b>Color</b>    | blue       | <b>Score</b>        | 2 pieces    |
| <b>Shape</b>    | DIAMOND    | <b>Size</b>         | 13mm        |
| <b>Flavor</b>   | PEPPERMINT | <b>Imprint Code</b> | TAGAMET;200 |
| <b>Contains</b> |            |                     |             |

## Packaging

| # | Item Code        | Package Description                                     | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:63029-223-30 | 3 in 1 BOX  | 02/01/2023           |                    |
| 1 |                  | 10 in 1 BLISTER PACK; Type 0: Not a Combination Product |                      |                    |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| NDA                | NDA020238                                | 02/01/2023           |                    |

**Labeler** - Medtech Products Inc. (122715688)