

**MEDIQUE AT HOME DECOREL FORTE- acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride tablet, film coated**  
**Unifirst First Aid Corporation**

-----  
**Medique @ Home Decorel Forte**

***Drug Facts***

***Active ingredients (in each caplet)***

Acetaminophen 325 mg  
Dextromethorphan HBr 10 mg  
Guaifenesin 100 mg  
Phenylephrine HCl 5 mg

***Purpose***

Pain reliever/fever reducer  
Cough suppressant  
Expectorant  
Nasal decongestant

***Uses***

Temporarily relieves these symptoms due to the common cold:

- headache
- nasal congestion
- cough
- minor aches and pains
- sore throat
- sinus congestion and pressure
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make cough more productive

Temporarily reduces fever.

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is

accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription).

If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug.

If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough that lasts as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin.

**When using this product do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed**

**Adults and children: (12 years and over):**

- take 2 caplets every 4 hours
- not to take more than 10 caplets in 24 hours

**Children under 12 years:** ask a doctor

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- **TAMPER EVIDENT** child resistant packets
- do not use any opened or torn packets

**Inactive ingredients**

corn starch, crospovidone, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

**Questions or comments?** 1-800-634-7680

Medique® @ Home

Decorel Forte

Acetaminophen 325 mg • **Pain Reliever/Fever Reducer**

Dextromethorphan HBr 10 mg • **Cough Suppressant**

Guaifenesin 100 mg • **Expectorant**

Phenylephrine HCl 5 mg • **Nasal Decongestant**

Pull to Open

Child-Resistant Packets

\*not actual size

50 Packets

(2 Caplets Per Packet)



**Medique Home**  
**Decorel Forte**  
Acetaminophen 325 mg • Pain Reliever/  
Fever Reducer  
Dextromethorphan HBr 10 mg • Cough  
Suppressant  
Guaifenesin 100 mg • Expectorant  
Phenylephrine HCl 5 mg • Nasal Decongestant  
50 Packets (2 Caplets Per Packet)



**Medique Home**  
**Decorel Forte**  
Acetaminophen 325 mg • Pain Reliever/  
Fever Reducer  
Dextromethorphan HBr 10 mg • Cough  
Suppressant  
Guaifenesin 100 mg • Expectorant  
Phenylephrine HCl 5 mg • Nasal Decongestant  
50 Packets (2 Caplets Per Packet)



**Medique Home**  
**Decorel Forte**  
Acetaminophen 325 mg • Pain Reliever/  
Fever Reducer  
Dextromethorphan HBr 10 mg • Cough  
Suppressant  
Guaifenesin 100 mg • Expectorant  
Phenylephrine HCl 5 mg • Nasal Decongestant  
50 Packets (2 Caplets Per Packet)

**Drug Facts**

Active Ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan hydrobromide 10 mg	Cough suppressant
Guaifenesin 100 mg	Expectorant
Phenylephrine hydrochloride 5 mg	Nasal decongestant

**Uses**  
Temporarily relieves these symptoms due to the common cold:  
 • headache • nasal congestion • cough • minor aches and pains • sore throat • sinus congestion and pressure  
 • helps loosen phlegm (mucus) and thin bronchial secretions to clear bronchial tubes and make coughs more productive  
 • temporarily reduces fever

**Warnings**  
 • Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:  
 • more than 4,000 mg of acetaminophen in 24 hours  
 • with other drugs containing acetaminophen  
 • 3 or more alcoholic drinks every day while using this product  
 • Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • blisters • rash • skin redness  
 If a skin reaction occurs, stop use and seek medical help right away.  
 • **See throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

**Do not use**  
 • with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 • if you have ever had an allergic reaction to the product or any of its ingredients  
 • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have:**  
 • liver disease • heart disease • high blood pressure • thyroid disease • diabetes  
 • difficulty in urination due to enlargement of the prostate gland  
 • cough that occurs with too much phlegm (mucus)  
 • persistent or chronic cough that lasts or occurs with smoking, asthma, chronic bronchitis or emphysema

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

**When using this product do not exceed recommended dosage.**

**Stop use and ask a doctor if:**  
 • nervousness, dizziness, or sleeplessness occur • pain, cough or nasal congestion gets worse or lasts for more than 7 days  
 • fever gets worse or lasts more than 3 days • redness or swelling is present  
 • new symptoms occur • cough comes back or occurs with rash or headache that lasts  
 These could be signs of a serious condition. (continued on back panel)

**Drug Facts (continued)**

If pregnant or breast feeding, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
 • do not take more than directed  
 Adults and children (12 years and over) • take 2 caplets every 4 hours  
 • not to take more than 10 caplets in 24 hours  
 Children under 12 years ask a doctor

**Other information**  
 • store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)  
 • **TRIPLEX EVIDENT** child resistant packets  
 • do not use any opened or torn packets

**Inactive ingredients**  
 corn starch, croscopolone, FD&C yellow #6 aluminum lake, magnesium stearate, methocel, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, polyvinyl alcohol, sodium starch glycolate, stearic acid, talc, titanium dioxide

**Questions or comments?** 1-800-634-7353

MEDIQUE AT HOME DECOREL FORTE			
acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride tablet, film coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-735
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg	
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg	
Inactive Ingredients			
Ingredient Name	Strength		
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)			
<b>CROSPVIDONE</b> (UNII: 2S7830E561)			
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)			
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)			
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)			
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)			

<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	

### Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (Caplet)	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	44;546
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-735-33	50 in 1 BOX	09/26/2022	
1		2 in 1 PACKET; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/26/2022	

**Labeler** - Unifirst First Aid Corporation (832947092)

Revised: 1/2024

Unifirst First Aid Corporation