ACETAMINOPHEN AND IBUPROFEN (NSAID)- acetaminophen and ibuprofen (nsaid) tablet RITE AID

Active ingredients (in each caplet)

Acetaminophen 250 mg

Ibuprofen 125 mg (NSAID**)

**nonsteroidal anti-inflammatory drug

Purposes

Pain reliever

Pain reliever

Uses

- temporarily relieves minor aches and pains due to:
- headache toothache backache
- menstrual cramps muscular aches minor pain of arthritis

Acetaminophen liver damage warning:

This product contains acetaminophen. Severe liver damage may occur if you take:

- with other drugs containing acetaminophen
- more than 6 caplets in 24 hours, which is the maximum daily amount for this product
- 3 or more alcoholic drinks every day while using this product

Acetaminophen allergy alert:

may cause severe skin reactions. Symptoms may include:

■ skin reddening ■ blisters ■ rash

If skin reaction occurs, stop use and seek medical help right away.

NSAID allergy alert:

ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

■ hives ■ facial swelling ■ asthma (wheezing)

 \blacksquare shock \blacksquare skin reddening \blacksquare rash \blacksquare blisters

If an allergic reaction occurs, stop use and seek medical help right away.

NSAID stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

are age 60 or older

■ have had stomach ulcers or bleeding problems

■ take a blood thinning (anticoagulant) or steroid drug

■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprefer_paperover_or_others]

- ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning:

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

 \blacksquare if you have ever had an allergic reaction to acetaminophen or any other pain reliever

right before or after heart surgery

Ask a doctor before use if

■ you have liver disease

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers
- you have a history of stomach problems, such as heartburn

■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke

you are taking a diuretic

Ask a doctor or pharmacist before use if you are

■ under a doctor's care for any serious condition

■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin

taking any other drug

When using this product

Take with food or milk if stomach upset occurs

Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding:

- \blacksquare feel faint \blacksquare vomit blood \blacksquare have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
- \blacksquare chest pain \blacksquare trouble breathing
- \blacksquare weakness in one part or side of body \blacksquare slurred speech \blacksquare leg swelling
- pain gets worse or lasts more than 10 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ do not take more than directed

adults and children 12 years and over

take 2 caplets every 8 hours while symptoms persist

children under 12 years

- ask a doctor
- do not take more than 6 caplets in 24 hours, unless directed by a doctor

Other information

- read all warnings and directions before use. Keep carton.
- store at 20 to 25°C (68 to 77°F)
- avoid excessive heat above 40°C (104°F)


Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, crospovidone, ferric oxide red, ferric oxide yellow, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, povidone, pregelatinized starch, sodium lauryl sulfate, stearic acid and titanium dioxide.

Questions or comments?

Principal display panel

| 7000000341 20000004129 | | ACETAMINOPHEN + IBUPROFEN ACETAMINOPHEN 250 mg AND BUPROFEN (NSAID) 125 mg TALLETS PAIN RELIEVER | CONTAINS 2 MEDICINES |
|--|--|--|---|
| Market location Day factor location Day factor location Day factor location Day fact | Construction Description Remain and and and and and and and and and an | ACETAMINOPHEN + IBUPROFEN ACETAMINOPHEN 250 mg AND IBUPROFEN (NSAID) 125 mg TABLETS PRIN RELEVER NORTH | Image: State of the state o |
| 200000004179 70000003241 | | 0 11822-77104 7 | |



ACETAMINOPHEN AND IBUPROFEN (NSAID)

acetaminophen and ibuprofen (nsaid) tablet

| acetaminophen and ibuprofer | i (iisaiu) tablet | | | | |
|---------------------------------|--------------------------|-----------------|-------------|----------|----------|
| | | | | | |
| Product Information | | | | | |
| Product Type | HUMAN OTC DRUG | ltem Code (Sou | rce) | NDC:1182 | 2-1131 |
| Route of Administration | ORAL | | | | |
| | | | | | |
| Active Ingredient/Active | Majaty | | | | |
| Active ingredient/Active | Molecy | | | | |
| Ingre | edient Name | | Basis of S | trength | Strength |
| IBUPROFEN (UNII: WK2XYI10QM) (I | BUPROFEN - UNII:WK2XYI10 | QM) | IBUPROFEN | | 125 mg |
| ACETAMINOPHEN (UNII: 36209ITL | 9D) (ACETAMINOPHEN - UN | III:36209ITL9D) | ACETAMINOPH | HEN | 250 mg |
| | | | | | |
| | | | | | |
| Inactive Ingredients | | | | | |
| | Ingredient Name | | | St | rength |
| CROSCARMELLOSE SODIUM (UN | II: M28OL1HH48) | | | | |
| | | | | | |

| | YELLOW (UNII: EX43802MRT) | | |
|-------------------|---|--------------|----------|
| | IE GLYCOL 400 (UNII: B697894SGQ) | | |
| STEARIC ACID | (UNII: 4ELV7Z65AP) | | |
| TITANIUM DIO | XIDE (UNII: 15FIX9V2JP) | | |
| CARNAUBA W | AX (UNII: R12CBM0EIZ) | | |
| SILICON DIOX | IDE (UNII: ETJ7Z6XBU4) | | |
| CROSPOVIDO | NE (UNII: 2S7830E561) | | |
| HYPROMELLO | SE, UNSPECIFIED (UNII: 3NXW29V3WO) | | |
| POVIDONE K9 | 0 (UNII: RDH86HJV5Z) | | |
| STARCH, COR | N (UNII: 08232NY3SJ) | | |
| FERRIC OXIDE | RED (UNII: 1K09F3G675) | | |
| MAGNESIUM S | TEARATE (UNII: 70097M6I30) | | |
| SODIUM LAUR | YL SULFATE (UNII: 368GB5141J) | | |
| | | | |
| | | | |
| Product Ch | naracteristics | | |
| Color | yellow (Light yellow to yellow colored) | Score | no score |
| Shape | CAPSULE | Size | 14mm |
| Flavor | | Imprint Code | G;131 |
| Contains | | | |

Packaging

| # | ltem Code | Package Description | Marketing Start Date | Marketing End Date |
|---|-------------------------------------|--|-------------------------|-----------------------|
| | NDC:11822- 1131-6 | 216 in 1 BOTTLE; Type 0: Not a Combination Product | 07/21/2023 | |
| | NDC:11822- 1131-4 | 144 in 1 BOTTLE; Type 0: Not a Combination Product | 07/21/2023 | |
| | | | | |
| | | | | |
| M | arketing | Information | | |
| M | arketing Marketing Category | I nformation Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |

Labeler - RITE AID (014578892)

Revised: 1/2024

RITE AID