

PYRANTEL PAMOATE- pyrantel pamoate suspension

ProMex LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Pyrantel Pamoate

Drug Facts

Active Ingredient (in each 1mL)

Pyrantel Pamoate 144 mg/mL
(50mg pyrantel base per mL)

Purpose

Anthelmintic

Use

for the treatment of pinworms.

Warnings

*Abdominal cramps, nausea, vomiting, diarrhea, headache, or dizziness sometimes occurs after taking this drug. If any of these conditions persist, consult a doctor.

*If you are pregnant or have liver disease, do not take this product unless directed by a doctor.

Directions

Shake well before use. Adults, children 12 years of age and over, and children 2 years to under 12 years of age: Oral dosage is a single dose of 5 milligrams of pyrantel base per pound, or 11 milligrams per kilogram, of body weight. Do not exceed 1 gram (1000 mg).

Weight	Dosage (Taken as a single dose)
Under 25 lbs or 2 yrs.	Don't use unless directed by a doctor
25-37 lbs	½ teaspoonful
38-62 lbs	1 teaspoonful
63-87 lbs	1 ½ teaspoonful
88-112 lbs	2 teaspoonful

113-137 lbs	2 ½ teaspoonful
138-162 lbs	3 teaspoonful
163-187lbs	3 ½ teaspoonful
over 188 lbs	4 teaspoonful

***Read package insert carefully before taking this medication.** Take only according to directions and do not exceed the recommended dosage unless directed by a doctor. Medication should only be taken one time as a single dose; do not repeat treatment unless directed by a doctor. When one individual in a household has pinworms, the entire household should be treated unless otherwise advised. See Warnings. If any worms other than pinworms are present before or after treatment consult a doctor. If any symptoms or pinworms are still present after treatment, consult a doctor.

*This product can be taken any time of day, with or without meals, it may be taken alone or with milk or fruit juice. Use of a laxative is not necessary prior to, during or after medication.

Keep out of Reach of Children. In case of accidental overdose/ingestion contact a poison control center immediately.

Inactive Ingredients

Purified Water, Glycerin, Sugar, Flavor, Xanthan Gum, Citric Acid, Sodium Benzoate and Potassium Sorbate


PRINCIPAL DISPLAY PANEL - 30 ml Bottle Carton

VERMISOL

SUSPENSION

1 oz (30 ml)





S U S P E N S I O N

1 oz (30 ml)

Almacénesse a temperaturas entre 15°-30°C (59°-86°F)

DATOS DEL MEDICAMENTO (continuado)

Mantener fuera del alcance de los niños. En caso de sobredosis / ingestión accidental contacte un centro de control de envenenamiento inmediatamente.

Ingredientes inactivos: Agua purificada, glicerina, azúcar, goma Xanthan, sabor, ácido cítrico, benzoato de sodio, sorbato de potasio

Distributed By:

PRO-MEX®

P.O. Box 63-0716 - Miami, FL. 33163

UPC
184125-00026

mayores de 2 a menores de 12 años de edad: la dosis oral es una sola dosis de 5 miligramos de la base de pirantel por libra, o 11 miligramos por kilogramo de peso corporal. No exceda de 1 gramo (1,000 mg).

Peso	Dosis (tome una sola dosis)
Menos de 25 lbs o 2 años	No usar sin aviso de un médico
25-37 lb	½ cucharadita
38-62 lbs	1 cucharadita
63-87 lbs	1 ½ cucharadita
88-112 lbs	2 cucharadita
113-137 lbs	2 ½ cucharadita
138-162 lbs	3 cucharadita
163-187 lbs	3 ½ cucharadita
188 lbs o mas	4 cucharadita

Lea el inserto cuidadosamente antes de tomar este medicamento. Tome solamente según las instrucciones y no exceder la dosis recomendada a menos que lo indique un médico. Medicamento sólo debe tomarse una vez en una sola dosis, no repetir el tratamiento a menos que lo indique un médico. Cuando una persona en un hogar tiene oxiuros, toda la familia debe ser tratada a menos que se aconseje lo contrario. Véase Advertencias. Si algún gusano que no sea pinworms están presentes antes o después del tratamiento, consulte a un médico. Si los síntomas o los oxiuros todavía están presentes después del tratamiento, consulte a un médico.

Este producto se puede tomar a cualquier hora del día, con o sin comidas. Se puede tomar solo o con leche o jugo de frutas. El uso de un laxante no es necesario antes, durante o después de la medicación.

S U S P E N S I O N

(Pyrantel Pamoate)

Tamper Evident: Do not use if seal is broken
No Usar Si el Sello esta roto

VERMISOL

VERMISOL



S U S P E N S I O N

1 oz (30 ml)

Store at 15°-30°C (59°-86°F)



P
VCOE_009

Drug Facts

Active Ingredient (in each 1mL) **PURPOSE**
Pyrantel Pamoate 144 mg/mL Anthelmintic
(50mg pyrantel base per mL)

Use: for the treatment of pinworms.

Warnings: ■ Abdominal cramps, nausea, vomiting, diarrhea, headache, or dizziness sometimes occur after taking this drug. If any of these conditions persist consult a doctor.
■ If you are pregnant or have liver disease, do not take this product unless directed by a doctor.

Directions: Shake well before use. Adults, children 12 years of age and over, and children 2 years to under 12 years of age: Oral dosage is a single dose of 5 milligrams of pyrantel base per pound, or 11 milligrams per kilogram, of body weight. Do not exceed 1 gram (1000 mg).

Weight	Dosage (Taken as a single dose)
Under 25 lbs or 2 yrs	Don't use unless directed by a doctor
25-37 lbs	½ teaspoonful
38-62 lbs	1 teaspoonful
63-87 lbs	1 ½ teaspoonful
88-112 lbs	2 teaspoonful
113-137 lbs	2 ½ teaspoonful
138-162 lbs	3 teaspoonful
163-187 lbs	3 ½ teaspoonful
over 188 lbs	4 teaspoonful

■ Read package insert carefully before taking this medication. Take only according to directions and do not exceed the recommended dosage unless directed by a doctor. Medication should only be taken one time as a single dose; do not repeat treatment unless directed by a doctor. When one individual in a household has pinworms, the entire household should be treated unless otherwise advised. See Warnings. If any worms other than pinworms are present before or after treatment, consult a doctor. If any symptoms or pinworms are still present after treatment, consult a doctor.

■ This product can be taken any time of day, with or without meals. It may be taken alone or with milk or fruit juice. Use of a laxative is not necessary prior to, during, or after medication.

Keep out of Reach of Children. In case of accidental overdose/ingestion contact a poison control center immediately.

Inactive Ingredients: Purified Water, Glycerin, Sugar, Flavor, Xanthan Gum, Citric Acid, Sodium Benzoate and Potassium Sorbate

PC-134.00

PYRANTEL PAMOATE

pyrantel pamoate suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58988-1770	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
Pyrantel Pamoate (UNII: 81BK194Z5M) (Pyrantel - UNII:4QIH0N49E7)		Pyrantel	144 mg in 1 mL	
Inactive Ingredients				
Ingredient Name			Strength	
Water (UNII: 059QF0KO0R)				
Citric Acid Monohydrate (UNII: 2968PHW8QP)				
Glycerin (UNII: PDC6A3C0OX)				
Sucrose (UNII: C151H8M554)				
Xanthan Gum (UNII: TTV12P4NEE)				
Sodium Benzoate (UNII: OJ245FE5EU)				
Potassium Sorbate (UNII: 1VPU26JZZ4)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58988-1770-1	1 in 1 CARTON	09/27/2002	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC MONOGRAPH FINAL	part357B	09/27/2002		

Labeler - ProMex LLC (789974388)

Revised: 1/2022

ProMex LLC