

**ACID REDUCER 150- ranitidine hydrochloride tablets 150mg tablet, coated**  
**Safeway Inc**

-----  
**Drug Facts**

**Active ingredient (in each tablet)**

Ranitidine 150 mg (as ranitidine hydrochloride, USP 168 mg)

**Purpose**

Acid reducer

**Use(s)**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by certain foods and beverages

**Warnings**

**Allergy alert:** Do not use if you are allergic to ranitidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor

**Ask a doctor before use if you have**

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

**Stop use and ask doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding**

ask a health professional before use.

## **Keep out of reach of children**

In case of overdose, get medical help or contact a Poison Control Center right away.

## **Directions**

- adults and children 12 years and over:
  - to **relieve** symptoms, swallow 1 tablet with a glass of water
  - to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
  - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

## **Other information**

- do not use if printed foil under bottle cap is open or torn
- store at 20°-25°C (68°-77°F)
- avoid excessive heat or humidity
- protect from light
- this product is sugar free

## **Inactive ingredients**

FD&C red #40 aluminum lake, hypromellose, iron oxide black, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide


## **Questions**

call **1-888-375-3784**

**SEALED WITH PRINTED FOIL UNDER CAP FOR YOUR PROTECTION.**

**IMPORTANT: Read the directions, consumer information leaflet and warnings before use. Keep the carton. It contains important information.**

**container label**



## CARTON LABELING - PRINCIPAL DISPLAY PANEL



**Signature Care**<sup>TM</sup>  
Quality Guaranteed

NDC 21130-404-34

**Maximum Strength Ranitidine Tablets USP, 150 mg**

**Acid Reducer**

Prevents & relieves heartburn associated with acid indigestion & sour stomach

**24 TABLETS (24 DOSES)**

**SEAL WITH PRINTED FOIL UNDER CAP FOR YOUR PROTECTION**

**IMPORTANT:** This label does not contain full product information. See carton for complete information. Read the directions, consumer information leaflet and warnings before use. Retain carton for reference.

**Active ingredient (in each tablet)** Ranitidine 150 mg (as ranitidine hydrochloride, USP 168 mg) **Purpose** Acid reducer

**Uses** ■ relieves heartburn associated with acid indigestion and sour stomach ■ prevents heartburn associated with acid indigestion and sour stomach brought on by certain foods and beverages ■ prevents heartburn associated with acid indigestion and sour stomach brought on by certain foods and beverages

**Warnings** ■ **Allergy alert:** Do not use if you are allergic to ranitidine or other acid reducers

**Do not use** ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ with other acid reducers ■ if you have kidney disease, except under the advice and supervision of a doctor

**Ask a doctor before use** if you have ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

**Drug Facts (continued)**

**Stop use and ask a doctor if** ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days

**If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.**

**Directions** ■ adults and children 12 years and over: ■ to relieve symptoms, swallow 1 tablet with a glass of water ■ to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn ■ can be used up to twice daily (do not take more than 2 tablets in 24 hours) ■ children under 12 years: ask a doctor

**Other information** ■ do not use if printed foil under bottle cap is open or torn ■ store at 20°-25°C (68°-77°F) ■ avoid excessive heat or humidity ■ protect from light ■ this product is sodium and sugar free

**Inactive ingredients** FD&C red #40 aluminum lake, hypromellose, iron oxide black, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide.

**Questions?** Call 1-888-375-3784

\*This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., distributor of Zantac 150® Tablets. Zantac 150® is a registered trademark of Boehringer Ingelheim Pharmaceuticals, Inc.

**Actual Size**



**24 TABLETS (24 DOSES)**

**OUR PROMISE**  
QUALITY & SATISFACTION  
**100% GUARANTEED**  
OR YOUR MONEY BACK.

DISTRIBUTED BY:  
BETTER LIVING BRANDS LLC  
P.O. BOX 99  
PLEASANTON, CA 94566-0099  
1-888-723-3929  
www.betterlivingbrandsLLC.com

**150060428**

LOT

EXP

## ACID REDUCER 150

ranitidine hydrochloride tablets 150mg tablet, coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:21130-404(NDC:55111-404)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
RANITIDINE HYDROCHLORIDE (UNII: BK76465IHM) (RANITIDINE - UNII:884KT10YB7)	RANITIDINE	150 mg

### Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HYPROMELLOSE 2910 (6 MPAS) (UNII: 0WZ8WG20P6)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
magnesium stearate (UNII: 70097M6I30)	
cellulose, microcrystalline (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
titanium dioxide (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	PINK	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	R150
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-404-34	1 in 1 CARTON	05/14/2012	09/30/2019
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:21130-404-50	1 in 1 CARTON	05/14/2012	09/30/2019
2		50 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:21130-404-02	1 in 1 CARTON	05/14/2012	09/30/2019
3		200 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078192	05/14/2012	

**Labeler** - Safeway Inc (009137209)

Revised: 9/2019

Safeway Inc