

**HEADACHE, CIRCLE K- acetaminophen, caffeine, aspirin tablet tablet
Lil' Drug Store Products, Inc.**

Headache, Circle K™

Drug Facts

Active ingredients

Active ingredients (in each caplet)

Acetaminophen 250 mg

Aspirin (NSAID*) 250 mg

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purposes

Purposes

Pain reliever

Pain reliever

Pain reliever aid

Uses

Uses

temporarily relieves minor aches and pains due to:

- headache
- a cold
- arthritis
- muscular aches
- toothache
- premenstrual & menstrual cramps

Warnings

Reye's syndrome

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- any new symptoms appear

These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Directions

- **do not use more than directed**
- drink a full glass of water with each dose

adults and children (12 years and over)	take 2 caplets every 6 hours as needed; not more than 8 caplets in 24 hours
children under 12 years	ask a doctor

Other information

Other information

- store at 68°-77°F (20°-25°C); excursions permitted between 59°-86°F (15°-30°C)
- avoid excessive heat and humidity
- **each caplet contains:** Magnesium 0.58 mg

Inactive ingredients

Inactive ingredients colloidal silicon dioxide, hypromellose, microcrystalline cellulose, polyethylene glycol, povidone K30, pregelatinized starch, stearic acid, talc

Questions or comments?

Questions or comments?

Call toll-free **1-877-507-6516 (M-F 8 AM-4:30PM CST)**

Product manufactured for:

Lil' Drug Store Products, Inc., 9300 Earhart Lane SW, Cedar Rapids, IA 52404
www.lildrugstore.com 54383C-US-08-21

Proudly distributed by Circle K Stores Inc.

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Excedrin® Extra Strength.

PDP/Package

TEMPORARILY RELIEVES HEADACHES

Compare to the active ingredients in
Excedrin® Extra Strength*

CIRCLE K™

extra strength

headache

Acetaminophen 250 mg, Pain Reliever
Aspirin (NSAID) 250 mg, Pain Reliever
Caffeine 65 mg, Pain Reliever Aid

[caplet image]

actual size

6 caplets (2 per pouch)

Drug Facts (continued)	
• pain gets worse or lasts for more than 10 days • fever gets worse or lasts for more than 3 days • any new symptoms appear These could be signs of a serious condition	
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
Directions - do not use more than directed • drink a full glass of water with each dose	adults and children (12 years and over) take 2 caplets every 6 hours as needed, not more than 8 caplets in 24 hours
children under 12 years ask a doctor	
Other information • store at 88°-77° (20°-25°C); excursions permitted between 59°-86° (15°-30°C) • avoid excessive heat and humidity • each caplet contains: Magnesium 0.38 mg	
Inactive ingredients colloidal silicon dioxide, hypromellose, microcrystalline cellulose, polyethylene glycol, croscarmellose sodium, pregelatinized starch, stearic acid, talc	
Questions or comments? Call toll-free 1-877-507-6516 (M-F 8 AM-4:30PM CST)	
Product manufactured for: Lil' Drug Store Products, Inc., 9300 Earhart Lane SW, Cedar Rapids, IA 52404 www.lildrugstore.com 54383C-US-08-21 Proudly distributed by Circle K Stores Inc. *This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Excedrin® Extra Strength.	

TEMPORARILY RELIEVES HEADACHES
Compare to the active ingredients in
Excedrin® Extra Strength*

CIRCLE K™
extra strength
headache

Acetaminophen 250 mg, Pain Reliever
Aspirin (NSAID) 250 mg, Pain Reliever
Caffeine 65 mg, Pain Reliever Aid

actual size

6 caplets (2 per pouch)

6 caplets (2 per pouch)

extra strength headache

CIRCLE K™

DO NOT USE IF INNER POUCH OR CARTON IS OPEN OR DAMAGED!

Drug Facts	Purposes
Active ingredients (in each caplet)	
Acetaminophen 250 mg	Pain reliever
Aspirin (NSAID) 250 mg	Pain reliever
Caffeine 65 mg	Pain reliever aid
*nonsteroidal anti-inflammatory drug	
Uses temporarily relieves minor aches and pains due to: • headache • a cold • arthritis • muscle aches • toothache • premenstrual & menstrual cramps	
Warnings Key symptoms: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin redness • blisters • rash If a skin reaction occurs, stop use and seek medical help right away. Allergy alert: Aspirin (NSAIDs) may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • block Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: • more than 8 caplets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems	

Drug Facts (continued)

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caution warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic • you have asthma

Ask a doctor or pharmacist before use if you are taking

- a prescription drug for diabetes, gout, or arthritis
- any other drug, or are under a doctor's care for any serious condition

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - fast faint • vomit blood
 - have bloody or black stools • have stomach pain that does not get better
 - tingling in the ears or loss of hearing occurs
- painful area is red or swollen

LIFT PANEL FOR CONTINUED DRUG FACTS

HEADACHE, CIRCLE K

acetaminophen, caffeine, aspirin tablet tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-5438(NDC:68210-0107)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE K30 (UNII: U725QWY32X)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	white (white to off-white, with specks)	Score	no score
Shape	OVAL (capsule-shaped biconvex)	Size	18mm
Flavor		Imprint Code	S53
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-5438-3	3 in 1 CARTON	09/17/2021	
1		6 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	09/17/2021	

Labeler - Lil' Drug Store Products, Inc. (093103646)

Revised: 12/2023

Lil' Drug Store Products, Inc.