# ARTHRITIS PAIN RELIEVER- acetaminophen tablet, extended release WALGREENS

-----

### Acetaminophen Extended Release Tablets 650 mg - Arthritis Pain Walgreens

### **Drug Facts**

### Active ingredient (in each caplet)

Acetaminophen 650 mg

### **Purpose**

Pain reliever/fever reducer

### Uses

- temporarily relieves minor aches and pains due to:
- o minor pain of arthritis
- o muscular aches
- o backache
- o premenstrual and menstrual cramps
- o the common cold
- o headache
- o toothache
- temporarily reduces fever

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

### Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

### Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition

**If pregnant or breast-feeding,** ask a health professional before use.

### Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- Do not take more than directed
- See overdose warning

	<ul> <li>take 2 caplets every 8 hours with water</li> </ul>
	<ul> <li>swallow whole; do not crush, chew, split or dissolve</li> </ul>
adults	<ul> <li>do not take more than 6 caplets in 24 hours</li> </ul>
	<ul> <li>do not use for more than 10 days unless directed by a doctor</li> </ul>

under 18 years of age • ask a doctor

### Other information

- store between 20-25°C (68-77°F)
- The FDA approved Dissolution methods differ from USP

Inactive ingredients carnauba wax, hydroxyethyl cellulose, hypromellose, povidone, pregelatinized starch, magnesium stearate, microcrystalline cellulose, sodium starch glycolate, titanium dioxide, triacetin

Questions or comments? Call **1-877-290-4008** 

# Drug Facts (continued)

with any other drug containing acetamino phen (prescription or nonprescription). If you are not sure whether a drug contains Symptoms may include: ■ skin reddening ■ blisters ■ r If a skin reaction occurs, stop use and seek medical help right away Do not use Allergy a lert: Acetaminophen may cause severe skin reactions. ■ rash

Ask a doctor or pharmacist before use if you are taking the blood thinning

Ask a doctor before use if you have liver disease

ingredients in this product

■ if you are allergic to acetaminophen or any of the inactive

acetaminophen, aska doctor or pharmacist

Stop use and ask a doctor if

new symptoms occur

■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days

■ redness or swelling is present
These could be signs of a serious condition

Keep out of reach of children. pregnant or breast-feeding, ask a health professional before use.

notice any signs or symptoms attention is critical for ad ults as well as for children even if you do not Overdos e warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical

# Do not take more than directed Directions

See overdose warning

adults

■ take 2 caplets every 8 hours with water

swallow whole; do not crush, chew, split do not take more than 6 caplets in 24 hours or dissolve

do not use for more than 10 days unless directed by a doctor

ask a doctor

under 18 years of age

Other information

■ The FDA approved Dissolution methods differ from USP ■ store between 20-25°C (68-77°F)

**nactive ingredients** carrauba wax, hydroxyethylcellulose,

hypromeliose, magnesium stearate, microcrystalline cellulose, povidone, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

**Questions or comments?** Call 1-877-290-4008

Adhesive Area

### Inside



### signs or symptoms. \*\*Directions\*\* ■ Do not take more than directed overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Duick medical attention is critical for adults as compare to national brands. Our pharmacists recommend the Walgreens brand. We invite you to Walgreens Pharmacist Recommended! microcrystalline cellulose, sodium starch glycolate, pregelatinized starch, magnesium stearate, hydroxyethylcellulose, hypromellose, povidone, ■ store between 20-25°C (68-77°F) Other information under 18 years of age: ■ ask a doctor do not use for more than 10 days unless directed do not take more than 6 caplets in 24 hours See overdose warning well as for children even if you do not notice any professional before use. **Keep out of reach of children. Overdose warning:** In case of If pregnant or breast-feeding, ask a health signs of a serious condition worse or lasts more than 10 days 'Stop use and ask a doctor if ■ pain gets taking the blood thinning drug warfarin. Ask a doctor or pharmacist before use if you are titanium dioxide, triacetin *Inactive ingredients* carnauba wax, swallow whole; do not crush, chew, split or adults: ■ take 2 caplets every 8 hours with water ow2recycle.info *Questions or comments?* Call **1-877-290-4008** The FDA approved Dissolution methods differ redness or swelling is present. These could be new symptoms occur fever gets worse or lasts more than 3 days

## Inside

Walgreens

Adhesive Area

# Pain Reliever

ACETAMINOPHEN EXTENDED-RELEASE TABLETS USP, 650 mg PAIN RELIEVER / FEVER REDUCER

 For the temporary relief of minor arthritis pain CAPLETS\* 650 mg EACH (\*capsule-shaped bi-layer tablets)

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN

NDC 0363-9781-42

etaminophen 650 mg.....Pain reliever/fever reduce Active ingredient (in each caplet)

temporarily relieves minor aches and pains due to:

minor pain of arthritis

muscular aches

backache

premenstrual and menstrual cramps

the common cold

headache

Liver warning: This product contains acetaminophen temporarily reduces fever ■ toothache

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening Severe liver damage may occur if you take ■ more than 6 caplets in 24 hours, which is the maximum if you are allergic to acetaminophen or any of the ■ blisters ■ rash If a skin reaction occurs, stop acetaminophen (prescription or nonprescription) daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks Do not use ■ with any other drug containing you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist use and seek medical help right away. every day while using this product

No Aspirin MADE IN INDIA walgreens.com ©2022 Walgreen Co. 699R 0922 REV-0423 TEM 810295\_W00000-0000-0 W3ORG1022-F inued under label) 200 WILMOT RD., DEERFIELD, IL 60015 100% SATISFACTION GUARANTEED DISTRIBUTED BY: WALGREEN CO. Drug Facts (con

nactive ingredients in this product

Contains

Ask a doctor before use if you have liver disease."

*Drug Facts* (continued)

Lot No.:

Exp Date:

Varnish Omit Area



### **ARTHRITIS PAIN RELIEVER**

acetaminophen tablet, extended release

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-9781
Route of Administration	ORAL		

### **Active Ingredient/Active Moiety**

1000	<b>40 4</b>	iant	NI -	
mg	reu	ient	Na	me

Dubib of Street

**Basis of Strength Strength** 

ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)

ACETAMINOPHEN

650 mg

Inactive Ingredients				
Ingredient Name	Strength			
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)				
STARCH, CORN (UNII: O8232NY3SJ)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
CARNAUBA WAX (UNII: R12CBM0EIZ)				
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)				
POVIDONE K30 (UNII: U725QWY32X)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
TRIACETIN (UNII: XHX3C3X673)				

Product Characteristics				
Color	white (White to off-white)	Score	no score	
Shape	CAPSULE (Capsule-shaped tablet)	Size	19mm	
Flavor		Imprint Code	71	
Contains				

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-9781- 01	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/28/2022	
2	NDC:0363-9781- 42	1 in 1 CARTON	11/22/2022	
2		24 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA215486	09/28/2022		

## Labeler - WALGREENS (008965063)

# Registrant - TIME CAP LABORATORIES INC. (037052099)

Establishment					
Name	Address	ID/FEI	<b>Business Operations</b>		
MARKSANS PHARMA LIMITED		925822975	manufacture(0363-9781)		

Revised: 3/2024 WALGREENS