

NITROGLYCERIN LINGUAL- nitroglycerin spray

Padagis Israel Pharmaceuticals Ltd

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use NITROGLYCERIN LINGUAL SPRAY safely and effectively.

See full prescribing information for NITROGLYCERIN LINGUAL SPRAY.

NITROGLYCERIN lingual spray

Initial U.S. Approval: 1997

INDICATIONS AND USAGE

Nitroglycerin Lingual Spray is a nitrate vasodilator indicated for acute relief of an attack or prophylaxis of angina pectoris due to coronary artery disease (1). (1)

DOSAGE AND ADMINISTRATION

- At the onset of an attack, administer onto or under the tongue. Repeat every 5 minutes as needed (2.1).
- Do up to three metered sprays within a 15-minute period. If chest pain persists, advise prompt medical attention (2.1).
- May be used prophylactically 5 to 10 minutes prior to engaging in activities that might precipitate an acute attack (2.1).

DOSAGE FORMS AND STRENGTHS

Lingual spray, 400 mcg per spray, available in 60 or 200 metered sprays per container (3). (3)

CONTRAINDICATIONS

- Use of phosphodiesterase type 5 (PDE-5) inhibitors, such as avanafil, sildenafil, tadalafil, or vardenafil, or soluble guanylate cyclase (sGC) stimulator (riociguat). (4.1, 7.1)
- Severe anemia. (4.2)
- Increased intracranial pressure. (4.3)
- Hypersensitivity to Nitroglycerin Lingual Spray or to other nitrates or nitrites or any excipient. (4.4)

WARNINGS AND PRECAUTIONS

Tolerance: Excessive use may lead to tolerance. (5.1) (5)

Hypotension: Severe hypotension may occur. (5.2) (5)

ADVERSE REACTIONS

Most common adverse reactions occurring at a frequency greater than 2% are headache, dizziness and paresthesia (6).

To report SUSPECTED ADVERSE REACTIONS, contact Perrigo at 1-866-634-9120 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Antihypertensives: Possible additive hypotensive effects. (7.2)
- Ergotamine: increased bioavailability of ergotamine. Avoid concomitant use (7.3)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 10/2020

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage

2.2 Priming

2.3 Administration

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

4.1 PDE-5-Inhibitors and sGC-Stimulators

4.2 Severe Anemia

4.3 Increased Intracranial Pressure

4.4 Hypersensitivity

4.5 Circulatory Failure and Shock

5 WARNINGS AND PRECAUTIONS

5.1 Tolerance

5.2 Hypotension

5.3 Hypertrophic Obstructive Cardiomyopathy

5.4 Headache

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

6.2 Postmarketing Experience

7 DRUG INTERACTIONS

7.1 PDE-5-Inhibitors and sGC-Stimulators

7.2 Antihypertensives

7.3 Ergotamine

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

10 OVERDOSAGE

10.1 Signs and Symptoms, Methemoglobinemia

10.2 Treatment of Overdosage

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Nitroglycerin Lingual Spray is indicated for acute relief of an attack or prophylaxis of

angina pectoris due to coronary artery disease.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage

Instruct the patient to administer one or two metered sprays (400 mcg of nitroglycerin per spray) at the onset of an attack onto or under the tongue. A spray may be repeated approximately every five minutes as needed. No more than three metered sprays are recommended within a 15-minute period. If the chest pain persists after a total of three sprays, advise prompt medical attention. Nitroglycerin Lingual Spray may be used prophylactically 5 to 10 minutes prior to engaging in activities that might precipitate an acute attack.

2.2 Priming

The pump must be primed prior to the first use. Each metered spray of Nitroglycerin Lingual Spray delivers 48 mg of solution containing 400 mcg of nitroglycerin after an initial priming of five sprays. It will remain adequately primed for 6 weeks. If the product is not used within 6 weeks it can be adequately re-primed with one spray. If the product is not used within 3 months it can be adequately re-primed with up to five sprays. There are 60 or 200 metered sprays per bottle. The total number of available doses is dependent, however, on the number of sprays per use (1 or 2 sprays), and the frequency of priming.

2.3 Administration

Instruct patients that during administration, the patient should rest, ideally in the sitting position. Hold the container vertically with the valve head uppermost and the spray orifice as close to the mouth as possible. Spray the dose preferably onto or under the tongue by pressing the grooved-button firmly and the mouth closed immediately after each dose. **THE SPRAY SHOULD NOT BE INHALED.** The medication should not be expectorated or the mouth rinsed for 5 to 10 minutes following administration. Instruct patients to familiarize themselves with the position of the spray orifice, which can be identified by the finger rest on top of the valve, in order to facilitate orientation for administration at night [*see Patient Information (17)*].

The amount of liquid remaining in the container should be checked periodically. The transparent container can be used for continuous monitoring of the consumption. With the container upright and level, check to be sure the end of the center tube extends below the level of the liquid. Once fluid falls below the level of the center tube, remaining sprays will not deliver intended dose.

3 DOSAGE FORMS AND STRENGTHS

Lingual spray, 400 mcg per spray available in either 60 or 200 metered sprays per container.

4 CONTRAINDICATIONS

4.1 PDE-5-Inhibitors and sGC-Stimulators

Do not use Nitroglycerin Lingual Spray in patients who are taking PDE-5-Inhibitors, such as avanafil, sildenafil, tadalafil, or vardenafil. Concomitant use can cause severe hypotension, syncope, or myocardial ischemia [*see Drug Interactions (7.1)*].

Do not use Nitroglycerin Lingual Spray in patients who are taking soluble guanylate cyclase (sGC) stimulators, such as riociguat. Concomitant use can cause hypotension.

4.2 Severe Anemia

Nitroglycerin Lingual Spray is contraindicated in patients with severe anemia (large doses of nitroglycerin may cause oxidation of hemoglobin to methemoglobin and could exacerbate anemia).

4.3 Increased Intracranial Pressure

Nitroglycerin Lingual Spray may precipitate or aggravate increased intracranial pressure and thus should not be used in patients with possible increased intracranial pressure (e. g. cerebral hemorrhage or traumatic brain injury).

4.4 Hypersensitivity

Nitroglycerin Lingual Spray is contraindicated in patients who are allergic to nitroglycerin, other nitrates or nitrites or any excipient.

4.5 Circulatory Failure and Shock

Nitroglycerin Lingual Spray is contraindicated in patients with acute circulatory failure or shock.

5 WARNINGS AND PRECAUTIONS

5.1 Tolerance

Excessive use may lead to the development of tolerance. Only the smallest number of doses required for effective relief of the acute angina attack should be used [*see Dosage and Administration (2.1)*].

5.2 Hypotension

Severe hypotension, particularly with upright posture, may occur even with small doses of nitroglycerin particularly in patients with constrictive pericarditis, aortic or mitral stenosis, patients who may be volume-depleted, or are already hypotensive.

Hypotension induced by nitroglycerin may be accompanied by paradoxical bradycardia and increased angina pectoris. Symptoms of severe hypotension (nausea, vomiting, weakness, pallor, perspiration and collapse/syncope) may occur even with therapeutic doses.

5.3 Hypertrophic Obstructive Cardiomyopathy

Nitrate therapy may aggravate the angina caused by hypertrophic obstructive cardiomyopathy.

5.4 Headache

Nitroglycerin produces dose-related headaches, especially at the start of nitroglycerin therapy, which may be severe and persistent but usually subside with continued use.

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

Adverse reactions occurring at a frequency greater than 2% and greater than placebo included: headache, dizziness, and paresthesia.

6.2 Postmarketing Experience

The following adverse reactions have been identified during post-approval use of nitroglycerin lingual spray and other nitroglycerin drugs. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to estimate their frequency reliably or establish a causal relationship to drug exposure.

Neurologic: weakness, drowsiness

Dermatologic: cutaneous vasodilation, flushing, drug rash, exfoliative dermatitis

Gastrointestinal: nausea, vomiting

Respiratory: transient hypoxemia

Cardiovascular: tachycardia

7 DRUG INTERACTIONS

7.1 PDE-5-Inhibitors and sGC-Stimulators

Nitroglycerin Lingual Spray is contraindicated in patients who are using a selective inhibitor of cyclic guanosine monophosphate (cGMP)-specific phosphodiesterase type 5 (PDE-5). PDE-5-Inhibitors such as avanafil, sildenafil, vardenafil, and tadalafil have been shown to potentiate the hypotensive effects of organic nitrates.

Nitroglycerin Lingual Spray is contraindicated in patients who are taking soluble guanylate cyclase (sGC) stimulators. Concomitant use can cause hypotension.

The time course and dose dependence of these interactions have not been studied, and use within a few days of one another is not recommended. Appropriate supportive care for the severe hypotension has not been studied, but it seems reasonable to treat this as a nitrate overdose, with elevation of the extremities and with central volume expansion.

7.2 Antihypertensives

Patients receiving antihypertensive drugs, beta-adrenergic blockers, and nitrates should

be observed for possible additive hypotensive effects. Marked orthostatic hypotension has been reported when calcium channel blockers and organic nitrates were used concomitantly.

Beta-adrenergic blockers blunt the reflex tachycardia produced by nitroglycerin without preventing its hypotensive effects. If beta-blockers are used with nitroglycerin in patients with angina pectoris, additional hypotensive effects may occur.

7.3 Ergotamine

Oral administration of nitroglycerin markedly decreases the first-pass metabolism of dihydroergotamine and subsequently increases its oral bioavailability.

Ergotamine is known to precipitate angina pectoris. Therefore, patients receiving sublingual nitroglycerin should avoid ergotamine and related drugs or be monitored for symptoms of ergotism if this is not possible.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk summary

Limited published data on the use of nitroglycerin are insufficient to determine a drug associated risk of major birth defects or miscarriage. In animal reproduction studies, there were no adverse developmental effects when nitroglycerin was administered intravenously to rabbits or intraperitoneally to rats during organogenesis at doses greater than 64-times the human dose [see *Data*].

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 - 4% and 15 - 20%, respectively.

Data

Animal Data

No embryotoxic or postnatal development effects were observed with transdermal application in pregnant rabbits and rats at doses up to 240 mg/kg/day for 13 days, at intraperitoneal doses in pregnant rats up to 20 mg/kg/day for 11 days, and at intravenous doses in pregnant rabbits up to 4 mg/kg/day for 13 days.

8.2 Lactation

Risk summary

Sublingual nitroglycerin has not been studied in lactating women. It is not known if nitroglycerin is present in human milk or if nitroglycerin has effects on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for nitroglycerin and any potential adverse effects on the breastfed child from nitroglycerin or from the underlying maternal condition.

8.4 Pediatric Use

Safety and effectiveness of nitroglycerin in pediatric patients have not been established.

8.5 Geriatric Use

Clinical studies did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between elderly (greater than or equal to 65 years) and younger (less than 65 years) patients. In general, dose selection for an elderly patient should start at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

10 OVERDOSAGE

10.1 Signs and Symptoms, Methemoglobinemia

Nitrate overdosage may result in: severe hypotension, persistent throbbing headache, vertigo, palpitation, visual disturbance, flushing and perspiring skin (later becoming cold and cyanotic), nausea and vomiting (possibly with colic and even bloody diarrhea), syncope (especially in the upright posture), methemoglobinemia with cyanosis and anorexia, initial hyperpnea, dyspnea and slow breathing, slow pulse (dicrotic and intermittent), heart block, increased intracranial pressure with cerebral symptoms of confusion and moderate fever, paralysis and coma followed by clonic convulsions, and possibly death due to circulatory collapse.

Case reports of clinically significant methemoglobinemia are rare at conventional doses of organic nitrates. The formation of methemoglobin is dose-related and in the case of genetic abnormalities of hemoglobin that favor methemoglobin formation, even conventional doses of organic nitrates could produce harmful concentrations of methemoglobin.

10.2 Treatment of Overdosage

As hypotension associated with nitroglycerin overdose is the result of venodilatation and arterial hypovolemia, prudent therapy in this situation should be directed toward increase in central fluid volume. No specific antagonist to the vasodilator effects of nitroglycerin is known. Keep the patient recumbent in a shock position and comfortably warm. Passive movement of the extremities may aid venous return. Intravenous infusion of normal saline or similar fluid may also be necessary. Administer oxygen and artificial ventilation, if necessary. If methemoglobinemia is present, administration of methylene blue (1% solution), 1-2 mg per kilogram of body weight intravenously, may be required unless the patient is known to have G-6-PD deficiency. If an excessive quantity of Nitroglycerin Lingual Spray has been recently swallowed gastric lavage may be of use.

As epinephrine is ineffective in reversing the severe hypotensive events associated with overdosage, it is not recommended for resuscitation.

11 DESCRIPTION

Nitroglycerin, an organic nitrate, is a vasodilator which has effects on both arteries and veins.

The chemical name for nitroglycerin is 1,2,3-propanetriol trinitrate (C₃H₅N₃O₉).

The compound has a molecular weight of 227.09. The chemical structure is:

CH₂-ONO₂

|

CH-ONO₂

|

CH₂-ONO₂

Nitroglycerin Lingual Spray (nitroglycerin lingual spray 400 mcg) is a metered dose spray containing nitroglycerin. This product delivers nitroglycerin (400 mcg per spray, 60 or 200 metered sprays) in the form of spray droplets onto or under the tongue. Inactive ingredients: medium-chain triglycerides, dehydrated alcohol, medium-chain partial glycerides, peppermint oil.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Nitroglycerin forms free radical nitric oxide (NO), which activates guanylate cyclase, resulting in an increase of guanosine 3',5'-monophosphate (cyclic GMP) in smooth muscle and other tissues. This eventually leads to dephosphorylation of myosin light chains, which regulates the contractile state in smooth muscle and results in vasodilatation.

12.2 Pharmacodynamics

The principal pharmacological action of nitroglycerin is relaxation of vascular smooth muscle. Although venous effects predominate, nitroglycerin produces, in a dose-related manner, dilation of both arterial and venous beds. Dilation of the postcapillary vessels, including large veins, promotes peripheral pooling of blood, decreases venous return to the heart, and reduces left ventricular end-diastolic pressure (preload). Nitroglycerin also produces arteriolar relaxation, thereby reducing peripheral vascular resistance and arterial pressure (after load), and dilates large epicardial coronary arteries; however, the extent to which this latter effect contributes to the relief of exertional angina is unclear.

Therapeutic doses of nitroglycerin may reduce systolic, diastolic and mean arterial blood pressure. Effective coronary perfusion pressure is usually maintained, but can be compromised if blood pressure falls excessively or increased heart rate decreases diastolic filling time.

Elevated central venous and pulmonary capillary wedge pressures, and pulmonary and systemic vascular resistance are also reduced by nitroglycerin therapy. Heart rate is usually slightly increased, presumably a reflex response to the fall in blood pressure. Cardiac index may be increased, decreased, or unchanged. Myocardial oxygen consumption or demand (as measured by the pressure-rate product, tension-time index, and stroke-work index) is decreased and a more favorable supply-demand ratio can be achieved. Patients with elevated left ventricular filling pressure and increased systemic vascular resistance in association with a depressed cardiac index are likely to

experience an improvement in cardiac index. In contrast, when filling pressures and cardiac index are normal, cardiac index may be slightly reduced following nitroglycerin administration.

12.3 Pharmacokinetics

A liver reductase enzyme is of primary importance in the metabolism of nitroglycerin to glycerol di- and mononitrate metabolites and ultimately to glycerol and organic nitrate. Known sites of extrahepatic metabolism include red blood cells and vascular walls. In addition to nitroglycerin, 2 major metabolites, 1,2- and 1,3-dinitroglycerin are found in plasma. The mean elimination half-life of both 1,2- and 1,3-dinitroglycerin is about 40 minutes. The 1,2- and 1,3-dinitroglycerin metabolites have been reported to possess some pharmacological activity, whereas the glycerol mononitrate metabolites of nitroglycerin are essentially inactive. Higher plasma concentrations of the dinitro metabolites, with their nearly 8-fold longer elimination half-lives, may contribute significantly to the duration of pharmacologic effect.

In a pharmacokinetic study when a single 0.8 mg dose of nitroglycerin lingual spray was administered to healthy volunteers ($n = 24$), the mean C_{max} and t_{max} were 1.041 pg/mL and 7.5 minutes, respectively. Additionally, in these subjects the mean area under the curve (AUC) was 12.769 pg/mL * min.

The volume of distribution of nitroglycerin following intravenous administration is 3.3 L/kg.

Drug interactions

Aspirin: Coadministration of nitroglycerin with high dose aspirin (1000 mg) results in increased exposure to nitroglycerin. The vasodilatory and hemodynamic effects of nitroglycerin may be enhanced by concomitant administration of nitroglycerin with high dose aspirin.

Tissue-type plasminogen activator (t-PA): Concomitant administration of t-PA and intravenous nitroglycerin has been shown to reduce plasma levels of t-PA and its thrombolytic effect.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Animal carcinogenesis studies with sublingual nitroglycerin have not been performed.

Rats receiving up to 434 mg/kg/day of dietary nitroglycerin for 2 years developed dose-related fibrotic and neoplastic changes in liver, including carcinomas, and interstitial cell tumors in testes. At high dose, the incidences of hepatocellular carcinomas in both sexes were 52% vs. 0% in controls, and incidences of testicular tumors were 52% vs. 8% in controls. Lifetime dietary administration of up to 1058 mg/kg/day of nitroglycerin was not tumorigenic in mice.

Nitroglycerin was weakly mutagenic in Ames tests performed in two different laboratories. There was no evidence of mutagenicity in an *in vivo* dominant lethal assay with male rats treated with doses up to about 363 mg/kg/day, p.o., or in *in vitro* cytogenic tests in rat and dog tissues and for chromosomal aberration in Chinese

hamster ovary cells.

In a three-generation reproduction study, rats received dietary nitroglycerin at doses up to about 434 mg/kg/day for six months prior to mating of the F₀ generation with treatment continuing through successive F₁ and F₂ generations. The high dose was associated with decreased feed intake and body weight gain in both sexes at all matings. No specific effect on the fertility of the F₀ generation was seen. Infertility noted in subsequent generations, however, was attributed to increased interstitial cell tissue and aspermatogenesis in the high-dose males. In this three-generation study there was no clear evidence of teratogenicity.

14 CLINICAL STUDIES

In a randomized, double-blind single-dose, 5-period cross-over study in 51 patients with exertional angina pectoris significant dose-related increases in exercise tolerance, time to onset of angina and ST-segment depression were seen following doses of 0.2, 0.4, 0.8 and 1.6 mg of nitroglycerin delivered by metered pumpspray as compared to placebo.

The drug showed a profile of mild to moderate adverse events.

16 HOW SUPPLIED/STORAGE AND HANDLING

Each box of Nitroglycerin Lingual Spray contains one glass bottle coated with orange transparent plastic which assists in containing the glass and medication should the bottle be shattered. Each bottle contains 4.9 g or 12 g (Net Contents) of nitroglycerin lingual spray which will deliver 60 or 200 metered sprays containing 400 mcg of nitroglycerin per spray after priming.

Nitroglycerin Lingual Spray is available as:

- 60-dose (4.9 g) single bottle NDC 45802-**210**-01
- 200-dose (12 g) single bottle NDC 45802-**210**-02

Store at 20-25°C (68-77°F) [see USP Controlled Room Temperature].

Note: Nitroglycerin Lingual Spray contains 20% alcohol. Do not forcefully open or burn container after use. Do not spray toward flames.

Rx Only

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Instructions for Use).

Made in Israel

Manufactured by Perrigo

Yeruham, Israel

Distributed By

Perrigo®

Instructions for Use

Nitroglycerin Lingual Spray


Read this Instructions for Use before you start using Nitroglycerin Lingual Spray and each time you get your prescription refilled. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or your treatment.

Important information:

- **Nitroglycerin Lingual Spray is for use onto or under the tongue.** Do not inhale Nitroglycerin Lingual Spray.
- A dose of Nitroglycerin Lingual Spray may be either 1 or 2 sprays. Follow your healthcare provider's instructions about how many sprays you should use for each dose. Doses should be separated by approximately 5 minutes.
- **You should not use more than 3 sprays of Nitroglycerin Lingual Spray within 15 minutes.**
- **Get emergency medical help right away if you still have chest pain after using a total of 3 sprays of Nitroglycerin Lingual Spray.**

Nitroglycerin Lingual Spray parts:

FIGURE A



How should I use Nitroglycerin Lingual Spray?

- It is best to use Nitroglycerin Lingual Spray while you are resting and in a sitting position.
- **Do not** shake the Nitroglycerin Lingual Spray container.


Priming Nitroglycerin Lingual Spray:

Before you use Nitroglycerin Lingual Spray for the first time, you must prime it. To prime your Nitroglycerin Lingual Spray, follow the steps below:

Step 1. Remove the plastic cap from the container.

(See Figure B)


FIGURE B



Step 2. Hold the container upright and facing away from yourself and others. Press down on the top of the grooved button 5 times.

(See Figure C)

FIGURE C



- Your Nitroglycerin Lingual Spray is now primed. You are ready to give your first dose.
- If you do not use your Nitroglycerin Lingual Spray within 6 weeks, you will need to prime it again by pressing down on the top of the grooved button 1 time.
- If you do not use your Nitroglycerin Lingual Spray within 3 months, you will need to re-prime it by pressing down on the top of the grooved button up to 5 times.

Giving a dose of Nitroglycerin Lingual Spray:

Step 3. Hold your Nitroglycerin Lingual Spray container upright with your index finger on top of the grooved button.

Step 4. Open your mouth and bring the Nitroglycerin Lingual Spray container as close to your mouth as possible.

Step 5. Press down on the top of the grooved button firmly with your index finger to release the spray onto or under your tongue. (See Figure D). The grooved button can help you make sure the canister is turned to the correct position if you are administering the spray in the dark.

FIGURE D



Step 6. Release the grooved button and close your mouth right away. Avoid swallowing right after using Nitroglycerin Lingual Spray. Do not spit out Nitroglycerin Lingual Spray or rinse your mouth for 5 to 10 minutes after using Nitroglycerin Lingual Spray.

Step 7. If a second dose of Nitroglycerin Lingual Spray is needed, repeat Steps 3 through 6 above.

Step 8. Replace the plastic cap.

Check the level of the fluid in your Nitroglycerin Lingual Spray container regularly.

- Check the container in an upright position.
- The end of the center tube should be covered by the fluid in the Nitroglycerin Lingual Spray container. If the level of the fluid falls below the end of the center tube, sprays will not provide enough Nitroglycerin Lingual Spray.
- Replace your Nitroglycerin Lingual Spray container before the fluid level falls below the end of the center tube.

How should I store Nitroglycerin Lingual Spray?

- Store Nitroglycerin Lingual Spray at room temperature 68°F - 77°F (20°C - 25°C) [see USP Controlled Room Temperature].
- Do not forcefully open or burn the Nitroglycerin Lingual Spray container after use.
- Do not spray Nitroglycerin Lingual Spray toward flames.

Keep Nitroglycerin Lingual Spray and all medicines out of the reach of children.

This Instructions for Use has been approved by the U.S. Food and Drug Administration.

Made in Israel

Manufactured by Perrigo

Yeruham, Israel

Distributed By

Perrigo®

Allegan, MI 49010 • www.perrigo.com

Rev 09-19

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL- 12 g Carton

Rx Only

NDC 45802-210-02

Nitroglycerin Lingual Spray

400 mcg per Spray


200 Metered Sprays

DO NOT SHAKE

HOLD CONTAINER UPRIGHT

PRIME BEFORE USE

12 g Net Contents



The following image is a placeholder representing the product identifier that is either affixed or imprinted on the drug package label during the packaging operation.

S/N [insert product's serial number]
 Lot [insert product's lot number]
 Exp [insert product's expiration date]

NITROGLYCERIN LINGUAL

nitroglycerin spray

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:45802-210
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NITROGLYCERIN (UNII: G59M7S0WS3) (NITROGLYCERIN - UNII:G59M7S0WS3)	NITROGLYCERIN	400 ug

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
PEPPERMINT OIL (UNII: AV092KU4JH)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:45802-210-01	1 in 1 CARTON	09/20/2013	
1		60 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:45802-210-02	1 in 1 CARTON	09/20/2013	
2		200 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091496	09/20/2013	

