

DIGESTIVE RELIEF REGULAR STRENGTH- bismuth subsalicylate liquid

TARGET Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient (in each 30 mL)

Bismuth subsalicylate 525 mg

Purpose

Upset stomach reliever/Antidiarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
 - heartburn
 - indigestion
 - nausea
 - gas
 - fullness
 - belching

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- mL = milliliter
- shake well before using
- measure only with dosing cup provided. do not use any other dosing device.
- keep dosing cup with product
- adults and children 12 years and over
 - 30 mL (1 dose) every 1/2 or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
 - 30 mL (1 dose) every 1/2 as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
 - do not exceed 8 doses (240 mL) in 24 hours
 - use until diarrhea stops but not more than 2 days
- children under 12 years of age: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- **each 30 mL contains:** potassium 8 mg sodium 6 mg
- **each 30 mL contains:** salicylate 206 mg
- low sodium

- keep tightly closed
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

Inactive ingredients

benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

Questions or comments?

Call **1-800-910-6874**

Principal Display Panel

Compare to the active ingredient in Pepto-Bismol®*

5-symptom

digestive relief

Bismuth subsalicylate 525 mg per 30 mL

upset stomach reliever/antidiarrheal

Soothing relief of heartburn, indigestion, nausea,

upset stomach and diarrhea

FL OZ (mL)

GLUTEN FREE

*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

TAMPER EVIDENT; DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

Dist. by Target Corp., Mpls., MN 55403

TM ©2020 Target Brands, Inc.

Package Label

Drug Facts (continued)

Directions ■ mL = milliliter ■ shake well before using
 ■ measure only with dosing cup provided. Do not use any other dosing device.
 ■ keep dosing cup with product
 ■ adults and children 12 years and over
 ■ 30 mL (1 dose) every ½ hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
 ■ 30 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
 ■ do not exceed 8 doses (240 mL) in 24 hours
 ■ use until diarrhea stops but not more than 2 days
 ■ children under 12 years of age: ask a doctor
 ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information ■ each 30 mL contains: potassium 8 mg, sodium 9 mg
 ■ each 30 mL contains: salicylate 206 mg
 ■ low sodium ■ keep tightly closed ■ protect from freezing
 ■ avoid excessive heat (over 104°F or 40°C)

Inactive ingredients benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

Questions or comments? Call 1-800-910-6874

*This product is not manufactured or distributed by The Procter & Gamble Company.
 Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.
 Dist. by Target Corp., Mpls., MN 55403
 TM & ©2020 Target Brands, Inc.

GLUTEN FREE

245 05 0365 R03 C-000703-01-111



3 49580 37218 4



PLD-84928 LB007728

Compare to the active ingredient
 in **Pepto-Bismol®***

**5-symptom
 digestive relief**

bismuth subsalicylate 525 mg per 30 mL
 upset stomach reliever/antidiarrheal

soothing relief of heartburn, indigestion, nausea,
 upset stomach and diarrhea



16 FL OZ (1 PT) 473 mL

TARGET Digestive Relief

NDC 11673-724-16

Drug Facts

Active ingredient (in each 30 mL dose) Bismuth subsalicylate 525 mg.....Upset stomach reliever/Antidiarrheal **Purpose**

Uses relieves ■ diarrhea ■ travelers' diarrhea
 ■ upset stomach due to overindulgence in food and drink, including
 ■ heartburn ■ gas ■ indigestion
 ■ nausea ■ fullness ■ belching

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Contains salicylate. Do not take if you are:
 ■ allergic to salicylates (including aspirin) ■ taking other salicylate products

Do not use if you have
 ■ an ulcer ■ a bleeding problem ■ bloody or black stool

Ask a doctor before use if you have ■ fever ■ mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for
 ■ diabetes ■ gout ■ arthritis ■ anticoagulation (thinning the blood)

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if
 ■ symptoms get worse or last more than 2 days
 ■ ringing in the ears or loss of hearing occurs
 ■ diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
 In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL
 AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

DIGESTIVE RELIEF REGULAR STRENGTH

bismuth subsalicylate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-724
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
BENZOIC ACID (UNII: 8SKN0B0MIM)	
D&C RED NO. 22 (UNII: 1678RKX8RT)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0KO0R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:11673-724-16	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/30/2019	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC monograph final	part335		09/30/2019	

Labeler - TARGET Corporation (006961700)

Revised: 11/2022

TARGET Corporation