

MEDI FIRST PLUS PAIN ACHE- aspirin,caffeine tablet, film coated
MEDI FIRST PAIN ACHE- aspirin,caffeine tablet, film coated
Unifirst First Aid Corporation

Medi-First/Medi-First Plus Pain & Ache

Drug Facts

Active ingredients (in each caplet)

Aspirin (NSAID)* 500 mg

Caffeine 32.5 mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Pain reliever aid

Uses

for the temporary relief of

- backache pain
- muscle aches and pains
- minor aches and pains of arthritis

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert:

Aspirin may cause a severe allergic reaction which may include:

- hives
- shock
- facial swelling
- asthma (wheezing)

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- have 3 or more alcohol drinks every day while using this product

- have had stomach ulcers or bleeding problems

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer.

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you

- you have a history of stomach problems, such as heartburn

- you have asthma

- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

- you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout

- diabetes

- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.

- you experience any of the following signs of stomach bleeding: ■ vomit blood ■ have bloody or black stools ■ feel faint ■ have stomach pain that does not get better

- ringing in the ears or a loss of hearing occurs

- pain gets worse or lasts more than 10 days

- fever gets worse or lasts more than 3 days

- redness or swelling is present

- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **drink a full glass of water with each dose**

Adults and children (12 years and over): take 2 caplets every 6 hours, not to exceed 8 caplets in 24 hours

Children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- tamper evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

corn starch, hypromellose, maltodextrin, microcrystalline cellulose, polyethylene glycol, sodium lauryl sulfate

Questions or comments? 1-800-634-7680

Medi-First®

Pain & Ache

Pain Reliever • Aspirin (NSAID) 500 mg

Pain Reliever Aid • Caffeine 32.5 mg

Fast Acting Advanced Formula

Pull to Open

This Package is for Households Without Young Children.

Tamper Evident Unit Dose Packets

80 Caplets

(40 x 2)

MEDI-FIRST® Pain & Ache
Fast Acting Advanced Formula
✓ Pain Reliever - Aspirin (NSAID) 500 mg
✓ Pain Reliever Aid - Caffeine 32.5 mg
THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.
Tamper Evident Unit Dose Packets

Drug Facts
Active ingredients (in each caplet) Purpose
Aspirin (NSAID) 500 mg Pain reliever
Caffeine 32.5 mg Pain reliever aid
*nonsteroidal anti-inflammatory drug

Uses
For the temporary relief of
■ headache pain
■ muscle aches and pains,
■ minor aches and pains of arthritis

Warnings
Roye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Roye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:
■ hives
■ shock
■ facial swelling
■ asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
■ take more or for a longer time than directed
■ take a blood thinning (anticoagulant) or steroid drug
■ are age 60 or older
■ have 3 or more alcohol drinks every day while using this product
■ have had stomach ulcers or bleeding problems
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use
■ if you are allergic to aspirin or any other pain reliever/fever reducer.
■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if
■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn
■ you have asthma
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
■ you are taking a diuretic (continued on opposite panel)

Drug Facts (continued)
Ask a doctor or pharmacist before use if you are taking a prescription drug for
■ gout
■ diabetes
■ arthritis

Stop use and ask a doctor if
■ an allergic reaction occurs. Seek medical help right away.
■ you experience any of the following signs of stomach bleeding:
■ vomit blood
■ have bloody or black stools
■ feel faint
■ have stomach pain that does not get better
■ ringing in the ears or a loss of hearing occurs
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ new symptoms occur
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions
■ drink a full glass of water with each dose
Adults and children (12 years and over): Take 2 caplets every 6 hours, not to exceed 8 caplets in 24 hours.
Children under 12 years: do not use unless directed by a doctor.

Other information
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
■ tamper evident sealed packets
■ do not use any opened or torn packets

Inactive ingredients
corn starch, hypromellose, maltodextrin, microcrystalline cellulose, polyethylene glycol, sodium lauryl sulfate

Questions or comments? 1-800-634-7680

MEDI-FIRST® Pain & Ache
Aspirin 500 mg
Caffeine 32.5 mg

8210800X
Rev. 8/15/22
Mfg. 24

3 47682 82180 7

Retain carton for complete product information

Medi-First® Plus

Pain & Ache

Pain Reliever • Aspirin (NSAID) 500 mg

Pain Reliever Aid • Caffeine 32.5 mg

Extra Strength Pain Relief

Fast Acting Advanced Formula

Pull to Open

This Package is for Households Without Young Children.

Tamper Evident Unit Dose Packets

80 Caplets

(40 x 2's)



MEDI FIRST PLUS PAIN ACHE

aspirin, caffeine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-921
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	500 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	32.5 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3S)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	

MALTODEXTRIN (UNII: 7CVR7L4A2D)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	

Product Characteristics

Color	white	Score	no score
Shape	CAPSULE (Caplet)	Size	17mm
Flavor		Imprint Code	44;413
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-921-80	40 in 1 BOX	12/12/2022	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-921-36	75 in 1 BOX	12/12/2022	
2		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	09/26/2022	

MEDI FIRST PAIN ACHE

aspirin,caffeine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-821
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	500 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	32.5 mg

Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	

Product Characteristics

Color	white	Score	no score
Shape	CAPSULE (Caplet)	Size	17mm
Flavor		Imprint Code	44;413
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-821-80	40 in 1 BOX	09/26/2022	
1	NDC:47682-821-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-821-36	75 in 1 BOX	12/12/2022	
2	NDC:47682-821-99	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	09/26/2022	

Labeler - Unifirst First Aid Corporation (832947092)

Revised: 1/2024

Unifirst First Aid Corporation