

THOMPSON XS PAIN ACHE- aspirin, caffeine tablet, film coated

Thompson Safety

Thompson XS Pain Relief

Drug Facts

Active ingredients (in each caplet)

Aspirin 500 mg (NSAID)*

Caffeine 32.5 mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Pain reliever aid

Uses

for the temporary relief of

- backache pain
- muscle aches and pains
- minor aches and pains of arthritis

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert:

Aspirin may cause a severe allergic reaction which may include:

- hives
- shock
- facial swelling
- asthma (wheezing)

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- have 3 or more alcohol drinks every day while using this product

- have had stomach ulcers or bleeding problems

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer.

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you

- you have a history of stomach problems, such as heartburn

- you have asthma

- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

- you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout

- diabetes

- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.

- you experience any of the following signs of stomach bleeding: ■ vomit blood ■ have bloody or black stools ■ feel faint ■ have stomach pain that does not get better

- ringing in the ears or a loss of hearing occurs

- pain gets worse or lasts more than 10 days

- fever gets worse or lasts more than 3 days

- redness or swelling is present

- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **drink a full glass of water with each dose**

Adults and children (12 years and over): take 2 caplets every 6 hours, not to exceed 8 caplets in 24 hours

Children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- tamper evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

corn starch, hypromellose, maltodextrin, microcrystalline cellulose, polyethylene glycol, sodium lauryl sulfate

Questions or comments? 877.506.4291

Thompson

Extra Strength Pain & Ache Relief

Pain Relief

Aspirin (NSAID) 500 mg

Pain Reliever Aid

Caffeine 32.5 mg

Safety Works | Business Thrives™

Fast Acting Advanced Formula

Pull to Open

This Package is for Households Without Young Children.

80 Caplets

Tamper Evident Sealed Packets

EXTRA STRENGTH PAIN & ACHE
FAST ACTING ADVANCED FORMULA
THOMPSON
SAFETY WORKS | BUSINESS THRIVES
PAIN RELIEF
ASPIRIN (NSAID) 500 MG
PAIN RELIEVER AID
CAFFEINE 32.5 MG
THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.
80 Caplets
TAMPER EVIDENT SEALED PACKETS

Drug Facts
Active ingredients (in each caplet) Purpose
Aspirin (NSAID) 500 mg Pain reliever
Caffeine 32.5 mg Pain reliever and
nonsteroidal anti-inflammatory drug
Uses
For the temporary relief of
• backache pain
• muscle aches and pains
• minor aches and pains of arthritis
Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction which may include:
• hives
• itches
• shock
• facial swelling
• asthma (wheezing)
Do not use
• if you are allergic to aspirin or any other pain reliever/fever reducer.
• if you have ever had an allergic reaction to this product or any of its ingredients.
Ask a doctor before use if
• stomach bleeding warning applies to you
• you have a history of stomach problems, such as heartburn
• you have asthma
• you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
• you are taking a diuretic (continued on opposite panel)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
• take more or for a longer time than directed
• take a blood thinner (anticoagulant) or steroid drug
• are age 60 or older
• have 3 or more alcohol drinks every day while using this product
• have had stomach ulcers or bleeding problems
• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
Stop use and ask a doctor if
• an allergic reaction occurs. Seek medical help right away.
• you experience any of the following signs of stomach bleeding:
• vomit blood
• have bloody or black stools
• feel faint
• have stomach pain that does not get better
• ringing in the ears or a loss of hearing occurs
• pain gets worse or lasts more than 10 days
• fever gets worse or lasts more than 3 days
• redness or swelling is present
• new symptoms occur
These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 30 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.
Directions
• drink a full glass of water with each dose
Adults and children (12 years and over) take 2 caplets every 6 hours, not to exceed 8 caplets in 24 hours
Children under 12 years: do not use unless directed by a doctor
Other information
• store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
• tamper evident sealed packets
• do not use any opened or torn packets
Inactive ingredients
corn starch, hypromellose, maltodextrin, microcrystalline cellulose, polyethylene glycol, sodium lauryl sulfate
Questions or comments? 877.506.4291
C592180
Rev. 8/15/22
Mfg. 24

EXTRA STRENGTH PAIN & ACHE
FAST ACTING ADVANCED FORMULA
THOMPSON
SAFETY WORKS | BUSINESS THRIVES
Pull to Open
THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.
PAIN RELIEF
ASPIRIN (NSAID) 500 MG
PAIN RELIEVER AID
CAFFEINE 32.5 MG
80 Caplets
TAMPER EVIDENT SEALED PACKETS

Drug Facts (continued)
Ask a doctor or pharmacist before use if you are taking a prescription drug for
• gout
• diabetes
• arthritis
Drug Facts (continued)
Retain carton for complete product information

Drug Facts (continued)
Retain carton for complete product information

0 73408 92180 2

Reorder C592180
Distributed by
Thompson Safety
Houston, TX 77073
877.506.4291
www.thompson-safety.com

THOMPSON XS PAIN ACHE

aspirin, caffeine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73408-821
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	32.5 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	500 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

Product Characteristics

Color	white	Score	no score
Shape	CAPSULE (Caplet)	Size	17mm
Flavor		Imprint Code	44;413
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73408-821-80	40 in 1 BOX	09/26/2022	
1		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	09/26/2022	

Labeler - Thompson Safety (080998015)

