

PREVNAR 13- pneumococcal 13-valent conjugate vaccine injection, suspension

A-S Medication Solutions

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use PREVNAR 13 safely and effectively. See full prescribing information for PREVNAR 13.

**PREVNAR 13 (Pneumococcal 13-valent Conjugate Vaccine [Diphtheria CRM₁₉₇ Protein])
Suspension for intramuscular injection
Initial U.S. Approval: 2010**

RECENT MAJOR CHANGES

Indications and Usage (1.3)	7/2016
Vaccination Schedule for Children Previously Vaccinated With Prevnar Pneumococcal 7-valent Conjugate Vaccine (Diphtheria CRM ₁₉₇ Protein) (2.5) Removal	3/2017
Vaccination Schedule for Adults 18 Years of Age and Older (2.6)	7/2016
Contraindications (4)	7/2016

INDICATIONS AND USAGE

In children 6 weeks through 5 years of age (prior to the 6th birthday), Prevnar 13 is indicated for:

- active immunization for the prevention of invasive disease caused by *Streptococcus pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F. (1.1)
- active immunization for the prevention of otitis media caused by *S. pneumoniae* serotypes 4, 6B, 9V, 14, 18C, 19F, and 23F. No otitis media efficacy data are available for serotypes 1, 3, 5, 6A, 7F, and 19A. (1.1)

In children 6 years through 17 years of age (prior to the 18th birthday), Prevnar 13 is indicated for:

- active immunization for the prevention of invasive disease caused by *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F. (1.2)

In adults 18 years of age and older, Prevnar 13 is indicated for:

- active immunization for the prevention of pneumonia and invasive disease caused by *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F. (1.3)

Limitations of Prevnar 13 Use and Effectiveness

- Prevnar 13 does not protect against disease caused by *S. pneumoniae* serotypes that are not in the vaccine. (1.4)

DOSAGE AND ADMINISTRATION

Children 6 weeks through 5 years: The four-dose immunization series consists of a 0.5 mL intramuscular injection administered at 2, 4, 6, and 12–15 months of age. (2.3)

Children 6 through 17 years of age: a single dose. (2.5)

Adults 18 years and older: a single dose. (2.6)

DOSAGE FORMS AND STRENGTHS

0.5 mL suspension for intramuscular injection, supplied in a single-dose prefilled syringe. (3)

CONTRAINDICATIONS

Severe allergic reaction (e.g., anaphylaxis) to any component of Prevnar 13 or any diphtheria toxoid-containing vaccine. (4)

WARNINGS AND PRECAUTIONS

Apnea following intramuscular vaccination has been observed in some infants born prematurely. Decisions about when to administer an intramuscular vaccine, including Prevnar 13, to infants born prematurely should be based on consideration of the individual infant's medical status, and the potential benefits and possible risks of vaccination. (5.3)

ADVERSE REACTIONS

- In infants and toddlers vaccinated at 2, 4, 6, and 12–15 months of age in US clinical trials, the most commonly reported solicited adverse reactions (>5%) were irritability (>70%), injection site tenderness (>50%), decreased appetite (>40%), decreased sleep (>40%), increased sleep (>40%), fever (>20%), injection site redness (>20%), and injection site swelling (>20%). (6.1)
- In children aged 5 through 17 years, the most commonly reported solicited adverse reactions (>5%) were injection site tenderness (>80%), injection site redness (>30%), injection site swelling (>30%), irritability (>20%), decreased appetite (>20%), increased sleep (>20%), fever (>5%), and decreased sleep (>5%). (6.1)

- In adults aged 18 years and older, the most commonly reported solicited adverse reactions (>5%) were pain at the injection site (>50%), fatigue (>30%), headache (>20%), muscle pain (>20%), joint pain (>10%), decreased appetite (>10%), injection site redness (>10%), injection site swelling (>10%), limitation of arm movement (>10%), vomiting (>5%), fever (>5%), chills (>5%), and rash (>5%). (6.2)

To report SUSPECTED ADVERSE REACTIONS, contact Wyeth Pharmaceuticals Inc. at 1-800-438-1985 or VAERS at 1-800-822-7967 or <http://vaers.hhs.gov>.

----- USE IN SPECIFIC POPULATIONS -----

Pediatric Use: Safety and effectiveness of Prevnar 13 in children below the age of 6 weeks have not been established. (8.4)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 8/2018

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

- 1.1 Children 6 Weeks Through 5 Years of Age
- 1.2 Children 6 Years Through 17 Years of Age
- 1.3 Adults 18 Years of Age and Older
- 1.4 Limitations of Prevnar 13 Use and Effectiveness

2 DOSAGE AND ADMINISTRATION

- 2.1 Preparation for Administration
- 2.2 Administration Information
- 2.3 Vaccination Schedule for Infants and Toddlers
- 2.4 Vaccination Schedule for Unvaccinated Children 7 Months Through 5 Years of Age
- 2.5 Vaccination Schedule for Children 6 Years Through 17 Years of Age
- 2.6 Vaccination Schedule for Adults 18 Years of Age and Older

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Management of Allergic Reactions
- 5.2 Altered Immunocompetence
- 5.3 Apnea in Premature Infants

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience With Prevnar 13 in Children 6 Weeks Through 17 Years of Age
- 6.2 Clinical Trials Experience With Prevnar 13 in Adults \geq 18 Years of Age
- 6.3 Post-marketing Experience With Prevnar 13 in Infants and Toddlers

7 DRUG INTERACTIONS

- 7.1 Concomitant Immunizations
- 7.2 Immunosuppressive Therapies
- 7.3 Antipyretics
- 7.4 Prior Vaccination with PPSV23

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 High Risk Populations

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

14.1 Efficacy Data

14.2 Prevnar 13 Clinical Trials in Children 6 Weeks Through 17 Years of Age

14.3 Prevnar 13 Immunogenicity Clinical Trials in Adults

14.4 Concomitant Vaccine Administration

15 REFERENCES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

1.1 Children 6 Weeks Through 5 Years of Age

In children 6 weeks through 5 years of age (prior to the 6th birthday), Prevnar 13[®] is indicated for:

- active immunization for the prevention of invasive disease caused by *Streptococcus pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F.
- active immunization for the prevention of otitis media caused by *S. pneumoniae* serotypes 4, 6B, 9V, 14, 18C, 19F, and 23F. No otitis media efficacy data are available for serotypes 1, 3, 5, 6A, 7F, and 19A.

1.2 Children 6 Years Through 17 Years of Age

In children 6 years through 17 years of age (prior to the 18th birthday), Prevnar 13 is indicated for:

- active immunization for the prevention of invasive disease caused by *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F.

1.3 Adults 18 Years of Age and Older

In adults 18 years of age and older, Prevnar 13 is indicated for:

- active immunization for the prevention of pneumonia and invasive disease caused by *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F.

1.4 Limitations of Prevnar 13 Use and Effectiveness

- Prevnar 13 does not protect against disease caused by *S. pneumoniae* serotypes that are not in the vaccine.

2 DOSAGE AND ADMINISTRATION

2.1 Preparation for Administration

Since this product is a suspension containing an adjuvant, shake vigorously immediately prior to use to obtain a homogenous, white suspension in the vaccine container. Do not use the vaccine if it cannot be resuspended. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration [see Description (11)]. This product should not be used if particulate matter or discoloration is found.

Do not mix Prevnar 13 with other vaccines/products in the same syringe.

2.2 Administration Information

For intramuscular injection only.

Each 0.5 mL dose is to be injected intramuscularly using a sterile needle attached to the supplied prefilled syringe. The preferred sites for injection are the anterolateral aspect of the thigh in infants and the deltoid muscle of the upper arm in toddlers, children and adults. The vaccine should not be injected in the gluteal area or areas where there may be a major nerve trunk and/or blood vessel.

2.3 Vaccination Schedule for Infants and Toddlers

Prevnar 13 is to be administered as a four-dose series at 2, 4, 6, and 12–15 months of age.

Table 1: Vaccination Schedule for Infants and Toddlers

Dose	Dose 1 ^{*,†}	Dose 2 [†]	Dose 3 [†]	Dose 4 [‡]
Age at Dose	2 months	4 months	6 months	12–15 months

* Dose 1 may be given as early as 6 weeks of age.

† The recommended dosing interval is 4 to 8 weeks.

‡ The fourth dose should be administered at approximately 12–15 months of age, and at least 2 months after the third dose.

2.4 Vaccination Schedule for Unvaccinated Children 7 Months Through 5 Years of Age

For children 7 months through 5 years of age who have not received Prevnar[®] or Prevnar 13, the catch-up schedule in Table 2 applies:

Table 2: Vaccination Schedule for Unvaccinated Children 7 Months of Age Through 5 Years of Age

Age at First Dose	Total Number of 0.5 mL Doses
7–11 months of age	3 [*]
12–23 months of age	2 [†]
24 months through 5 years of age (prior to the 6 th birthday)	1

* The first 2 doses at least 4 weeks apart; third dose after the one-year birthday, separated from the second dose by at least 2 months.

† Two doses at least 2 months apart.

The immune responses induced by this catch-up schedule may result in lower antibody concentrations for some serotypes, compared to antibody concentrations following 4 doses of Prevnar 13 (given at 2, 4, 6, and 12–15 months). In children 24 months through 5 years of age, lower antibody concentrations were observed for some serotypes, compared to antibody concentrations following 3 doses of Prevnar 13 (given at 2, 4, and 6 months).

2.5 Vaccination Schedule for Children 6 Years Through 17 Years of Age

In children 6 years through 17 years of age, Prevnar 13 is administered as single dose. If Prevnar was previously administered, then at least 8 weeks should elapse before receiving Prevnar 13.

2.6 Vaccination Schedule for Adults 18 Years of Age and Older

Prevnar 13 is administered as a single dose.

3 DOSAGE FORMS AND STRENGTHS

Pevnar 13 is a suspension for intramuscular injection available in 0.5 mL single-dose prefilled syringes.

4 CONTRAINDICATIONS

Severe allergic reaction (e.g., anaphylaxis) to any component of Pevnar 13 or any diphtheria toxoid-containing vaccine [see *Description (11)*].

5 WARNINGS AND PRECAUTIONS

5.1 Management of Allergic Reactions

Epinephrine and other appropriate agents used to manage immediate allergic reactions must be immediately available should an acute anaphylactic reaction occur following administration of Pevnar 13.

5.2 Altered Immunocompetence

Individuals with altered immunocompetence, including those at higher risk for invasive pneumococcal disease (e.g., individuals with congenital or acquired splenic dysfunction, HIV infection, malignancy, hematopoietic stem cell transplant, nephrotic syndrome), may have reduced antibody responses to immunization with Pevnar 13 [see *Use in Specific Populations (8.6)*].

5.3 Apnea in Premature Infants

Apnea following intramuscular vaccination has been observed in some infants born prematurely. Decisions about when to administer an intramuscular vaccine, including Pevnar 13, to infants born prematurely should be based on consideration of the individual infant's medical status and the potential benefits and possible risks of vaccination.

6 ADVERSE REACTIONS

Because clinical trials are conducted under widely varying conditions, adverse-reaction rates observed in the clinical trials of a vaccine cannot be directly compared to rates in the clinical trials of another vaccine and may not reflect the rates observed in practice.

6.1 Clinical Trials Experience With Pevnar 13 in Children 6 Weeks Through 17 Years of Age

The safety of Pevnar 13 was evaluated in 13 clinical trials in which 4,729 infants (6 weeks through 11 months of age) and toddlers (12 months through 15 months of age) received at least one dose of Pevnar 13 and 2,760 infants and toddlers received at least one dose of Pevnar active control. Safety data for the first three doses are available for all 13 infant studies; dose 4 data are available for 10 studies; and data for the 6-month follow-up are available for 7 studies. The vaccination schedule and concomitant vaccinations used in these infant trials were consistent with country-specific recommendations and local clinical practice. There were no substantive differences in demographic characteristics between the vaccine groups. By race, 84.0% of subjects were White, 6.0% were Black or African-American, 5.8% were Asian and 3.8% were of 'Other' race (most of these being biracial). Overall, 52.3% of subjects were male infants.

Three studies in the US (Studies 1, 2 and 3)^{1,2,3} evaluated the safety of Pevnar 13 when administered concomitantly with routine US pediatric vaccinations at 2, 4, 6, and 12–15 months of age. Solicited local and systemic adverse reactions were recorded daily by parents/guardians using an electronic diary for 7 consecutive days following each vaccination. For unsolicited adverse events, study subjects were monitored from administration of the first dose until one month after the infant series, and for one month after the administration of the toddler dose. Information regarding unsolicited and serious adverse events, newly diagnosed chronic medical conditions, and hospitalizations since the last visit were

collected during the clinic visit for the fourth-study dose and during a scripted telephone interview 6 months after the fourth-study dose. Serious adverse events were also collected throughout the study period. Overall, the safety data show a similar proportion of Prevnar 13 and Prevnar subjects reporting serious adverse events. Among US study subjects, a similar proportion of Prevnar 13 and Prevnar recipients reported solicited local and systemic adverse reactions as well as unsolicited adverse events.

Serious Adverse Events in All Infant and Toddler Clinical Studies

Serious adverse events were collected throughout the study period for all 13 clinical trials. This reporting period is longer than the 30-day post-vaccination period used in some vaccine trials. The longer reporting period may have resulted in serious adverse events being reported in a higher percentage of subjects than for other vaccines. Serious adverse events reported following vaccination in infants and toddlers occurred in 8.2% among Prevnar 13 recipients and 7.2% among Prevnar recipients. Serious adverse events observed during different study periods for Prevnar 13 and Prevnar respectively were: 1) 3.7% and 3.5% from dose 1 to the blood draw approximately 1 month after the infant series; 2) 3.6% and 2.7% from the blood draw after the infant series to the toddler dose; 3) 0.9% and 0.8% from the toddler dose to the blood draw approximately 1 month after the toddler dose and 4) 2.5% and 2.8% during the 6 month follow-up period after the last dose.

The most commonly reported serious adverse events were in the 'Infections and infestations' system organ class including bronchiolitis (0.9%, 1.1%), gastroenteritis, (0.9%, 0.9%), and pneumonia (0.9%, 0.5%) for Prevnar 13 and Prevnar respectively.

There were 3 (0.063%) deaths among Prevnar 13 recipients, and 1 (0.036%) death in Prevnar recipients, all as a result of sudden infant death syndrome (SIDS). These SIDS rates are consistent with published age specific background rates of SIDS from the year 2000.

Among 6,839 subjects who received at least 1 dose of Prevnar 13 in clinical trials conducted globally, there was 1 hypotonic-hyporesponsive episode adverse reaction reported (0.015%). Among 4,204 subjects who received at least 1 dose of Prevnar in clinical trials conducted globally, there were 3 hypotonic-hyporesponsive episode adverse reactions reported (0.071%). All 4 events occurred in a single clinical trial in Brazil in which subjects received whole cell pertussis vaccine at the same time as Prevnar 13 or Prevnar.

Solicited Adverse Reactions in the Three US Infant and Toddler Studies

A total of 1,907 subjects received at least 1 dose of Prevnar 13 and 701 subjects received at least 1 dose of Prevnar in the three US studies (Studies 1, 2 and 3)^{1,2,3}. Most subjects were White (77.3%), 14.2% were Black or African-American, and 1.7% were Asian; 79.1% of subjects were non-Hispanic and non-Latino and 14.6% were Hispanic or Latino. Overall, 53.6% of subjects were male infants.

The incidence and severity of solicited adverse reactions that occurred within 7 days following each dose of Prevnar 13 or Prevnar administered to US infants and toddlers are shown in Tables 3 and 4.

Table 3: Percentage of US Infant and Toddler Subjects Reporting Solicited Local Reactions at the Prevnar 13 or Prevnar Injection Sites Within 7 Days After Each Vaccination at 2, 4, 6, and 12–15 Months of Age*

Graded Local Reaction	Dose 1		Dose 2		Dose 3		Dose 4	
	Prevnar 13 (N†=1375-1612) %	Prevnar (N†=516-606) %	Prevnar 13 (N†=1069-1331) %	Prevnar (N†=405-510) %	Prevnar 13 (N†=998-1206) %	Prevnar (N†=348-446) %	Prevnar 13 (N†=874-1060) %	Prevnar (N†=283-379) %
Redness‡								
Any	24.3	26.0	33.3	29.7	37.1	36.6	42.3	45.5
Mild	23.1	25.2	31.9	28.7	35.3	35.3	39.5	42.7

Moderate	2.2	1.5	2.7	2.2	4.6	5.1	9.6	13.4 [§]
Severe	0	0	0	0	0	0	0	0
Swelling [‡]								
Any	20.1	20.7	25.2	22.5	26.8	28.4	31.6	36.0 [§]
Mild	17.2	18.7	23.8	20.5	25.2	27.5	29.4	33.8
Moderate	4.9	3.9	3.7	4.9	3.8	5.8	8.3	11.2 [§]
Severe	0	0	0.1	0	0	0	0	0
Tenderness								
Any	62.5	64.5	64.7	62.9	59.2	60.8	57.8	62.5
Interferes with limb movement	10.4	9.6	9.0	10.5	8.4	9.0	6.9	5.7

* Data are from three primary US safety studies (the US Phase 2 infant study [National Clinical Trial (NCT) number NCT00205803] Study 1, the US noninferiority study [NCT00373958] Study 2, and the US lot consistency study [NCT00444457] Study 3). All infants received concomitant routine infant immunizations. Concomitant vaccines and pneumococcal conjugate vaccines were administered in different limbs.

† Number of subjects reporting Yes for at least 1 day or No for all days.

‡ Diameters were measured in caliper units of whole numbers from 1 to 14 or 14+. One caliper unit = 0.5 cm. Measurements were rounded up to the nearest whole number. Intensity of induration and erythema were then characterized as Mild (0.5–2.0 cm), Moderate (2.5–7.0 cm), or Severe (>7.0 cm).

§ Statistically significant difference $p < 0.05$. No adjustments for multiplicity.

Table 4: Percentage of US Infant and Toddler Subjects Reporting Solicited Systemic Adverse Reactions Within 7 Days After Each Vaccination at 2, 4, 6, and 12–15 Months of Age*,†

Graded Systemic Events	Dose 1		Dose 2		Dose 3		Dose 4	
	Prevnar 13 (N* = 1360 – 1707) %	Prevnar (N* = 497–640) %	Prevnar 13 (N* = 1084–1469) %	Prevnar (N* = 409–555) %	Prevnar 13 (N* = 997–1361) %	Prevnar (N* = 354–521) %	Prevnar 13 (N* = 850–1227) %	Prevnar (N* = 278–436) %
Fever [‡]								
Any	24.3	22.1	36.5	32.8	30.3	31.6	31.9	30.6
Mild	23.6	21.7	34.9	31.6	29.1	30.2	30.3	30.0
Moderate	1.1	0.6	3.4	2.8	4.2	3.3	4.4	4.6
Severe	0.1	0.2	0.1	0.3	0.1	0.7	1.0	0
Decreased appetite	48.3	43.6	47.8	43.6	47.6	47.6	51.0	49.4
Irritability	85.6	83.6	84.8	80.4	79.8	80.8	80.4	77.8
Increased sleep	71.5	71.5	66.6	63.4	57.7	55.2	48.7	55.1
Decreased sleep	42.5	40.6	45.6	43.7	46.5	47.7	45.3	40.3

* Number of subjects reporting Yes for at least 1 day or No for all days.

† Data are from three primary US safety studies (the US Phase 2 infant study [NCT00205803] Study 1, the US noninferiority study [NCT00373958] Study 2, and the US lot consistency study [NCT00444457] Study 3). All infants received concomitant routine infant immunizations. Concomitant vaccines and pneumococcal conjugate vaccines were administered in different limbs.

‡ Fever gradings: Mild ($\geq 38^{\circ}\text{C}$ but $\leq 39^{\circ}\text{C}$), Moderate ($> 39^{\circ}\text{C}$ but $\leq 40^{\circ}\text{C}$), and Severe ($> 40^{\circ}\text{C}$). No other systemic event other than fever was graded. Parents reported the use of antipyretic medication to treat or

prevent symptoms in 62 to 75% of subjects after any of the 4 doses. There were no statistical differences in frequencies of adverse reactions reported between the Prevnar 13 and Prevnar groups.

The incidence rates of any fever ($\geq 38.0^{\circ}\text{C}$) were similar on days 1 and 2 following each dose of Prevnar 13 compared to after each dose of Prevnar administered to US infants and toddlers (day 1 = day of vaccination). After dose 1, fever was reported in 11.0–12.7% on day 1 and 6.4–6.8% on day 2. After dose 2, fever was reported in 12.3–13.1% on day 1 and 12.5–12.8% on day 2. After dose 3, fever was reported in 8.0–9.6% on day 1 and 9.1–10.5% on day 2. And after dose 4, fever was reported in 6.3–6.4% on day 1 and 7.3–9.7% on day 2.

Unsolicited Adverse Reactions in the Three US Infant and Toddler Safety Studies

The following were determined to be adverse drug reactions based on experience with Prevnar 13 in clinical trials.

Reactions occurring in greater than 1% of infants and toddlers: diarrhea, vomiting, and rash.

Reactions occurring in less than 1% of infants and toddlers: crying, hypersensitivity reaction (including face edema, dyspnea, and bronchospasm), seizures (including febrile seizures), and urticaria or urticaria-like rash.

Safety Assessments in the Catch-Up Studies in Infants and Children Through 5 Years of Age

In a catch-up study⁴ conducted in Poland (Study 4), 354 children (7 months through 5 years of age) receiving at least one dose of Prevnar 13 were also monitored for safety. All subjects in this study were White and non-Hispanic. Overall, 49.6% of subjects were male infants. The incidence and severity of solicited adverse reactions that occurred within 4 days following each dose of Prevnar 13 administered to pneumococcal-vaccine naïve children 7 months through 5 years of age are shown in Tables 5 and 6.

Table 5: Percentage of Subjects 7 Months Through 5 Years of Age Reporting Solicited Local Reactions Within 4 Days After Each Catch-Up Prevnar 13 Vaccination*

Graded Local Reaction	7 through 11 months			12 through 23 months		24 months through 5 years
	Dose 1 N [†] =86 %	Dose 2 N [†] =86– 87 %	Dose 3 N [†] =78– 82 %	Dose 1 N [†] =108–110 %	Dose 2 N [†] =98– 106 %	Dose 1 N [†] =147–149 %
Redness [‡]						
Any	48.8	46.0	37.8	70.0	54.7	50.0
Mild	41.9	40.2	31.3	55.5	44.7	37.4
Moderate	16.3	9.3	12.5	38.2	25.5	25.7
Severe	0.0	0.0	0.0	0.0	0.0	0.0
Swelling [‡]						
Any	36.0	32.2	25.0	44.5	41.0	36.9
Mild	32.6	28.7	20.5	36.7	36.2	28.2
Moderate	11.6	14.0	11.3	24.8	12.1	20.3
Severe	0.0	0.0	0.0	0.0	0.0	0.0
Tenderness						
Any	15.1	15.1	15.2	33.3	43.7	42.3
Interferes with limb movement	1.2	3.5	6.4	0.0	4.1	4.1

* Study conducted in Poland (NCT00452452) Study 4.

[†] Number of subjects reporting Yes for at least 1 day or No for all days.

[‡] Diameters were measured in caliper units of whole numbers from 1 to 14 or 14+. One caliper unit =

0.5 cm. Measurements were rounded up to the nearest whole number. Intensity of redness and swelling were then characterized as Mild (0.5–2.0 cm), Moderate (2.5–7.0 cm), or Severe (>7.0 cm).

Table 6: Percentage of Subjects 7 Months Through 5 Years of Age Reporting Solicited Systemic Adverse Reactions Within 4 Days After Each Catch-Up Prevnar 13 Vaccination*

Systemic Reaction	7 through 11 months			12 through 23 months		24 months through 5 years
	Dose 1 N [†] =86– 87 %	Dose 2 N [†] =86– 87 %	Dose 3 N [†] =78– 81 %	Dose 1 N [†] =108 %	Dose 2 N [†] =98– 100 %	Dose 1 N [†] =147–148 %
Fever [‡]						
Mild	3.4	8.1	5.1	3.7	5.1	0.7
Moderate	1.2	2.3	1.3	0.9	0.0	0.7
Severe	0.0	0.0	0.0	0.0	0.0	0.0
Decreased appetite	19.5	17.2	17.5	22.2	25.5	16.3
Irritability	24.1	34.5	24.7	30.6	34.0	14.3
Increased sleep	9.2	9.3	2.6	13.0	10.1	11.6
Decreased sleep	24.1	18.4	15.0	19.4	20.4	6.8

* Study conducted in Poland (NCT00452452) Study 4.

† Number of subjects reporting Yes for at least 1 day or No for all days.

‡ Fever gradings: Mild ($\geq 38^{\circ}\text{C}$ but $\leq 39^{\circ}\text{C}$), Moderate ($> 39^{\circ}\text{C}$ but $\leq 40^{\circ}\text{C}$), and Severe ($> 40^{\circ}\text{C}$). No other systemic event other than fever was graded.

A US study⁵ (Study 5) evaluated the use of Prevnar 13 in children previously immunized with Prevnar. In this open label trial, 596 healthy children 15 through 59 months of age previously vaccinated with at least 3 doses of Prevnar, received 1 or 2 doses of Prevnar 13. Children 15 months through 23 months of age (group 1) received 2 doses, and children 24 months through 59 months of age (group 2) received one dose. Most subjects were White (74.3%), 14.9% were Black or African-American, and 1.2% were Asian; 89.3% of subjects were non-Hispanic and non-Latino and 10.7% were Hispanic or Latino. Overall, 52.2% of subjects were male.

The incidence and severity of solicited adverse reactions that occurred within 7 days following one dose of Prevnar 13 administered to children 15 months through 59 months of age are shown in Tables 7 and 8.

Table 7: Percentage of Subjects 15 Months Through 59 Months of Age, Previously Vaccinated With 3 or 4 Prior Infant Doses of Prevnar, Reporting Solicited Local Reactions Within 7 Days After One Supplemental Prevnar 13 Vaccination*

Graded Local Reaction	15 months through 23 months [†]		24 months through 59 months [‡]
	1 dose Prevnar 13 3 prior Prevnar doses N [§] =67–72 %	1 dose Prevnar 13 4 prior Prevnar doses N [§] =154–184 %	1 dose Prevnar 13 3 or 4 prior Prevnar doses N [§] =209–238 %
Redness [¶]			
Any	26.4	28.2	35.4
Mild	18.8	24.3	31.1

Moderate	11.4	7.5	12.1
Severe	1.5	0.0	0.0
Swelling[¶]			
Any	23.9	19.6	20.7
Mild	18.6	16.4	17.2
Moderate	8.8	8.1	7.5
Severe	0.0	0.0	0.0
Tenderness			
Any	48.6	47.3	62.6
Interferes with limb movement	5.9	6.4	10.7

* Study conducted in US NCT00761631 (Study 5).

† Dose 2 data not shown.

‡ The data for this age group are only represented as a single result as 95% of children received 4 doses of Prevnar prior to enrollment.

§ Number of subjects reporting Yes for at least 1 day or No for all days.

¶ Diameters were measured in caliper units of whole numbers from 1 to 14 or 14+. One caliper unit = 0.5 cm. Measurements were rounded up to the nearest whole number. Intensity of redness and swelling were then characterized as Mild (0.5–2.0 cm), Moderate (2.5–7.0 cm), or Severe (>7.0 cm).

Table 8: Percentage of Subjects 15 Months Through 59 Months of Age, Previously Vaccinated With 3 or 4 Prior Infant Prevnar Doses, Reporting Solicited Systemic Adverse Reactions Within 7 Days After One Supplemental Prevnar 13 Vaccination*

	15 through 23 months[†]		24 months through 59 months[‡]
Systemic Reaction	1 dose Prevnar 13 3 prior Prevnar doses N[§]=66–75 %	1 dose Prevnar 13 4 prior Prevnar doses N[§]=154–189 %	1 dose Prevnar 13 3 or 4 prior Prevnar doses N[§]=209–236 %
Fever[¶]			
Any	19.1	19.9	8.1
Mild	16.2	17.4	7.6
Moderate	6.1	3.9	1.9
Severe	0.0	0.0	0.5
Decreased appetite	44.4	39.3	28.1
Irritability	73.3	65.1	45.8
Increased sleep	35.2	35.3	18.8
Decreased sleep	25.0	29.7	14.8

* Study conducted in US NCT00761631 (Study 5).

† Dose 2 data not shown.

‡ The data for this age group are only represented as a single result as 95% of children received 4 doses of Prevnar prior to enrollment.

§ Number of subjects reporting Yes for at least 1 day or No for all days.

¶ Fever gradings: Mild ($\geq 38^{\circ}\text{C}$ but $\leq 39^{\circ}\text{C}$), Moderate ($> 39^{\circ}\text{C}$ but $\leq 40^{\circ}\text{C}$), and Severe ($> 40^{\circ}\text{C}$). No other systemic event other than fever was graded.

Clinical Trials Experience With Prevnar 13 in Children 5 Through 17 Years of Age

In a US study⁵ (Study 5), the safety of Prevnar 13 was evaluated in children 5 through 9 years of age previously immunized with at least one dose of Prevnar, and in children 10 through 17 years of age with no prior pneumococcal vaccination. In this open label trial, 592 children, including those with asthma, received a single dose of Prevnar 13. The percentage of children 5 through 9 years of age who received 3 and 4 prior doses of Prevnar was 29.1% and 54.5% respectively.

Most subjects were White (72.8%), 21.8% were Black or African-American, and 1.5% were Asian; 91.4% of subjects were non-Hispanic and non-Latino and 8.6% were Hispanic or Latino. Overall, 51.2% of subjects were male.

The incidence and severity of solicited adverse reactions that occurred within 7 days following one dose of Prevnar 13 administered to children 5 through 17 years of age are shown in Tables 9 and 10.

Table 9: Percentage of Subjects 5 Through 17 Years of Age, Reporting Solicited Local Reactions Within 7 Days After Prevnar 13 Vaccination*

Local Reaction	Vaccine Group (as Administered)					
	Prevnar 13 (5 Through 9 Years)			Prevnar 13 (10 Through 17 Years)		
	N [†]	n [‡]	%	N [†]	n [‡]	%
Redness						
Any	233	100	42.9	232	70	30.2
Mild [§]	226	63	27.9	226	48	21.2
Moderate [§]	218	48	22.0	221	31	14.0
Severe [§]	212	7	3.3	213	4	1.9
Swelling						
Any	226	85	37.6	233	86	36.9
Mild [§]	220	48	21.8	221	50	22.6
Moderate [§]	219	48	21.9	226	48	21.2
Severe [§]	211	7	3.3	214	4	1.9
Tenderness						
Any	265	230	86.8	283	252	89.0
Significant [¶]	221	43	19.5	242	106	43.8

* Study conducted in US NCT00761631 (Study 5).

† N = number of subjects reporting Yes for at least 1 day or No for all days.

‡ n = Number of subjects reporting the specific characteristic.

§ Mild, 0.5 – 2.0 cm; moderate, 2.5 – 7.0 cm; severe, >7.0 cm.

¶ Significant = present and interfered with limb movement.

Table 10: Percentage of Subjects 5 Through 17 Years of Age, Reporting Solicited Systemic Adverse Reactions Within 7 Days After Prevnar 13 Vaccination*

Systemic Event	Vaccine Group (as Administered)					
	Prevnar 13 (5 Through 9 Years)			Prevnar 13 (10 Through 17 Years)		
	N [†]	n [‡]	%	N [†]	n [‡]	%
Any fever ≥38°C	214	13	6.1	214	12	5.6
Mild [§]	212	9	4.2	214	11	5.1
Moderate [§]	212	5	2.4	212	1	0.5

Severe [§]	210	1	0.5	212	1	0.5
Decreased appetite	227	52	22.9	223	51	22.9
Irritability	234	73	31.2	234	59	25.2
Increased sleep	226	48	21.2	229	61	26.6
Decreased sleep	212	12	5.7	224	42	18.8
Hives (urticaria)	213	4	1.9	214	3	1.4

* Study conducted in US NCT00761631 (Study 5).

† N = number of subjects reporting Yes for at least 1 day or No for all days.

‡ n = Number of subjects reporting the event.

§ Fever gradings: Mild ($\geq 38^{\circ}\text{C}$ but $\leq 39^{\circ}\text{C}$), Moderate ($> 39^{\circ}\text{C}$ but $\leq 40^{\circ}\text{C}$), and Severe ($> 40^{\circ}\text{C}$). No other systemic event other than fever was graded. Parents reported the use of antipyretic medication to treat or prevent symptoms in 45.1% and 33.1% of subjects 5 through 9 years of age and 10 through 17 years of age, respectively.

6.2 Clinical Trials Experience With Prevnar 13 in Adults ≥ 18 Years of Age

The safety of Prevnar 13 was assessed in 7 clinical studies (Studies 6–12)^{6–12} conducted in the US and Europe which included 91,593 adults (48,806 received Prevnar 13) ranging in age from 18 through 101 years.

The 48,806 Prevnar 13 recipients included 899 adults who were aged 18 through 49 years, 2,616 adults who were aged 50 through 64 years, 45,291 adults aged 65 years and older. Of the 48,806 Prevnar 13 recipients, 46,890 adults had not previously received Pneumovax[®] 23 (pneumococcal polysaccharide vaccine [23-valent], PPSV23) ("PPSV23 unvaccinated") and 1,916 adults were previously vaccinated ("PPSV23 previously vaccinated") with PPSV23 at least 3 years prior to enrollment.

Safety and Immunogenicity Studies

Safety and immunogenicity of Prevnar 13 is supported by 6 clinical studies. Study 6⁶ evaluated the safety and immunogenicity of Prevnar 13 in adults 18 through 64 years of age who had not received a previous dose of pneumococcal vaccine. Adults 18 through 59 years of age received a single dose of Prevnar 13, and adults 60 through 64 years of age received a single dose of Prevnar 13 or PPSV23.

Study 7 was randomized and compared the safety and immunogenicity of Prevnar 13 with PPSV23 as a single dose in adults ≥ 70 years vaccinated with PPSV23 (≥ 5 years prior to enrollment).⁷ Study 8 was randomized and evaluated the safety and immunogenicity of Prevnar 13 and PPSV23 in different sequential order in PPSV23 naive adults aged 60 through 64 years⁸.

One clinical safety study⁹ (Study 9) of Prevnar 13, conducted in PPSV23 previously vaccinated (≥ 3 years prior to enrollment) adults aged ≥ 68 years was a single arm study. Two studies, one in the US¹⁰ (Study 10) in adults aged 50 through 59 years and the other in Europe¹¹ (Study 11) in adults aged ≥ 65 years, evaluated the concomitant administration of Prevnar 13 with inactivated influenza vaccine, trivalent (Fluarix[®], A/H1N1, A/H3N2, and B, Fall 2007/Spring 2008: IIV3) in these two age groups in PPSV23 unvaccinated adults.

The total safety population in the 6 safety and immunogenicity studies was 7,097. In 5 of the 6 safety and immunogenicity studies, more females than males were enrolled (50.2% – 61.8%). Across the 6 studies the racial distribution included: $> 85\%$ White; 0.2%–10.7% Black or African American; 0%–1.7% Asian; $< 1\%$ Native Hawaiian or other Pacific Islander; $\leq 1\%$, American Indian or Alaskan Native. Ethnicity data were not collected in Study 11; in the 5 other studies 0.6%–4.8% were Hispanic or Latino.

In five studies,^{6–8,10,11} subjects with pre-existing underlying diseases were enrolled if the medical condition was stable (did not require a change in therapy or hospitalization for worsening disease for 12 weeks before receipt of study vaccine) except in Study 9 where subjects were enrolled if the medical

condition was stable for 6 or more weeks before receipt of study vaccine.

In the 6 safety and immunogenicity studies,^{6–11} subjects were excluded from study participation due to prior receipt of diphtheria toxoid-containing vaccines within 6 months of study vaccine. However, the time of prior receipt of a diphtheria toxoid-containing vaccine was not recorded.

Solicited adverse reactions for Prevnar 13 in the safety and immunogenicity studies were monitored by subjects recording local adverse reactions and systemic reactions daily using an electronic diary for 14 consecutive days following vaccination. Unsolicited serious and non-serious adverse events were collected for one month after each vaccination. In addition, serious adverse events were collected for an additional 5 months after each vaccination (at the 6-month follow-up phone contact) in all studies except Study 11.

Following licensure of Prevnar 13 in adults ≥ 50 years of age, a randomized, double-blind, placebo-controlled US study (Study 13) was conducted to evaluate concomitant administration of Prevnar 13 with inactivated influenza vaccine, quadrivalent (Fluzone[®] Quadrivalent, A/H1N1, A/H3N2, B/Brisbane, and B/Massachusetts, Fall 2014/Spring 2015: IIV4) in PPSV23 previously vaccinated adults ≥ 50 years of age. Unsolicited serious and non-serious adverse events were collected as described above for Studies 6–10.

Efficacy Study

Study 12¹² was a randomized double-blind placebo-controlled study conducted in the Netherlands in community-dwelling adults aged 65 years and older with no prior pneumococcal vaccination history. A total of 84,496 subjects received either a single dose of Prevnar 13 (42,240) or placebo (42,256) in a 1:1 randomization. Among the 84,496 subjects, 58,072 (68.7%) were ≥ 65 to < 75 years of age, 23,481 (27.8%) were ≥ 75 and < 85 years of age, and 2,943 (3.5%) were ≥ 85 years of age. In the total safety population, more males (55.9%) were enrolled than females. The racial distribution was 98.5% White, 0.3% Black, 0.7% Asian, 0.5% Other, with $< 0.1\%$ having missing data.

Adults with immunocompromising conditions or receiving immunosuppressive therapy and adults residing in a long-term care facility or requiring semiskilled nursing care were excluded. Adults with pre-existing medical conditions, as well as subjects with a history of smoking were eligible for enrollment. In the safety population, 42.3% of subjects had pre-existing medical conditions including heart disease (25.4%), lung disease or asthma (15.1%) and type 1 and type 2 diabetes mellitus (12.5%). Smoking was reported at baseline by 12.3% of the subjects.

For a subset of 2,011 subjects (1,006 Prevnar 13 recipients and 1,005 placebo recipients), solicited adverse reactions were monitored by recording local and systemic events using electronic diaries for 7 days after vaccination; unsolicited adverse events were collected for 28 days after vaccination, and serious adverse events were collected for 6 months after vaccination. For the remaining 41,231 Prevnar 13 and 41,250 placebo vaccinated subjects, serious adverse events were collected for 28 days after vaccination.

Serious Adverse Events in Adult Clinical Studies

Safety and Immunogenicity Studies

Across the 6 safety and immunogenicity studies,^{6–11} serious adverse events within 1 month of vaccination were reported after an initial study dose in 0.2%–1.4% of 5,057 subjects vaccinated with Prevnar 13, and in 0.4%–1.7% of 1,124 subjects vaccinated after an initial study dose of PPSV23. From 1 month to 6 months after an initial study dose, serious adverse events were reported in 0.2%–5.8% of subjects vaccinated during the studies with Prevnar 13 and in 2.4%–5.5% of subjects vaccinated with PPSV23. One case of erythema multiforme occurred 34 days after receipt of a second dose of Prevnar 13.

Twelve of 5,667 (0.21%) Prevnar 13 recipients and 4 of 1,391 (0.29%) PPSV23 recipients died. Deaths occurred between Day 3 and Day 309 after study vaccination with Prevnar 13 or PPSV23. Two of 12 deaths occurred within 30 days of vaccination and both deaths were in subjects > 65 years of age.

Any	96.7	88.8	80.1	73.4	69.2 [¶]	58.3	36.1 [¶]	6.1
Mild	93.2	85.9	78.6 [¶]	68.6	66.1 [¶]	52.9	32.9 [¶]	5.6
Moderate	77.1	39.5	23.3	30.0	20.1	21.7	7.7 [¶]	0.6
Severe	16.0	3.6	1.7	8.6 [¶]	2.3	0.8	0.3	0.1
Limitation of arm movement [Ⓓ]								
Any	75.2	40.7	28.5	30.8	23.5	28.2	14.1 [¶]	3.2
Mild	71.5	38.6	26.9	29.3	22.7	26.1	12.4 [¶]	2.5
Moderate	18.5	2.9	2.2	3.8	1.2	3.1	1.7 [¶]	0.5
Severe	15.6	2.9	1.7	4.3	1.1	2.3	1.2	0.7

* Studies conducted in US NCT00427895 (Study 6) and NCT00574548 (Study 8) reported local reactions within 14 days. Study conducted in the Netherlands NCT00744263 (Study 12) reported local reactions within 7 days.

† Open label administration of Prevnar 13.

‡ Number of subjects with known values (number of subjects reporting yes for at least one day or no for all days).

§ Diameters were measured in caliper units of whole numbers from 1 to 21 or 21+. One caliper unit = 0.5 cm. Measurements were rounded up to the nearest whole number. Intensity of redness and swelling were then characterized as Mild = 2.5 to 5.0 cm, Moderate = 5.1 to 10.0 cm, and Severe is >10.0 cm.

¶ Statistically significant difference $p < 0.05$. No adjustments for multiplicity.

Mild = awareness of symptom but easily tolerated, Moderate = discomfort enough to cause interference with usual activity, Severe = incapacitating with inability to do usual activity.

Ⓓ Mild = some limitation of arm movement, Moderate = unable to move arm above head but able to move arm above shoulder, and Severe = unable to move arm above shoulder.

Table 12 - Percentage of Subjects With Solicited Local Adverse Reactions in PPSV23 Previously Vaccinated Adults *

Age in Years	Study 7		Study 9
	≥70		≥68
Local Reaction	Prevnar 13 N [†] =306–362 %	PPSV23 N [†] =324–383 %	Prevnar 13 [‡] N [†] =664–777 %
Redness [§]			
Any	10.8	22.2 [¶]	14.3
Mild	9.5	13.5	12.6
Moderate	4.7	11.5 [¶]	6.5
Severe	1.7	4.8 [¶]	1.1
Swelling [§]			
Any	10.4	23.1 [¶]	12.8
Mild	8.9	14.0 [¶]	10.9
Moderate	4.0	13.6 [¶]	5.5
Severe	0.0	4.8 [¶]	0.6
Pain [#]			
Any	51.7	58.5	51.0
Mild	50.1	54.1	49.4
Moderate	7.5	23.6 [¶]	9.0
Severe	1.3	2.3	0.2

new muscle pain	82.0	61.8	56.2	57.8	46.9	51.5	18.4 [§]	8.4
Generalized aggravated muscle pain	55.9	39.9	32.6	37.3	22.0	32.5 [§]	9.1 [§]	4.4
Generalized new joint pain	41.7	31.5	24.4	30.1	15.5	23.8 [§]	7.4	5.4
Generalized aggravated joint pain	28.6	25.6	24.9	21.4	14.0	21.1	5.2	4.2

* Studies conducted in US NCT00427895 (Study 6) and NCT00574548 (Study 8) reported systemic events within 14 days. Study conducted in the Netherlands NCT00744263 (Study 12) reported systemic events within 7 days.

† Open label administration of Prevnar 13.

‡ Number of subjects with known values (number of subjects reporting yes for at least one day or no for all days).

§ Statistically significant difference $p < 0.05$. No adjustments for multiplicity.

¶ Fevers $>40.0^{\circ}\text{C}$ were confirmed to be data entry errors and remain in the table for the following: 1 case in the 18- to 49- year-old cohort (Study 6), and 7 cases in the Prevnar 13 group and 3 cases in placebo group (Study 12). For the other cohorts in Study 6 and for Study 8, data entry errors were removed.

Table 14 - Percentage of Subjects With Systemic Events in PPSV23 Previously Vaccinated Adults*

Age in Years	Study 7 ≥70		Study 9 ≥68
	Prevnar 13 N [†] =299–350 %	PPSV23 N [†] =303–367 %	Prevnar 13 [‡] N [†] =635–733 %
Systemic Event			
Fever			
≥38.0°C	1.0	2.3	1.1
38.0°C to 38.4°C	1.0	2.0	0.8
38.5°C to 38.9°C	0.0	0.0	0.0
39.0°C to 40.0°C	0.0	0.3	0.3
>40.0°C	0.0	0.0	0.0
Fatigue	34.0	43.3 [§]	34.4
Headache	23.7	26.0	26.1
Chills	7.9	11.2	7.5
Rash	7.3	16.4 [§]	8.4
Vomiting	1.7	1.3	0.9
Decreased appetite	10.4	11.5	11.2
Generalized new muscle pain	36.8	44.7 [§]	25.3
Generalized aggravated muscle pain	20.6	27.5 [§]	12.3
Generalized new joint pain	12.6	14.9	12.8

Generalized aggravated joint pain	11.6	16.5	9.7
-----------------------------------	------	------	-----

* Study conducted in US and Sweden NCT00546572 (Study 7) reported systemic events within 14 days. Study conducted in US, Sweden and Germany NCT00500266 (Study 9) reported systemic events within 14 days.

† Number of subjects with known values.

‡ Open label administration of Prevnar 13.

§ Statistically significant difference $p < 0.05$. No adjustments for multiplicity.

Safety Results from Adult Clinical Study of Concomitant Administration of Prevnar 13 and IIV4 (Fluzone Quadrivalent) (Study 13)

The safety profile of Prevnar 13 when administered concomitantly with seasonal inactivated influenza vaccine, quadrivalent, to PPSV23 previously vaccinated adults ≥ 50 years of age was generally consistent with the known safety profile of Prevnar 13.

6.3 Post-marketing Experience With Prevnar 13 in Infants and Toddlers

The following adverse events have been reported through passive surveillance since market introduction of Prevnar 13. Because these events are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to the vaccine. The following adverse events were included based on one or more of the following factors: severity, frequency of reporting, or strength of evidence for a causal relationship to Prevnar 13 vaccine.

Administration site conditions: Vaccination-site dermatitis, vaccination-site pruritus, vaccination-site urticaria

Blood and lymphatic system disorders: Lymphadenopathy localized to the region of the injection site

Cardiac disorders: Cyanosis

Immune system disorders: Anaphylactic/anaphylactoid reaction including shock

Nervous system disorders: Hypotonia

Skin and subcutaneous tissue disorders: Angioneurotic edema, erythema multiforme

Respiratory: Apnea

Vascular disorders: Pallor

7 DRUG INTERACTIONS

7.1 Concomitant Immunizations

In clinical trials with infants and toddlers, Prevnar 13 was administered concomitantly with the following US-licensed vaccines: Pediarix [Diphtheria and Tetanus Toxoids and Acellular Pertussis Adsorbed, Hepatitis B (Recombinant) and Inactivated Poliovirus Vaccine Combined] (DTaP-HBV-IPV) and ActHIB [Haemophilus b Conjugate Vaccine (Tetanus Toxoid Conjugate)] (PRP-T) for the first three doses and with PedvaxHIB [Haemophilus b Conjugate Vaccine (Meningococcal Protein Conjugate)] (PRP-OMP), M-M-R II [Measles, Mumps, Rubella Virus Vaccine Live] (MMR) and Varivax [Varicella Virus Vaccine Live], or ProQuad [Measles, Mumps, Rubella and Varicella Virus Vaccine Live] (MMRV) and VAQTA [Hepatitis A vaccine, Inactivated] (HepA) for dose 4 [see *Clinical Studies (14.2) and Adverse Reactions (6.1)*].

In children and adolescents, data are insufficient to assess the concomitant administration of Prevnar 13 with Human Papillomavirus Vaccine (HPV), Meningococcal Conjugate Vaccine (MCV4) and Tetanus Toxoid, Reduced Diphtheria Toxoid and Acellular Pertussis Vaccine, Adsorbed (Tdap).

In adults, Prevnar 13 was administered concomitantly with US-licensed inactivated influenza vaccines, trivalent and quadrivalent (Studies 10, 11 and 13)[*see Clinical Studies (14.4) and Adverse Reactions (6.2)*]. There are no data on the concomitant administration of Prevnar 13 with diphtheria toxoid-containing vaccines and other vaccines licensed for use in adults 50 years of age and older.

When Prevnar 13 is administered at the same time as another injectable vaccine(s), the vaccines should always be administered with different syringes and given at different injection sites.

Do not mix Prevnar 13 with other vaccines/products in the same syringe.

7.2 Immunosuppressive Therapies

Individuals with impaired immune responsiveness due to the use of immunosuppressive therapy (including irradiation, corticosteroids, antimetabolites, alkylating agents, and cytotoxic agents) may not respond optimally to active immunization.

7.3 Antipyretics

A post-marketing clinical study conducted in Poland using a non-US vaccination schedule (2, 3, 4, and 12 months of age) evaluated the impact of prophylactic oral acetaminophen on antibody responses to Prevnar 13. The data show that 3 doses of acetaminophen (the first dose administered at the time of each vaccination and the subsequent doses at 6 to 8 hour intervals) reduced the antibody response to some serotypes following the third dose of Prevnar 13, compared with responses among infants who received antipyretics only as needed for treatment. Reduced antibody responses were not observed after the fourth dose of Prevnar 13 when acetaminophen was administered prophylactically.

7.4 Prior Vaccination with PPSV23

Prior receipt of PPSV23 within 1 year results in diminished immune responses to Prevnar 13 compared to PPSV23 naïve individuals [*see Clinical Studies (14.3)*].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

All pregnancies have a risk of birth defect, loss, or other adverse outcomes. In the US general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively. Available data on Prevnar 13 administered to pregnant women are insufficient to inform vaccine-associated risks in pregnancy.

A developmental toxicity study has been performed in female rabbits administered Prevnar 13 prior to mating and during gestation. Each dose was approximately 20 times the human dose. This study revealed no evidence of harm to the fetus due to Prevnar 13 (see 8.1 Data).

Data

Animal

In a developmental toxicity study, female rabbits were administered Prevnar 13 by intramuscular injection twice prior to mating (17 days and 3 days prior to mating) and twice during gestation (gestation days 10 and 24), 0.5 mL/rabbit/occasion (each dose approximately 20 times the human dose). No adverse effects on pre-weaning development were observed. There were no vaccine-related fetal malformations or variations.

8.2 Lactation

Risk Summary

Data are not available to assess the effects of Prevnar 13 on the breastfed infant or on milk production/excretion. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Prevnar 13 and any potential adverse effects on the breastfed child from Prevnar 13 or from the underlying maternal condition. For preventive vaccines, the underlying maternal condition is susceptibility to disease prevented by the vaccine.

8.4 Pediatric Use

Safety and effectiveness of Prevnar 13 in children below the age of 6 weeks have not been established.

8.5 Geriatric Use

Of the total number of Prevnar 13 recipients aged 50 years and older in clinical studies (N=47,907), 94.5% (45,291 of 47,907) were 65 years and older and 30.3% (14,498 of 47,907) were 75 years and older [see *Clinical Studies (14.1) and (14.3)*].

8.6 High Risk Populations

Individuals with the diseases or conditions listed below are at increased risk of pneumococcal disease. Immunogenicity and safety data in these populations are limited.

Infants Born Prematurely

Immune responses elicited by Prevnar 13 administered on a US schedule to preterm infants have not been studied. When preterm infants (<37 weeks gestational age, N=100) were administered 4 doses of Prevnar 13 on a non-US schedule, the serotype-specific IgG antibody responses after the third and fourth dose were lower compared to responses among term infants (≥ 37 weeks gestational age, N=100) for some serotypes; the effectiveness of Prevnar 13 in preterm infants cannot be established from this study.

Children with Sickle Cell Disease

In an open-label, single-arm, descriptive study, 2 doses of Prevnar 13 were administered 6 months apart to children ≥ 6 to <18 years of age with sickle cell disease who previously received PPSV23 at least 6 months prior to enrollment. Children with a prior history of pneumococcal conjugate vaccination were excluded. For all vaccine serotypes, anti-pneumococcal opsonophagocytic activity (OPA) geometric mean antibody titers (GMTs) were higher after the first dose compared to pre-vaccination (N=95–131); OPA GMTs following the first and second dose were comparable. The effectiveness of Prevnar 13 in this specific population has not been established.

Individuals with Hematopoietic Stem Cell Transplant

In an open-label, single-arm, descriptive study, 4 doses of Prevnar 13 were administered to subjects ≥ 2 years of age (range 2 to 71 years) who had received an allogeneic hematopoietic stem cell transplant 3 to 6 months prior to enrollment. All subjects had a history of stable engraftment (absolute neutrophil count $> 1000/\mu\text{L}$, platelet count $> 50,000/\mu\text{L}$), and did not have uncontrolled graft versus host disease. The first three doses of Prevnar 13 were administered one month apart, followed by a fourth dose of Prevnar 13 six months after the third dose. Sera were obtained approximately one month after each vaccination. Immune responses (IgG GMCs) after the first dose of Prevnar 13 were numerically higher for all serotypes compared with baseline. In addition, after each subsequent dose of Prevnar 13, IgG GMCs for all serotypes were numerically higher than responses after the previous dose. A post hoc analysis of the immune responses as measured by OPA antibody assay showed the pattern of functional antibody responses to be consistent with IgG responses for each serotype. The effectiveness of Prevnar 13 in this specific population has not been established.

Individuals with HIV Infection

In an open-label, single-arm, descriptive study, 3 doses of Prevnar 13 were administered 6 months apart to HIV-infected adults ≥ 18 years of age (median age 48 years), with CD4 counts ≥ 200 cells/ μL and

serum HIV RNA titer <50,000 copies/mL. All subjects had been vaccinated previously with PPSV23 at least 6 months prior to enrollment. For all vaccine serotypes anti-pneumococcal OPA GMTs were numerically higher after the first dose compared to pre-vaccination (N=227–253); OPA GMTs following the first, second and third dose were generally comparable. The effectiveness of Prevnar 13 in this specific population has not been established.

In an open-label, single-arm, descriptive study, 3 doses of Prevnar 13 were administered 1 month apart to HIV-infected subjects ≥ 6 years of age with CD4 counts ≥ 200 cells/ μ L, and serum HIV RNA titer <50,000 copies/mL. Subjects had not previously been vaccinated with a pneumococcal vaccine. For all vaccine serotypes anti-pneumococcal OPA GMTs were numerically higher after the first dose compared to pre-vaccination (N=197–257); OPA GMTs following the first, second and third dose were generally comparable. The effectiveness of Prevnar 13 in this specific population has not been established.

11 DESCRIPTION

Prevnar 13, Pneumococcal 13-valent Conjugate Vaccine (Diphtheria CRM₁₉₇ Protein) is a sterile suspension of saccharides of the capsular antigens of *Streptococcus pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, and 23F, individually linked to non-toxic diphtheria CRM₁₉₇ protein. Each serotype is grown in soy peptone broth. The individual polysaccharides are purified through centrifugation, precipitation, ultrafiltration, and column chromatography. The polysaccharides are chemically activated to make saccharides, which are directly conjugated by reductive amination to the protein carrier CRM₁₉₇, to form the glycoconjugate. CRM₁₉₇ is a nontoxic variant of diphtheria toxin isolated from cultures of *Corynebacterium diphtheriae* strain C7 (β 197) grown in a casamino acids and yeast extract-based medium or in a chemically-defined medium. CRM₁₉₇ is purified through ultrafiltration, ammonium sulfate precipitation, and ion-exchange chromatography. The individual glycoconjugates are purified by ultrafiltration and column chromatography and analyzed for saccharide to protein ratios, molecular size, free saccharide, and free protein.

The individual glycoconjugates are compounded to formulate Prevnar 13. Potency of the formulated vaccine is determined by quantification of each of the saccharide antigens and by the saccharide to protein ratios in the individual glycoconjugates. Each 0.5 mL dose of the vaccine is formulated to contain approximately 2.2 μ g of each of *Streptococcus pneumoniae* serotypes 1, 3, 4, 5, 6A, 7F, 9V, 14, 18C, 19A, 19F, 23F saccharides, 4.4 μ g of 6B saccharides, 34 μ g CRM₁₉₇ carrier protein, 100 μ g polysorbate 80, 295 μ g succinate buffer and 125 μ g aluminum as aluminum phosphate adjuvant.

The tip cap and rubber plunger of the prefilled syringe are not made with natural rubber latex.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Prevnar 13, comprised of pneumococcal polysaccharides conjugated to a carrier protein (CRM₁₉₇), elicits a T-cell dependent immune response. Protein carrier-specific T-cells provide the signals needed for maturation of the B-cell response.

Nonclinical and clinical data support opsonophagocytic activity, as measured by opsonophagocytic activity (OPA) antibody assay, as a contributor to protection against pneumococcal disease. The OPA antibody assay provides an in vitro measurement of the ability of serum antibodies to eliminate pneumococci by promoting complement-mediated phagocytosis and is believed to reflect relevant in vivo mechanisms of protection against pneumococcal disease. OPA antibody titers are expressed as the reciprocal of the highest serum dilution that reduces survival of the pneumococci by at least 50%.

In infants that have received Prevnar 13, opsonophagocytic activity correlates well with serotype specific anti-capsular polysaccharide IgG levels as measured by ELISA. A serum anti-capsular polysaccharide antibody concentration of 0.35 μ g/mL as measured by ELISA one month after the third

dose as a single antibody reference concentration was used to estimate the effectiveness of Prevnar 13 against invasive pneumococcal disease (IPD) in infants and children. The assay used for this determination is a standardized ELISA involving pre-absorption of the test sera with pneumococcal C-polysaccharide and serotype 22F polysaccharide to reduce non-specific background reactivity. The single antibody reference value was based on pooled efficacy estimates from three placebo-controlled IPD efficacy trials with either Prevnar or the investigational 9-valent CRM₁₉₇ conjugate pneumococcal polysaccharide vaccine. This reference concentration is only applicable on a population basis and cannot be used to predict protection against IPD on an individual basis. Functional antibodies elicited by the vaccine (as measured by a dribble opsonophagocytic activity [dOPA] antibody assay) were also evaluated in infants.

In adults, an antipolysaccharide binding antibody IgG level to predict protection against invasive pneumococcal disease or non-bacteremic pneumonia has not been defined. Noninferiority trials for Prevnar 13 were designed to show that functional OPA antibody responses (as measured by a microcolony OPA [mcOPA] antibody assay) for the Prevnar 13 serotypes are noninferior and for some serotypes superior to the common serotypes in the currently licensed pneumococcal polysaccharide vaccine (PPSV23). OPA antibody titers measured in the mcOPA antibody assay cannot be compared directly to titers measured in the dOPA antibody assay.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Prevnar 13 has not been evaluated for the potential to cause carcinogenicity, genotoxicity, or impairment of male fertility. In a study in rabbits, no vaccine-related effects were found regarding reproductive performance including female fertility [see *Use in Specific Populations (8.1)*].

14 CLINICAL STUDIES

14.1 Efficacy Data

Prevnar Efficacy Data

Invasive Pneumococcal Disease (IPD)

Prevnar (Pneumococcal 7-valent Conjugate Vaccine [Diphtheria CRM₁₉₇ Protein]) was licensed in the US for infants and children in 2000, following a randomized, double-blind clinical trial in a multiethnic population at Northern California Kaiser Permanente (NCKP) from October 1995 through August 20, 1998, in which 37,816 infants were randomized to receive either Prevnar or a control vaccine (an investigational meningococcal group C conjugate vaccine [MnCC]) at 2, 4, 6, and 12–15 months of age. In this study, the efficacy of Prevnar against invasive disease due to *S. pneumoniae* in cases accrued during this period was 100% in both the per-protocol and intent-to-treat analyses (95% confidence interval [CI]: 75.4%, 100% and 81.7%, 100%, respectively). Data accumulated through an extended follow-up period to April 20, 1999, resulted in similar efficacy estimates of 97.4% in the per-protocol analysis and 93.9% in the intent-to-treat analysis (95% CI: 82.7%, 99.9% and 79.6%, 98.5%, respectively).

Acute Otitis Media (AOM)

The efficacy of Prevnar against otitis media was assessed in two clinical trials: a trial in Finnish infants at the National Public Health Institute and the efficacy trial in US infants at Northern California Kaiser Permanente (NCKP).

The Finnish Otitis Media (FinOM) trial was a randomized, double-blind trial in which 1,662 infants were equally randomized to receive either Prevnar or a control vaccine Recombivax HB (Hepatitis B vaccine (Recombinant) [Hep B]) at 2, 4, 6, and 12–15 months of age. In this study, conducted between December

1995 and March 1999, parents of study participants were asked to bring their children to the study clinics if the child had respiratory infections or symptoms suggesting acute otitis media (AOM). If AOM was diagnosed, tympanocentesis was performed, and the middle-ear fluid was cultured. If *S. pneumoniae* was isolated, serotyping was performed; the primary endpoint was efficacy against AOM episodes caused by vaccine serotypes in the per-protocol population. In the NCKP trial, the efficacy of Prevnar against otitis media was assessed from the beginning of the trial in October 1995 through April 1998. The otitis media analysis included 34,146 infants randomized to receive either Prevnar (N=17,070), or the control vaccine (N=17,076), at 2, 4, 6, and 12–15 months of age. In this trial, no routine tympanocentesis was performed, and no standard definition of otitis media was used by study physicians. The primary otitis media endpoint was efficacy against all otitis media episodes in the per-protocol population.

The vaccine efficacy against AOM episodes due to vaccine serotypes assessed in the Finnish trial, was 57% (95% CI: 44%, 67%) in the per-protocol population and 54% (95% CI: 41%, 64%) in the intent-to-treat population. The vaccine efficacy against AOM episodes due to vaccine-related serotypes (6A, 9N, 18B, 19A, 23A), also assessed in the Finnish trial, was 51% (95% CI: 27, 67) in the per-protocol population and 44% (95% CI: 20, 62) in the intent-to-treat population. There was a nonsignificant increase in AOM episodes caused by serotypes unrelated to the vaccine in the per-protocol population, compared to children who received the control vaccine, suggesting that children who received Prevnar appeared to be at increased risk of otitis media due to pneumococcal serotypes not represented in the vaccine. However, vaccination with Prevnar reduced pneumococcal otitis media episodes overall. In the NCKP trial, in which the endpoint was all otitis media episodes regardless of etiology, vaccine efficacy was 7% (95% CI: 4%, 10%) and 6% (95% CI: 4%, 9%), respectively, in the per-protocol and intent-to-treat analyses. Several other otitis media endpoints were also assessed in the two trials.

Recurrent AOM, defined as 3 episodes in 6 months or 4 episodes in 12 months, was reduced by 9% in both the per-protocol and intent-to-treat populations (95% CI: 3%, 15% in per-protocol and 95% CI: 4%, 14% in intent-to-treat) in the NCKP trial; a similar trend was observed in the Finnish trial. The NCKP trial also demonstrated a 20% reduction (95% CI: 2, 35) in the placement of tympanostomy tubes in the per-protocol population and a 21% reduction (95% CI: 4, 34) in the intent-to-treat population. Data from the NCKP trial accumulated through an extended follow-up period to April 20, 1999, in which a total of 37,866 children were included (18,925 in Prevnar group and 18,941 in MnCC control group), resulted in similar otitis media efficacy estimates for all endpoints.

Prevnar 13 Adult Efficacy Data

The efficacy of Prevnar 13 against vaccine-type (VT) pneumococcal community-acquired pneumonia (CAP) and IPD was assessed in a randomized, double-blind, placebo-controlled study conducted over ~ 4 years in the Netherlands¹² (Study 12). A total of 84,496 subjects 65 years and older received a single dose of either Prevnar 13 or placebo in a 1:1 randomization; 42,240 subjects were vaccinated with Prevnar 13 and 42,256 subjects were vaccinated with placebo.

The primary objective was to demonstrate the efficacy of Prevnar 13 in the prevention of a first episode of confirmed VT-CAP (defined as presence of ≥ 2 specified clinical criteria; chest X-ray consistent with CAP as determined by a central committee of radiologists; and positive VT-specific Urinary Antigen Detection assay (UAD) or isolation of VT *S. pneumoniae* from blood or other sterile site). The secondary objectives were to demonstrate the efficacy of Prevnar 13 in the prevention of a first episode of 1) confirmed nonbacteremic/noninvasive (NB/NI) VT-CAP (an episode of VT-CAP for which the blood culture result and any other sterile site culture results were negative for *S. pneumoniae*) and 2) VT-IPD (the presence of *S. pneumoniae* in a sterile site).

Surveillance for suspected pneumonia and IPD began immediately after vaccination and continued through identification of a prespecified number of cases. Subjects who had a CAP or IPD episode with symptom onset less than 14 days after vaccination were excluded from all analyses.

The median duration of follow-up per subject was 3.93 years. Prevnar 13 demonstrated statistically significant vaccine efficacy (VE) in preventing first episodes of VT pneumococcal CAP,

nonbacteremic/noninvasive (NB/NI) VT pneumococcal CAP, and VT-IPD (Table 15).

Table 15 - Vaccine Efficacy for the Primary and Secondary Efficacy Endpoints – Per-Protocol Population

		Vaccine Group			
		Prevnar 13	Placebo		
		N=42240	N=42256		
Efficacy Endpoint	Total Number of Episodes	n	n	VE (%)	(95.2% CI)
Primary endpoint: First case of confirmed VT pneumococcal CAP	139	49	90	45.6	(21.8, 62.5)
Secondary endpoint: First episode of confirmed NB/NI VT pneumococcal CAP	93	33	60	45	(14.2, 65.3)
Secondary endpoint: First episode of VT-IPD	35	7	28	75	(41.1, 90.9)

Abbreviations: CAP = community-acquired pneumonia; CI = confidence interval; NB/NI = nonbacteremic/noninvasive; IPD = invasive pneumococcal disease; VE = vaccine efficacy; VT = vaccine-type.

14.2 Prevnar 13 Clinical Trials in Children 6 Weeks Through 17 Years of Age

Infants and Children 6 Weeks Through 17 Months of Age

Prevnar 13 effectiveness against invasive pneumococcal disease was inferred from comparative studies to a US-licensed 7-valent pneumococcal conjugate vaccine, Prevnar, in which Prevnar 13 elicited antipolysaccharide binding and functional OPA antibodies, as measured by ELISA and dOPA assays, respectively. These studies were designed to evaluate immunologic noninferiority of Prevnar 13 to Prevnar.

Clinical trials have been conducted in the US using a 2, 4, 6, and 12–15 month schedule.

The US noninferiority study² (Study 2) was a randomized, double-blind, active-controlled trial in which 2 month-old infants were randomly assigned to receive either Prevnar 13 or Prevnar in a 1:1 ratio. The two vaccine groups were well balanced with respect to race, ethnicity, and age and weight at enrollment. Most subjects were White (69.1%), 19.6% were Black or African-American, and 2.4% were Asian; 82.1% of subjects were non-Hispanic and non-Latino and 17.3% were Hispanic or Latino. Overall, 54.0% of subjects were male infants.

In Study 2, immune responses were compared in subjects receiving either Prevnar 13 or Prevnar using a set of noninferiority criteria. Co-primary endpoints included the percentage of subjects with serum pneumococcal anti-capsular polysaccharide IgG ≥ 0.35 $\mu\text{g/mL}$ measured one month after the third dose and serum pneumococcal anti-capsular polysaccharide IgG geometric mean concentrations (GMCs) one month after the fourth dose. The assay used for this determination was a standardized ELISA involving pre-absorption of the test sera with pneumococcal C-polysaccharide and serotype 22F polysaccharide to reduce non-specific background reactivity. Responses to the 7 common serotypes in Prevnar 13 and Prevnar recipients were compared directly. Responses to the 6 additional serotypes in Prevnar 13 recipients were each compared to the lowest response observed among the Prevnar serotypes in Prevnar recipients.

Pneumococcal Immune Responses Following Three Doses

In Study 2, the noninferiority criterion for the proportion of subjects with pneumococcal anti-capsular polysaccharide IgG antibody concentrations ≥ 0.35 $\mu\text{g/mL}$ one month after the third dose was met for 10 of the 13 serotypes. The exceptions were serotypes 6B, 9V, and 3. Although the response to serotypes 6B and 9V did not meet the pre-specified noninferiority criterion, the differences were marginal.

The percentage of infants achieving pneumococcal anti-capsular polysaccharide IgG antibody concentrations ≥ 0.35 $\mu\text{g/mL}$ one month after the third dose is shown below (Table 16).

Table 16: Percentage of Subjects With Anti-capsular Antibody Concentration ≥ 0.35 $\mu\text{g/mL}$ One Month After a Three Dose Series Administered at 2, 4 and 6 Months of Age, Study 2^{*,†,‡,§}

Serotype	Pevnar 13 N=249–252 (95% CI)	Pevnar N=250–252 (95% CI)	Difference in % responders (95% CI)
Pevnar Serotypes			
4	94.4 (90.9, 96.9)	98.0 (95.4, 99.4)	-3.6 (-7.3, -0.1)
6B	87.3 (82.5, 91.1)	92.8 (88.9, 95.7)	-5.5 (-10.9, -0.1)
9V	90.5 (86.2, 93.8)	98.4 (96.0, 99.6)	-7.9 (-12.4, -4.0)
14	97.6 (94.9, 99.1)	97.2 (94.4, 98.9)	0.4 (-2.7, 3.5)
18C	96.8 (93.8, 98.6)	98.4 (96.0, 99.6)	-1.6 (-4.7, 1.2)
19F	98.0 (95.4, 99.4)	97.6 (99.4, 99.1)	0.4 (-2.4, 3.4)
23F	90.5 (86.2, 93.8)	94.0 (90.4, 96.6)	-3.6 (-8.5, 1.2)
Additional Serotypes [¶]			
1	95.6 (92.3, 97.8)	¶	2.8 (-1.3, 7.2)
3	63.5 (57.1, 69.4)	¶	-29.3 (-36.2, -22.4)
5	89.7 (85.2, 93.1)	¶	-3.1 (-8.3, 1.9)
6A	96.0 (92.8, 98.1)	¶	3.2 (-0.8, 7.6)
7F	98.4 (96.0, 99.6)	¶	5.6 (1.9, 9.7)
19A	98.4 (96.0, 99.6)	¶	5.6 (1.9, 9.7)

* Studies conducted in US NCT00373958 (Study 2).

† Evaluable Immunogenicity Population.

‡ Noninferiority was met when the lower limit of the 95% CI for the difference between groups (Pevnar 13 minus Pevnar) was greater than -10%.

§ Antibody measured by a standardized ELISA involving pre-absorption of the test sera with pneumococcal C-polysaccharide and serotype 22F polysaccharide to reduce non-specific background reactivity.

¶ Comparison for the 6 additional serotypes was to the lowest responder of the 7 common serotypes in Pevnar recipients, which for this analysis was serotype 6B (92.8%; 95% CI: 88.9, 95.7).

Functional dOPA antibody responses were elicited for all 13 serotypes, as shown in Table 17.

Table 17: Pneumococcal dOPA Antibody Geometric Mean Titers One Month After a Three Dose Series Administered at 2, 4 and 6 Months of Age, Study 2^{*,†,‡}

Serotype	Pevnar 13 N=91–94 (95% CI)	Pevnar N=89–94 (95% CI)
Pevnar Serotypes		
4	359 (276, 468)	536 (421, 681)
6B	1055 (817, 1361)	1514 (1207, 1899)
9V	4035 (2933, 5553)	3259 (2288, 4641)

14	1240 (935, 1646)	1481 (1133, 1934)
18C	276 (210, 361)	376 (292, 484)
19F	54 (40, 74)	45 (34, 60)
23F	791 (605, 1034)	924 (709, 1204)
Additional Serotypes		
1	52 (39, 69)	4 (4, 5)
3	121 (92, 158)	7 (5, 9)
5	91 (67, 123)	4 (4, 4)
6A	980 (783, 1226)	100 (66, 152)
7F	9494 (7339, 12281)	128 (80, 206)
19A	152 (105, 220)	7 (5, 9)

* Studies conducted in US NCT00373958 (Study 2).

† The dOPA (opsonophagocytic activity) antibody assay measures the ability of immune sera, in conjunction with complement, to mediate the uptake and killing of *S. pneumoniae* by phagocytic cells.

‡ Evaluable Immunogenicity Population.

Pneumococcal Immune Responses Following Four Doses

In Study 2, post-dose 4 antibody concentrations were higher for all 13 serotypes than those achieved after the third dose. The noninferiority criterion for pneumococcal anti-capsular polysaccharide GMCs after 4 doses was met for 12 of the 13 pneumococcal serotypes. The noninferiority criterion was not met for the response to serotype 3 (Table 18).

Table 18: Pneumococcal IgG GMCs (µg/mL) One Month After a Four Dose Series Administered at 2, 4, 6 and 12–15 Months, Study 2^{*,†,‡,§}

Serotype	Pprevnar 13 N=232–236 (95% CI)	Pprevnar N=222–223 (95% CI)	GMC Ratio (95% CI)
Pprevnar Serotypes			
4	3.73 (3.28, 4.24)	5.49 (4.91, 6.13)	0.68 (0.57, 0.80)
6B	11.53 (9.99, 13.30)	15.63 (13.80, 17.69)	0.74 (0.61, 0.89)
9V	2.62 (2.34, 2.94)	3.63 (3.25, 4.05)	0.72 (0.62, 0.85)
14	9.11 (7.95, 10.45)	12.72 (11.22, 14.41)	0.72 (0.60, 0.86)
18C	3.20 (2.82, 3.64)	4.70 (4.18, 5.28)	0.68 (0.57, 0.81)
19F	6.60 (5.85, 7.44)	5.60 (4.87, 6.43)	1.18 (0.98, 1.41)
23F	5.07 (4.41, 5.83)	7.84 (6.91, 8.90)	0.65 (0.54, 0.78)
Additional Serotypes [¶]			
1	5.06 (4.43, 5.80)	¶	1.40 (1.17, 1.66)
3	0.94 (0.83, 1.05)	¶	0.26 (0.22, 0.30)
5	3.72 (3.31, 4.18)	¶	1.03 (0.87, 1.20)
6A	8.20 (7.30, 9.20)	¶	2.26 (1.93, 2.65)
7F	5.67 (5.01, 6.42)	¶	1.56 (1.32, 1.85)
19A	8.55 (7.64, 9.56)	¶	2.36 (2.01, 2.76)

* Studies conducted in US NCT00373958 (Study 2).

† Evaluable Immunogenicity Population.

‡ Noninferiority was declared if the lower limit of the 2-sided 95% CI for Geometric Mean Ratio (Pprevnar 13:Pprevnar) was greater than 0.5.

§ Antibody measured by a standardized ELISA involving pre-absorption of the test sera with pneumococcal C-polysaccharide and serotype 22F polysaccharide to reduce non-specific background

reactivity.

¶ Comparison for the 6 additional serotypes was to the lowest responder of the 7 common serotypes in Prevnar recipients, which for this analysis was serotype 9V (3.63; 95% CI 3.25, 4.05).

Following the fourth dose, the functional dOPA antibody response for each serotype was quantitatively greater than the response following the third dose (see Table 19).

Table 19: Pneumococcal dOPA Antibody Geometric Mean Titers One Month After the Fourth Dose-Evaluable Toddler Immunogenicity Population, Study 2^{*,†}

Serotype	Prevnar 13 N=88–92 (95% CI)	Prevnar N=92–96 (95% CI)
Prevnar Serotypes		
4	1180 (847, 1643)	1492 (1114, 1999)
6B	3100 (2337, 4111)	4066 (3243, 5098)
9V	11856 (8810, 15955)	18032 (14125, 23021)
14	2002 (1453, 2760)	2366 (1871, 2992)
18C	993 (754, 1308)	1722 (1327, 2236)
19F	200 (144, 276)	167 (121, 230)
23F	2723 (1961, 3782)	4982 (3886, 6387)
Additional Serotypes		
1	164 (114, 237)	5 (4, 6)
3	380 (300, 482)	12 (9, 16)
5	300 (229, 393)	5 (4, 6)
6A	2242 (1707, 2945)	539 (375, 774)
7F	11629 (9054, 14938)	268 (164, 436)
19A	1024 (774, 1355)	29 (19, 44)

* Studies conducted in US NCT00373958 (Study 2).

† The dOPA (opsonophagocytic activity) antibody assay measures the ability of immune sera, in conjunction with complement, to mediate the uptake and killing of *S. pneumoniae* by phagocytic cells.

Previously Unvaccinated Older Infants and Children 7 Months Through 5 Years of Age

In an open-label descriptive study of Prevnar 13 in Poland⁴ (Study 4), children 7 months through 11 months of age, 12 months through 23 months of age and 24 months through 5 years of age (prior to the 6th birthday) who were naïve to pneumococcal conjugate vaccine, were given 3, 2 or 1 dose of Prevnar 13 respectively, according to the age-appropriate schedules in Table 2. Serum IgG concentrations were measured one month after the final dose in each age group and the data are shown in Table 20.

Table 20: Pneumococcal Anti-capsular Polysaccharide IgG Antibody Geometric Mean Concentrations (µg/mL) One Month After the Final Prevnar 13 Catch-Up Dose in Pneumococcal Vaccine Naïve Children 7 Months Through 5 Years of Age by Age Group, Study 4^{*,†}

Serotype	3 doses Prevnar 13 7 through 11 months N=83–84 (95% CI)	2 doses Prevnar 13 12 through 23 months N=104–110 (95% CI)	1 dose Prevnar 13 24 months through 5 years N=135–152 (95% CI)
1	2.88 (2.44, 3.39)	2.74 (2.37, 3.16)	1.78 (1.52, 2.08)

3	1.94 (1.68, 2.24)	1.86 (1.60, 2.15)	1.42 (1.23, 1.64)
4	3.63 (3.11, 4.23)	4.28 (3.78, 4.86)	3.37 (2.95, 3.85)
5	2.85 (2.34, 3.46)	2.16 (1.89, 2.47)	2.33 (2.05, 2.64)
6A	3.72 (3.12, 4.45)	2.62 (2.25, 3.06)	2.96 (2.52, 3.47)
6B	4.77 (3.90, 5.84)	3.38 (2.81, 4.06)	3.41 (2.80, 4.16)
7F	5.30 (4.54, 6.18)	5.99 (5.40, 6.65)	4.92 (4.26, 5.68)
9V	2.56 (2.21, 2.96)	3.08 (2.69, 3.53)	2.67 (2.32, 3.07)
14	8.04 (6.95, 9.30)	6.45 (5.48, 7.59)	2.24 (1.71, 2.93)
18C	2.77 (2.39, 3.23)	3.71 (3.29, 4.19)	2.56 (2.17, 3.03)
19A	4.77 (4.28, 5.33)	4.94 (4.31, 5.65)	6.03 (5.22, 6.97)
19F	2.88 (2.35, 3.54)	3.07 (2.68, 3.51)	2.53 (2.14, 2.99)
23F	2.16 (1.82, 2.55)	1.98 (1.64, 2.39)	1.55 (1.31, 1.85)

Note – ClinicalTrials.gov NCT number is as follows: NCT00452452 (Poland).

* Studies conducted in Poland NCT00452452 (Study 4).

† Open label administration of Pevnar 13.

Children 15 Months Through 59 Months of Age Previously Vaccinated with Pevnar

In an open-label descriptive study in the US⁵ (Study 5), children 15 months through 59 months previously vaccinated with 3 or 4 doses of Pevnar, received 2 doses of Pevnar 13 (children >15 through 23 months of age) or 1 dose of Pevnar 13 (children 24 months through 59 months of age). The data following one dose of Pevnar 13 in children 24 months through 59 months of age are shown in Table 21.

Table 21: Pneumococcal Anti-capsular Polysaccharide IgG Antibody Geometric Mean Concentrations (µg/mL) One Month After One Pevnar 13 Catch-Up Dose in Children 24 Through 59 Months of Age With 3 or 4 Prior Doses of Pevnar, US Catch-Up Study 5*,†

Serotype	1 dose Pevnar 13 24 months through 59 months N=173–175 (95% CI)
	1
3	1.38 (1.17, 1.61)
5	2.13 (1.89, 2.41)
6A	12.96 (11.04, 15.21)
7F	4.22 (3.74, 4.77)
19A	14.18 (12.37, 16.25)

* Studies conducted in US NCT00761631 (Study 5).

† Open label administration of Pevnar 13.

Children 5 Through 17 Years of Age

In a US study⁵ (Study 5), a single dose of Pevnar 13 was administered to children 5 through 9 years of age, who were previously vaccinated with at least one dose of Pevnar, and to pneumococcal vaccine-naïve children 10 through 17 years of age.

In children 5 through 9 years of age, serotype-specific IgG concentrations measured 1 month after vaccination were noninferior (i.e., the lower limit of the 2-sided 95% CI for the geometric mean ratio [GMR] of >0.5) to the corresponding IgG concentrations in toddlers (Study 3) 1 month after a fourth pneumococcal vaccination (after the 4th dose of Pevnar for the 7 common serotypes and after the 4th

dose of Prevnar 13 for the 6 additional serotypes) as shown in Tables 22 and 23 respectively.

Table 22: Pneumococcal IgG GMCs ($\mu\text{g/mL}$) One Month After Vaccination for 7 Common Serotypes, Prevnar 13 in Children 5 through 9 Years of Age in Study 5 Relative to Prevnar in Study 3 (Post-toddler)*,†,‡

Serotype	Vaccine Group (as Enrolled/Randomized)							
	Prevnar 13 5 Through 9 Years (Study 5)			Prevnar Post-Toddler Dose (Study 3)			GMC Ratio ^D	(95% CI ^B)
	n [§]	GMC [¶]	(95% CI [#])	n [§]	GMC [¶]	(95% CI [#])		
Common								
4	169	8.45	(7.24, 9.87)	173	2.79	(2.45, 3.18)	3.03	(2.48, 3.71)
6B	171	53.56	(45.48, 63.07)	173	9.47	(8.26, 10.86)	5.66	(4.57, 6.99)
9V	171	9.51	(8.38, 10.78)	172	1.97	(1.77, 2.19)	4.83	(4.10, 5.70)
14	169	29.36	(24.78, 34.78)	173	8.19	(7.31, 9.18)	3.58	(2.93, 4.39)
18C	171	8.23	(7.13, 9.51)	173	2.33	(2.05, 2.65)	3.53	(2.91, 4.29)
19F	171	17.58	(14.95, 20.67)	173	3.31	(2.87, 3.81)	5.31	(4.29, 6.58)
23F	169	11.26	(9.79, 12.95)	173	4.49	(3.86, 5.23)	2.51	(2.04, 3.08)

* Studies conducted in US NCT00761631 (Study 5) and NCT00444457 (Study 3).

† Evaluable Immunogenicity Population.

‡ Noninferiority was declared if the lower limit of the 2-sided 95% CI for geometric mean ratio was greater than 0.5.

§ n = Number of subjects with a determinate antibody concentration for the specified serotype.

¶ Geometric mean concentrations (GMCs) were calculated using all subjects with available data for the specified blood draw. GMC after a 4-dose vaccination series with Prevnar (Study 3, post-toddler).

Confidence intervals (CIs) are back transformations of a confidence interval based on the Student t distribution for the mean logarithm of the concentrations.

^D Ratio of GMCs: Prevnar 13 (Study 5) to Prevnar (Study 3) reference.

^B CIs for the ratio are back transformations of a confidence interval based on the Student t distribution for the mean difference of the logarithms of the measures [Prevnar 13 (Study 5) – Prevnar (Study 3)].

Table 23: Pneumococcal IgG GMCs ($\mu\text{g/mL}$) One Month After Vaccination for Additional 6 Serotypes, Prevnar 13 in Children 5 through 9 Years of Age in Study 5 Relative to Prevnar 13 in Study 3 (Post-toddler)*,†,‡

Serotype	Vaccine Group (as Enrolled/Randomized)							
	Prevnar 13 5 Through 9 Years (Study 5)			Prevnar 13 Post-Toddler Dose (Study 3)			GMC Ratio ^D	(95% CI ^B)
	n [§]	GMC [¶]	(95% CI [#])	n [§]	GMC [¶]	(95% CI [#])		
Additional								
1	171	3.57	(3.05, 4.18)	1068	2.90	(2.75, 3.05)	1.23	(1.07, 1.42)
3	171	2.38	(2.07, 2.74)	1065	0.75	(0.72, 0.79)	3.17	(2.78, 3.62)
5	171	5.52	(4.82, 6.32)	1068	2.85	(2.72, 2.98)	1.94	(1.71, 2.20)
6A	169	21.51	(18.15, 25.51)	1063	7.11	(6.78, 7.46)	3.03	(2.64, 3.47)
7F	170	6.24	(5.49, 7.08)	1067	4.39	(4.18, 4.61)	1.42	(1.24, 1.62)
19A	170	17.18	(15.01, 19.67)	1056	8.44	(8.05, 8.86)	2.03	(1.78, 2.32)

- * Studies conducted in US NCT00761631 (Study 5) and NCT00444457 (Study 3).
- † Evaluable Immunogenicity Population.
- ‡ Noninferiority was declared if the lower limit of the 2-sided 95% CI for geometric mean ratio was greater than 0.5.
- § n = Number of subjects with a determinate antibody concentration for the specified serotype.
- ¶ Geometric mean concentrations (GMCs) were calculated using all subjects with available data for the specified blood draw. GMC after a 4-dose vaccination series with Prevnar 13 (Study 3, post-toddler).
- # Confidence intervals (CIs) are back transformations of a confidence interval based on the Student t distribution for the mean logarithm of the concentrations.
- ᵀ Ratio of GMCs: Prevnar 13 (Study 5) to Prevnar 13 (Study 3).
- ᵆ CIs for the ratio are back transformations of a confidence interval based on the Student t distribution for the mean difference of the logarithms of the measures [Prevnar 13 (Study 5) – Prevnar 13 (Study 3)].

In children 10 through 17 years of age OPA GMTs, as measured by the mcOPA assay, 1 month after vaccination were noninferior (i.e., the lower limit of the 2-sided 95% CI for the GMR of >0.5) to mcOPA GMTs in the 5 through 9 year old group for 12 of 13 serotypes (except for serotype 3), as shown in Table 24.

Table 24: Comparison of Pneumococcal mcOPA GMTs One Month After Vaccination, Prevnar 13, in Children 10 through 17 Years of Age Relative to Prevnar 13 in Children 5 through 9 Years of Age^{*,†,‡,§}

Serotype	Vaccine Group (as Enrolled)						GMT Ratio ^ᵆ	(95% CI ^à)
	Prevnar 13 (10 through 17 Years)			Prevnar 13 (5 through 9 Years)				
	n [¶]	GMT [#]	(95% CI ^ᵀ)	n [¶]	GMT [#]	(95% CI ^ᵀ)		
Common								
4	188	6912	(6101, 7831)	181	4629	(4017, 5334)	1.5	(1.24, 1.80)
6B	183	14224	(12316, 16427)	178	14996	(13164, 17083)	0.9	(0.78, 1.15)
9V	186	4485	(4001, 5028)	180	4733	(4203, 5328)	0.9	(0.80, 1.12)
14	187	6894	(6028, 7884)	176	4759	(4120, 5497)	1.4	(1.19, 1.76)
18C	182	6263	(5436, 7215)	175	8815	(7738, 10041)	0.7	(0.59, 0.86)
19F	184	2280	(1949, 2668)	178	1591	(1336, 1893)	1.4	(1.14, 1.81)
23F	187	3808	(3355, 4323)	176	3245	(2819, 3736)	1.2	(0.97, 1.42)
Additional								
1	189	322	(275, 378)	179	191	(165, 221)	1.7	(1.36, 2.10)
3	181	114	(101, 130)	178	203	(182, 226)	0.6	(0.48, 0.67)
5	183	360	(298, 436)	178	498	(437, 568)	0.7	(0.57, 0.91)
6A	182	9928	(8457, 11655)	178	7514	(6351, 8891)	1.3	(1.05, 1.67)
7F	185	6584	(5829, 7436)	178	10334	(9099, 11737)	0.6	(0.53, 0.76)
19A	187	1276	(1132, 1439)	180	1180	(1048, 1329)	1.1	(0.91, 1.28)

- * Studies conducted in US NCT00761631 (Study 5).
- † Evaluable Immunogenicity Population.
- ‡ Noninferiority was declared if the lower limit of the 2-sided 95% CI for geometric mean ratio was greater than 0.5.
- § Individual mcOPA antibody assay values below the assay LLOQ (lower limit of quantitation) were set

at 0.50*LLOQ for the purpose of calculating the mcOPA antibody GMT.

- ¶ n= Number of subjects with a determinate antibody titer for the specified serotype.
- # Geometric mean titers (GMTs) were calculated using all subjects with available data for the specified blood draw.
- ▷ Confidence intervals (CIs) are back transformations of a confidence interval based on the Student t distribution for the mean logarithm of the titers.
- β Ratio of GMTs: Prevnar 13(10 through 17 years of age) to Prevnar 13 (5 through 9 years of age).
- à CIs for the ratio are back transformations of a confidence interval based on the Student t distribution for the mean difference of the logarithms of the measures [Prevnar 13(10 through 17 years of age) – Prevnar 13(5 through 9 years of age)] Study 5.

14.3 Prevnar 13 Immunogenicity Clinical Trials in Adults

Six Phase 3 or Phase 4 clinical trials^{6-8,10,11,13} were conducted in the US and Europe evaluating the immunogenicity of Prevnar 13 in different adult age groups, in individuals who were either not previously vaccinated with PPSV23 (PPSV23 unvaccinated) or who had received one dose of PPSV23 (PPSV23 previously vaccinated).

Each study included healthy adults and immunocompetent adults with stable underlying conditions including chronic cardiovascular disease, chronic pulmonary disease, renal disorders, diabetes mellitus, chronic liver disease, and medical risk conditions and behaviors (e.g., alcoholism and smoking) that are known to increase the risk of serious pneumococcal pneumonia and invasive pneumococcal disease. A stable medical condition was defined as a medical condition not requiring significant change in therapy (i.e., change to new therapy category due to worsening disease) or hospitalization for worsening disease 6–12 weeks prior to receipt of the study vaccine.

Immune responses elicited by Prevnar 13 and PPSV23 were measured by a mcOPA antibody assay for the 13 pneumococcal serotypes contained in Prevnar 13. Serotype-specific mcOPA antibody GMTs measured 1 month after each vaccination were calculated. For the 12 serotypes in common to both vaccines, noninferiority between vaccines was met if the lower limit of the 2-sided 95% confidence interval (CI) of the GMT ratio (Prevnar 13/PPSV23) was greater than 0.5.

The response to the additional serotype 6A, which is contained in Prevnar 13 but not in PPSV23, was assessed by demonstration of a ≥ 4 -fold increase in the anti-6A mcOPA antibody titer above preimmunization levels. A statistically significantly greater response for Prevnar 13 was defined, for the difference in percentages (Prevnar 13 minus PPSV23) of adults achieving a ≥ 4 -fold increase in anti-6A mcOPA antibody titer, as the lower limit of the 2-sided 95% CI greater than zero. For comparison of mcOPA antibody GMTs, a statistically greater response for serotype 6A was defined as the lower limit of the 2-sided 95% CI of the GMT ratio (Prevnar 13/PPSV23) greater than 2.

Of the 6 Phase 3 or Phase 4 clinical trials, 2 noninferiority trials^{6,7} were conducted in which the immune responses to Prevnar 13 were compared with the immune responses to PPSV23; one in PPSV23 unvaccinated adults aged 18 through 64 years⁶ (Study 6), and one in PPSV23 previously vaccinated adults aged ≥ 70 years⁷ (Study 7). A third study compared immune responses to a single dose of Prevnar 13 to the response to Prevnar 13 administered one year after a dose of PPSV23 in adults aged 60 through 64 years who were PPSV23 unvaccinated at enrollment⁸ (Study 8). The study also compared immune responses of PPSV23 as a single dose to the responses to PPSV23 administered one year after a dose of Prevnar 13. Two studies assessed the concomitant administration of Prevnar 13 with seasonal inactivated Fluarix (IIV3) in the US¹⁰ (Study 10) and Europe¹¹ (Study 11). One study (Study 13) assessed the concomitant administration of Prevnar 13 with seasonal inactivated Fluzone Quadrivalent (IIV4) in PPSV23 previously vaccinated adults ≥ 50 years of age in the US.

Overall across the clinical studies evaluating the immunogenicity of Prevnar 13 in adults, persons 18 through 64 years of age responded at least as well as persons 65 years and older, the age group evaluated in a clinical endpoint efficacy trial.

Clinical Trials Conducted in PPSV23 Unvaccinated Adults

In an active-controlled modified¹ double-blind clinical trial⁶ (Study 6) of Prevnar 13 in the US, PPSV23 unvaccinated adults aged 60 through 64 years were randomly assigned (1:1) to receive Prevnar 13 or PPSV23. In addition, adults aged 18 through 49 years and 50 through 59 years were enrolled and received one dose of Prevnar 13 (open-label).

In adults aged 60 through 64 years, the mcOPA antibody GMTs elicited by Prevnar 13 were noninferior to those elicited by PPSV23 for the 12 serotypes in common to both vaccines (see Table 24). In addition, the lower limit of the 95% confidence interval for the mcOPA antibody GMT ratio (Prevnar 13/PPSV23) was greater than 1 for 8 of the serotypes in common.

For serotype 6A, which is unique to Prevnar 13, the proportion of subjects with a ≥ 4 -fold increase after Prevnar 13 (88.5%) was statistically significantly greater than after PPSV23 (49.3%) in PPSV23-unvaccinated adults aged 60 through 64 years. OPA antibody GMTs for serotype 6A were statistically significantly greater after Prevnar 13 compared with after PPSV23 (see Table 25).

The mcOPA antibody GMTs elicited by Prevnar 13 in adults aged 50 through 59 years were noninferior to the corresponding mcOPA antibody GMTs elicited by Prevnar 13 in adults aged 60 through 64 years for all 13 serotypes (see Table 25).

In adults aged 18 through 49 years, the mcOPA antibody GMTs elicited by Prevnar 13 were noninferior to those elicited by Prevnar 13 in adults aged 60 through 64 years for all 13 serotypes (see Table 25).

Table 25: mcOPA Antibody GMTs in PPSV23-Unvaccinated Adults Aged 18 Through 49 Years or Aged 50 Through 59 Years Given Prevnar 13 and in Adults Aged 60 Through 64 Years Given Prevnar 13 or PPSV23 (Study 6)^{*,†,‡,§,¶}

Serotype	Prevnar 13	Prevnar 13	Prevnar 13	PPSV23	Prevnar 13	Prevnar 13	Prevnar 13
	18–49 Years ^p N=836–866	50–59 Years ^p N=350–384	60–64 Years N=359–404	60–64 Years N=367–402	18–49 Relative to 60–64 Years	50–59 Relative to 60–64 Years	Relative to PPSV23, 60–64 Years [#]
	GMT	GMT	GMT	GMT	GMT Ratio (95% CI)	GMT Ratio (95% CI)	GMT Ratio (95% CI)
1	353	211	158	119	2.4 (2.03, 2.87)	1.3 (1.07, 1.65)	1.3 (1.07, 1.65)
3	91	94	96	90	1.0 (0.84, 1.13)	1.0 (0.82, 1.18)	1.1 (0.89, 1.29)
4	4747	2904	2164	1405	2.3 (1.92, 2.76)	1.3 (1.06, 1.70)	1.5 (1.18, 2.00)
5	386	322	236	198	1.9 (1.55, 2.42)	1.4 (1.08, 1.74)	1.2 (0.95, 1.50)
6A ^β	5746	4469	2766	343	2.2 (1.84, 2.67)	1.6 (1.28, 2.03)	8.1 (6.11, 10.67)
6B	9813	3350	2212	998	4.9 (4.13, 5.93)	1.5 (1.20, 1.91)	2.2 (1.70, 2.89)

7F	3249	1807	1535	829	2.9 (2.41, 3.49)	1.2 (0.98, 1.41)	1.9 (1.52, 2.26)
9V	3339	2190	1701	1012	2.9 (2.34, 3.52)	1.3 (1.08, 1.53)	1.7 (1.40, 2.02)
14	2983	1078	733	819	4.9 (4.01, 5.93)	1.5 (1.14, 1.89)	0.9 (0.69, 1.16)
18C	3989	2077	1834	1074	2.3 (1.91, 2.79)	1.1 (0.89, 1.44)	1.7 (1.32, 2.21)
19A	1580	968	691	368	2.3 (2.02, 2.66)	1.4 (1.17, 1.68)	1.9 (1.53, 2.30)
19F	1533	697	622	636	3.0 (2.44, 3.60)	1.1 (0.89, 1.41)	1.0 (0.78, 1.23)
23F	1570	531	404	87	4.2 (3.31, 5.31)	1.3 (0.96, 1.80)	4.6 (3.37, 6.38)

GMT, Geometric Mean Titer.

* Study conducted in US NCT00427895 (Study 6).

† Noninferiority was defined for the 13 serotypes in adults aged 18 to 49 years, for the 12 common serotypes in adults aged 60 to 64 years and for the 13 serotypes in adults aged 50 to 59 years as the lower limit of the 2-sided 95% CI for GMT ratio greater than 0.5.

‡ mcOPA antibody for the 11 serotypes unique to PPSV23 but not contained in Prevnar 13 were not measured.

§ Individual mcOPA antibody assay values below the assay LLOQ (lower limit of quantitation) were set at 0.50*LLOQ for the purpose of calculating the mcOPA antibody GMT.

¶ Evaluable Immunogenicity Population.

For serotype 6A, which is unique to Prevnar 13, a statistically significantly greater response was defined for analysis in cohort 1 as the lower limit of the 2-sided 95% CI for the GMT ratio (Prevnar 13/PPSV23) greater than 2.

▷ Open label administration of Prevnar 13.

⋈ 6A is a serotype unique to Prevnar 13 but not contained in PPSV23.

¹ Modified double-blind means that the site staff dispensing and administering the vaccine were unblinded, but all other study personnel including the principal investigator and subject were blinded.

Clinical Trials Conducted in PPSV23 Previously Vaccinated Adults

In a Phase 3 active-controlled, modified double-blind clinical trial⁷ (Study 7) of Prevnar 13 in the US and Sweden, PPSV23 previously vaccinated adults aged ≥70 years who had received one dose of PPSV23 ≥5 years prior were randomly assigned (1:1) to receive either Prevnar 13 or PPSV23.

The mcOPA antibody GMTs elicited by Prevnar 13 were noninferior to those elicited by PPSV23 for the 12 serotypes in common, when Prevnar 13 or PPSV23 were administered at a minimum of 5 years after a prior dose of PPSV23. In addition, the lower limit of the 95% confidence interval for the mcOPA antibody GMT ratio (Prevnar 13/PPSV23) was greater than 1 for 9 of the serotypes in common.

For serotype 6A, which is unique to Prevnar 13, the proportion of subjects with a ≥4-fold increase in mcOPA antibody titers after Prevnar 13 (71.1%) was statistically significantly greater than after PPSV23 (27.3%) in PPSV23 previously vaccinated adults aged ≥70 years. mcOPA antibody GMTs for serotype

6A were statistically significantly greater after Prevnar 13 compared with after PPSV23.

This clinical trial demonstrated that in adults aged ≥ 70 years and previously vaccinated with PPSV23 ≥ 5 years prior, vaccination with Prevnar 13 elicited noninferior immune responses as compared with re-vaccination with PPSV23 (see Table 26).

Table 26: mcOPA Antibody GMTs in PPSV23-Previously Vaccinated Adults Aged ≥ 70 Years Given Prevnar 13 or PPSV23 (Study 7)*,†,‡,§,¶,#

Serotype	Prevnar 13 N=400–426 GMT	PPSV23 N=395–445 GMT	Prevnar 13 Relative to PPSV23	
			GMT Ratio	(95% CI)
1	93	66	1.4	(1.14, 1.72)
3	59	53	1.1	(0.92, 1.31)
4	613	263	2.3	(1.76, 3.10)
5	100	61	1.6	(1.35, 2.00)
6A [¶]	1056	160	6.6	(5.14, 8.49)
6B	1450	565	2.6	(2.00, 3.29)
7F	559	481	1.2	(0.97, 1.39)
9V	622	491	1.3	(1.08, 1.49)
14	355	366	1.0	(0.76, 1.23)
18C	972	573	1.7	(1.33, 2.16)
19A	366	216	1.7	(1.40, 2.07)
19F	422	295	1.4	(1.16, 1.77)
23F	177	53	3.3	(2.49, 4.47)

GMT, Geometric Mean Titer.

* Study conducted in US and Sweden NCT00546572 (Study 7).

† For the 12 common serotypes, noninferiority was defined as the lower limit of the 2-sided 95% CI for GMT ratio (Prevnar 13/PPSV23) greater than 0.5.

‡ For serotype 6A, which is unique to Prevnar 13, a statistically significantly greater response was defined as the lower limit of the 2-sided 95% CI for the GMT ratio (Prevnar 13/PPSV23) greater than 2.

§ mcOPA antibody for the 11 serotypes unique to PPSV23 but not contained in Prevnar 13 were not measured.

¶ Individual mcOPA antibody assay values below the assay LLOQ (lower limit of quantitation) were set at 0.50*LLOQ for the purpose of calculating the mcOPA antibody GMT.

Evaluable Immunogenicity Population.

[¶] 6A is a serotype unique to Prevnar 13 but not contained in PPSV23.

Clinical Trial of Sequential Vaccination of Prevnar 13 and PPSV23 in PPSV23 Unvaccinated Adults

In a randomized clinical trial conducted in PPSV23-unvaccinated adults 60 through 64 years of age⁸ (Study 8), 223 subjects received PPSV23 followed by Prevnar 13 one year later (PPSV23/Prevnar 13), and 478 received only Prevnar 13. mcOPA antibody titers were measured 1 month after vaccination with Prevnar 13 and are shown in Table 26. mcOPA antibody GMTs in those that received Prevnar 13 one year after PPSV23 were diminished when compared to those who received Prevnar 13 alone. Similarly, in exploratory analyses in PPSV23 previously vaccinated adults ≥ 70 years of age in Study 7, diminished mcOPA antibody GMTs were observed in those that received Prevnar 13 one year after PPSV23 when compared to those who received Prevnar 13 alone.

Table 27: mcOPA Antibody GMTs for the Prevnar 13 Serotypes in PPSV23

Unvaccinated Adults Aged 60 Through 64 Years Given Prevnar 13 Alone or Prevnar 13 One Year After PPSV23 (Study 8) (PPSV23/Prevnar 13)^{*,†,‡,§}

Serotype	Prevnar 13 N=410–457		PPSV23/Prevnar 13 N=180–196	
	GMT	(95% CI)	GMT	(95% CI)
1	219	(191, 252)	88	(72, 109)
3	78	(69, 88)	54	(45, 65)
4	2590	(2257, 2973)	988	(802, 1218)
5	258	(218, 305)	112	(90, 139)
6A [¶]	2947	(2536, 3426)	1210	(962, 1522)
6B	2165	(1845, 2540)	832	(654, 1059)
7F	1518	(1339, 1721)	407	(342, 485)
9V	1279	(1142, 1432)	495	(426, 575)
14	790	(663, 941)	515	(402, 659)
18C	1683	(1437, 1971)	650	(504, 839)
19A	717	(629, 818)	299	(248, 361)
19F	812	(702, 939)	360	(293, 442)
23F	384	(312, 472)	142	(104, 193)

GMT =Geometric Mean Titer.

* Study conducted in US NCT00574548 (Study 8).

† Evaluable Immunogenicity Population.

‡ mcOPA antibody for the 11 serotypes unique to PPSV23 but not contained in Prevnar 13 were not measured.

§ Individual mcOPA antibody assay values below the assay LLOQ (lower limit of quantitation) were set at 0.50*LLOQ for the purpose of calculating the mcOPA antibody GMT.

¶ 6A is a serotype unique to Prevnar 13 but not contained in PPSV23.

Also in Study 8, 266 subjects received Prevnar 13 followed by PPSV23 one year later (Prevnar 13/PPSV23). mcOPA antibody GMTs following PPSV23 administered one year after Prevnar 13 (Prevnar 13/PPSV23) were noninferior to those following a single dose of PPSV23 (N=237) for the 12 common serotypes [the lower limit of the 95% CI for the GMT ratio [Prevnar 13/PPSV23 relative to PPSV23] was >0.5] (see Table 27). In Study 6, which was conducted in PPSV23-unvaccinated adults 60 through 64 years of age, 108 subjects received PPSV23 3.5 to 4 years after Prevnar 13 (Prevnar 13/PPSV23) and 414 received a single dose of PPSV23. Higher serotype-specific mcOPA antibody GMT ratios [(Prevnar 13/PPSV23) / PPSV23] were generally observed compared to the one year dosing interval in Study 8.

Table 28: mcOPA Antibody GMTs for the Prevnar 13 Serotypes in PPSV23-Unvaccinated Adults Aged 60 Through 64 Years Given PPSV23 One Year After Prevnar 13 Relative to PPSV23 Alone (Study 8)^{*,†,‡,§}

Serotype	Prevnar 13/PPSV23 N=216–233		PPSV23 N=214–229		GMT Ratio (Prevnar 13/PPSV23) / PPSV23	
	GMT	95% CI	GMT	95% CI	Ratio	95% CI
1	155	(131, 182)	161	(131, 198)	1.0	(0.74, 1.25)
3	127	(111, 145)	83	(71, 98)	1.5	(1.23, 1.87)
4	1409	(1202, 1651)	1468	(1139, 1893)	1.0	(0.71, 1.29)
5	220	(184, 264)	178	(144, 222)	1.2	(0.93, 1.64)
6A [¶]	1366	(1122, 1663)	400	(306, 524)	3.4	(2.45, 4.77)
6B	1345	(1113, 1625)	875	(689, 1111)	1.5	(1.14, 2.08)

7F	748	(653, 857)	719	(598, 865)	1.0	(0.83, 1.31)
9V	848	(731, 984)	824	(694, 977)	1.0	(0.82, 1.29)
14	711	(580, 872)	869	(677, 1115)	0.8	(0.59, 1.13)
18C	1115	(925, 1344)	912	(707, 1177)	1.2	(0.89, 1.67)
19A	471	(408, 543)	390	(318, 477)	1.2	(0.94, 1.55)
19F	819	(697, 963)	626	(504, 779)	1.3	(1.00, 1.71)
23F	216	(169, 277)	84	(62, 114)	2.6	(1.74, 3.79)

GMT =Geometric Mean Titer.

* Study conducted in US NCT00574548 (Study 8).

† Evaluable Immunogenicity Population.

‡ mcOPA antibody for the 11 serotypes unique to PPSV23 but not contained in Prevnar 13 were not measured.

§ Individual mcOPA antibody assay values below the assay LLOQ (lower limit of quantitation) were set at 0.50*LLOQ for the purpose of calculating the mcOPA antibody GMT.

¶ 6A is a serotype unique to Prevnar 13 but not contained in PPSV23. Anti-6A mcOPA antibody GMTs were descriptive in nature.

14.4 Concomitant Vaccine Administration

Infants and Toddlers

The concomitant administration of routine US infant vaccines [see *Drug Interactions (7.1)*] with Prevnar 13 was evaluated in two studies: Study 2 [see *Clinical Studies (14.2)*], Pneumococcal Immune Responses Following Three Doses², and the US lot consistency study³ (Study 3). In Study 3, subjects were randomly assigned to receive one of 3 lots of Prevnar 13 or Prevnar in a 2:2:2:1 ratio. The total number of infants vaccinated was 663² (Study 2) and 1699³ (Study 3). Immune responses to concomitant vaccine antigens were compared in infants receiving Prevnar and Prevnar 13. Responses to diphtheria toxoid, tetanus toxoid, pertussis, polio types 1, 2, and 3, hepatitis B, PRP-T, PRP-OMP, measles, and varicella antigens in Prevnar 13 recipients were similar to those in Prevnar recipients. Based on limited data, responses to mumps and rubella antigens in Prevnar 13 recipients were similar to those in Prevnar recipients.

Adults ≥50 Years of Age

Concomitant Administration with QIV

Prevnar 13 was administered to PPSV23 previously vaccinated adults ≥50 years of age concomitantly with a US-licensed inactivated influenza vaccine, quadrivalent (IIV4) (Fluzone Quadrivalent) for the 2014/2015 influenza season (Study 13) [see *Adverse Reactions (6.2)* and *Drug Interactions (7.1)*]. One study group received Prevnar 13 and IIV4 concurrently, followed approximately one month later by placebo. A second study group received IIV4 and placebo concurrently, followed approximately one month later by Prevnar 13.

Serotype-specific pneumococcal antibody responses were measured one month after Prevnar 13 vaccination as OPA GMTs. Noninferiority was demonstrated for each pneumococcal serotype if the lower limit of the 2-sided 95% CI for the GMT ratio (Prevnar 13 + IIV4 relative to Prevnar 13 alone) was >0.5. Although OPA antibody responses to Prevnar 13 generally appeared to be slightly lower when Prevnar 13 was administered concomitantly with IIV4 compared to Prevnar 13 administered alone, noninferiority was demonstrated for all Prevnar 13 pneumococcal serotypes evaluated in Study 13.

Strain-specific influenza antibody responses were measured one month after IIV4 as hemagglutinin inhibition assay (HAI) titers. HAI GMTs were evaluated for each IIV4 strain in Study 13. Noninferiority was demonstrated if the lower limit of the 2-sided 95% CI for the HAI GMT ratio (Prevnar 13 + IIV4 relative to IIV4 + Placebo) was >0.5. Noninferiority was demonstrated for each IIV4 vaccine strain evaluated in Study 13.

Concomitant Administration with TIV

Two randomized, double-blind clinical trials evaluated the immunogenicity of Prevnar 13 given with IIV3 (Fall 2007/ Spring 2008 Fluarix, A/H1N1, A/H3N2, and B strains) in PPSV23 unvaccinated adults aged 50 through 59 years¹⁰ (Study 10, conducted in the US) and in adults ≥ 65 years¹¹ (Study 11, conducted in Europe). Based on analysis of the primary pre-specified comparison of serotype specific anti-capsular polysaccharide IgG GMCs, noninferiority was met for all serotypes in adults 50–59 years of age and for 12 of 13 serotypes in adults ≥ 65 years of age.

15 REFERENCES

ClinicalTrials.gov identifiers for studies included below:

1. Study 1 NCT00205803
2. Study 2 NCT00373958
3. Study 3 NCT00444457
4. Study 4 NCT00452452
5. Study 5 NCT00761631
6. Study 6 NCT00427895
7. Study 7 NCT00546572
8. Study 8 NCT00574548
9. Study 9 NCT00500266
10. Study 10 NCT00521586
11. Study 11 NCT00492557
12. Study 12 NCT00744263
13. Study 13 NCT02124161

16 HOW SUPPLIED/STORAGE AND HANDLING

Product: 50090-1944

NDC: 50090-1944-0 .5 mL in a VIAL / 1 in a POUCH

17 PATIENT COUNSELING INFORMATION

Prior to administration of this vaccine, inform the individual, parent, guardian, or other responsible adult of the following:

- The potential benefits and risks of immunization with Prevnar 13 [*see Warnings and Precautions (5) and Adverse Reactions (6)*].
- The importance of completing the immunization series unless contraindicated.
- Any suspected adverse reactions should be reported to their healthcare professional.

Provide the Vaccine Information Statements, which are available free of charge at the Centers for Disease Control and Prevention (CDC) website (www.cdc.gov/vaccines).

This product's label may have been updated. For current full prescribing information, please visit www.pfizer.com.

Manufactured by



Wyeth Pharmaceuticals Inc


A subsidiary of Pfizer Inc, Philadelphia, PA 19101

US Govt. License No. 3

LAB-0469-17.0

CPT Code 90670

PREVNAR 13 (PNEUMOCOCCAL 13-VALENT CONJUGATE VACCINE) INJECTION, SUSPENSION



PREVNAR 13

pneumococcal 13-valent conjugate vaccine injection, suspension

Product Information

Product Type	VACCINE	Item Code (Source)	NDC:50090-1944(NDC:0005-1971)
Route of Administration	INTRAMUSCULAR		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
STREPTOCOCCUS PNEUMONIAE TYPE 1 CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN (UNII: 54EC0SE5PZ) (STREPTOCOCCUS PNEUMONIAE TYPE 1 CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN - UNII:54EC0SE5PZ)	STREPTOCOCCUS PNEUMONIAE TYPE 1 CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN	2.2 ug in 0.5 mL
STREPTOCOCCUS PNEUMONIAE TYPE 3 CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN (UNII: 2VF3V7175U) (STREPTOCOCCUS PNEUMONIAE TYPE 3 CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN - UNII:2VF3V7175U)	STREPTOCOCCUS PNEUMONIAE TYPE 3 CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN	2.2 ug in 0.5 mL
STREPTOCOCCUS PNEUMONIAE TYPE 4 CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN (UNII: TGJ6YZC4W7) (STREPTOCOCCUS PNEUMONIAE TYPE 4 CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN - UNII:TGJ6YZC4W7)	STREPTOCOCCUS PNEUMONIAE TYPE 4 CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN	2.2 ug in 0.5 mL
STREPTOCOCCUS PNEUMONIAE TYPE 5 CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN (UNII: 5SKG37872O) (STREPTOCOCCUS PNEUMONIAE TYPE 5 CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN - UNII:5SKG37872O)	STREPTOCOCCUS PNEUMONIAE TYPE 5 CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN	2.2 ug in 0.5 mL
STREPTOCOCCUS PNEUMONIAE TYPE 6A CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN (UNII: Z9HK08690W) (STREPTOCOCCUS PNEUMONIAE TYPE 6A CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN - UNII:Z9HK08690W)	STREPTOCOCCUS PNEUMONIAE TYPE 6A CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN	2.2 ug in 0.5 mL

STREPTOCOCCUS PNEUMONIAE TYPE 6B CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN (UNII: 4M543JVT7G) (STREPTOCOCCUS PNEUMONIAE TYPE 6B CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN - UNII:4M543JVT7G)	STREPTOCOCCUS PNEUMONIAE TYPE 6B CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN	4.4 ug in 0.5 mL
STREPTOCOCCUS PNEUMONIAE TYPE 7F CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN (UNII: 0K0S2P98ZJ) (STREPTOCOCCUS PNEUMONIAE TYPE 7F CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN - UNII:0K0S2P98ZJ)	STREPTOCOCCUS PNEUMONIAE TYPE 7F CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN	2.2 ug in 0.5 mL
STREPTOCOCCUS PNEUMONIAE TYPE 9V CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN (UNII: 5Q768OY0GJ) (STREPTOCOCCUS PNEUMONIAE TYPE 9V CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN - UNII:5Q768OY0GJ)	STREPTOCOCCUS PNEUMONIAE TYPE 9V CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN	2.2 ug in 0.5 mL
STREPTOCOCCUS PNEUMONIAE TYPE 14 CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN (UNII: SK54I0S386) (STREPTOCOCCUS PNEUMONIAE TYPE 14 CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN - UNII:SK54I0S386)	STREPTOCOCCUS PNEUMONIAE TYPE 14 CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN	2.2 ug in 0.5 mL
STREPTOCOCCUS PNEUMONIAE TYPE 18C CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN (UNII: XK87D9J012) (STREPTOCOCCUS PNEUMONIAE TYPE 18C CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN - UNII:XK87D9J012)	STREPTOCOCCUS PNEUMONIAE TYPE 18C CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN	2.2 ug in 0.5 mL
STREPTOCOCCUS PNEUMONIAE TYPE 23F CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN (UNII: 25N8E57V6T) (STREPTOCOCCUS PNEUMONIAE TYPE 23F CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN - UNII:25N8E57V6T)	STREPTOCOCCUS PNEUMONIAE TYPE 23F CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN	2.2 ug in 0.5 mL
STREPTOCOCCUS PNEUMONIAE TYPE 19A CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN (UNII: B970MQT365) (STREPTOCOCCUS PNEUMONIAE TYPE 19A CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN - UNII:B970MQT365)	STREPTOCOCCUS PNEUMONIAE TYPE 19A CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN	2.2 ug in 0.5 mL
STREPTOCOCCUS PNEUMONIAE TYPE 19F CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN (UNII: 2E1M7F958B) (STREPTOCOCCUS PNEUMONIAE TYPE 19F CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN - UNII:2E1M7F958B)	STREPTOCOCCUS PNEUMONIAE TYPE 19F CAPSULAR POLYSACCHARIDE DIPHTHERIA CRM197 PROTEIN CONJUGATE ANTIGEN	2.2 ug in 0.5 mL

Inactive Ingredients

Ingredient Name	Strength
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	100 ug in 0.5 mL
ALUMINUM PHOSPHATE (UNII: F92V3S521O)	125 ug in 0.5 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50090-1944-0	1 in 1 POUCH		
1		.5 mL in 1 VIAL; Type 3: Prefilled Biologic Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

BLA	BLA125324	03/12/2010	
-----	-----------	------------	--

Labeler - A-S Medication Solutions (830016429)

Establishment

Name	Address	ID/FEI	Business Operations
A-S Medication Solutions		830016429	RELABEL(50090-1944)

Revised: 8/2018

A-S Medication Solutions