SEDAL3X- acetaminophen, aspirin, caffeine tablet OPMX LLC

Sedal 3X

Drug facts

Active ingredients (in each caplets)

Acetaminophen 250 mg Aspirin 250 mg (NSAID)* Caffeine 65 mg *nonsteroidal anti-inflammatory drug

Uses

Uses
temporarily relieves minor aches and pains due to:
headache
a cold
arthritis
muscular aches
sinusitis
toothache
premenstrual and menstrual cramps

Warnings

Allergy alert

Aspirin may cause a severe allergic reaction, which may include: hives shock facial swelling asthma (wheezing)

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take: more than 8 tablets in 24 hours, wich is the maximum daily amount with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older

have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed

Caffeine warning

The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness and, occasionally, rapid heartbeat.

Do not use

if you ever had an allergic reaction to acetaminophen, aspirin, or any other pain reliever/fever reducer

with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have

liver disease stomach bleeding warning applies to you you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis or kidney disease you have asthma you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking

a prescription drug for diabetes, gout, or arthritis

taking any other drug or are under a doctor's care for any serious condition

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding: feel faint have bloody or black stools vomit blood have stomach pain that does not get better an allergic reaction occurs. Seek medical help right away if: any new symptoms appear ringing in the ears or loss of hearing occurs pain gets worse or lasts for more than 10 days fever gets worse or lasts for more than 3 days painful area is red or swollen

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during

the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not use more than directed drink a full glass of water with each dose adults and children 12 years and over: take 2 tablets every 6 hours do not take more than 8 tablets in 24 hours children under 12 years: ask a doctor

Other information

store at 25°C (77°F) excursions permitted between 15-30°C (59-86°F) TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAPS IS BROKEN OR MISSING

Inactive ingredients

corn starch, crospovidone, hypromellose, microcrystalline cellulose, povidone, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide

Manufactured in: GMP Registered Facility in the USA

Made for:

inov FARMACEUTIQUE

Exclusively distributed by:

OPMX

Chula Vista, CA 91910 PHONE: 619-600-5632

Sedal3x

Acetaminophen 250 mg

Aspirin 250 mg Caffeine 65 mg

Pain Reliever

40 Caplets

PRINCIPAL DISPLAY PANEL - 40 CAPLETS

NDC 69729-055-40

Sedal3x

Acetaminophen 250 mg Pain reliever

Aspirin 250 mg Pain reliever Caffeine 65 mg Pain reliever

40 Caplets







Drug Facts

Active ingredient (in each caplet)

Purpose

Uses

Itemporarily relieves minor aches and pains due to:
headache a cold arthritis muscular aches sinusitis toothache premenstrual and

Warnings

Warnings

Rey's symdrome. Children and teernagers who have or are recovering from children prox or flu-like symproms. Should not use this product. When using this product, if changes in behavior with nause and vormfling occur, consult a doctor because these symproms could be an early sign of Rey's syndrome, a rate but serious liness. Allery's elert Aspirin may cause a severe a leirgic reaction which may include: If these I shock I facial swelling I establish Tay hours, which is read to the should be a should be a severe a leirgic reaction which may footloot contraine set detarinophen. Severe liver damage may occur if you take.

Inver warning: This product contains set'y day white using this product.

Severe liver damage may armon I with other drugs containing a celaminophen III of the contraining and the

this product

It is product

It is en more of for a honger time than directed
Caffeine variage; The recommended dose of the
product conditions about as much caffeine as a our of
ordise. Limit the use of ordinive containing medications,
flooks, or beverages white taking this product because the
output caffeine rings causes inervolvenses; initiability,
steeplessness, and, occasionally, rapid heartheat.

Do not use III in great exit and allegin periadron to
avalammorphen aspirin, or any other pain reliever them or other dury ordining ascelamicaphen
(prescription or comprescription). If you are not save whether
at drug ordinates ascelamicaphens, all dector or pharmacist.

Ask a dector before use III III you have five disease.

Ill so un have it followed pressure, hand desease, liver
ordinates for bidney disease.

Ill you have a shirt you are taking a price sity you are

I sking a prescription drug for diabetes, you, or
antimits III sking any other five use or
antimits III sking any other five use III you are
a triving a prescription drug for diabetes, you, or
antimits III sking any other five use or
and the contraction of the contraction o

Stop use and ask a doctor if
you experience any of the following signs of
stomach bleeding: ■ red faint ■ have bloody or black
stools ■ vormit blood ■ have stomach pain that

take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, Drug Facts (continued)

Drug Facts (continued)

have 3 or more alcoholic drinks every day while using

in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdrose, get medical help or contact a Poison
Control Center (1-800-222-222) right away Juck
medical attention is official for addits as well as for
children even if you do not notice any signs or symptoms. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directled to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keen and 1 reach of this flore.



do not use more than directed drink a full glass of water with each dose adults and children 12 years and over take 2 tablets every 6 hours do not take more than 8 tablets in 24 hours children under 12 years; ask a doctor

Other information
store at 25°C (77°F) excursions permitted between

7 ÉVIDENT: DO NOT USE IF IMPRINTED 7 SEAL UNDER CAP IS BROKEN OR MISSING

Inactive ingredients com starch, crospovidone, hypromellose, microcrystalline cellulose, povidone, propylene glycol, sodium lauryl sulfate, sodiu

SEDAL3X

acetaminophen, aspirin, caffeine tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69729-055
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg

Inactive Ingredients				
Ingredient Name	Strength			
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)				
CROSPOVIDONE (UNII: 2S7830E561)				
POVIDONE (UNII: FZ989GH94E)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
SODIUM LAURYL SULFATE (UNII: 368GB5141J)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)				
STARCH, CORN (UNII: O8232NY3SJ)				

Product Characteristics				
Color	white	Score	no score	
Shape	CAPSULE	Size	17mm	
Flavor		Imprint Code	S53	
Contains				

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69729-055- 40	1 in 1 CARTON	09/07/2022	
1		40 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing	Application Number or Monograph	Marketing Start	Marketing End
Category	Citation	Date	Date

OTC Monograph Drug	M013	09/07/2022	

Labeler - OPMX LLC (029918743)

Establishment				
	Name	Address	ID/FEI	Business Operations
	SIUM PHARMACEUTICALS		915664486	analysis(69729-055), label(69729-055), manufacture(69729-055), pack(69729-055)

Revised: 6/2024 OPMX LLC