

PRO-WORMER 2- pyrantel pamoate suspension
Foster and Smith Inc

Pro-Wormer•2®

ACTIVE INGREDIENT

4.54 mg of Pyrantel Base as Pyrantel Pamoate per mL

IMPORTANT

SHAKE WELL BEFORE USE

REFER TO PACKAGE INSERT FOR COMPLETE DIRECTIONS

Barcode 25141 20270

STORAGE

Store at controlled room temperature between 15°- 30°C (59°- 86°F). Keep container tightly closed when not in use.

FOR QUESTIONS OR TO REORDER

1-800-562-7169

DrsFosterSmith.com

Manufactured for:

Foster and Smith, Inc.

2253 Air Park Road

Rhineland, WI 54501

Item #30311 Iss. 05-12

INDICATIONS

For the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies. The presence of these parasites should be confirmed by laboratory fecal exam. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism. PRO-Wormer•2® may also be used to prevent reinfestation of *Toxocara canis* in puppies and adult dogs and in lactating bitches after whelping.

DIRECTIONS FOR USE

Administer 1 full teaspoon (5 mL) for each 10 lb of body weight. To assure proper dosage, weigh animal prior to treatment. It is not necessary to withhold food prior to or after treatment. Dogs usually find this dewormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2-4 weeks after first treatment. If your dog looks or acts sick, consult your veterinarian before treatment. For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8 and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *Toxocara canis* reinfestation.

TAKE TIME



OBSERVE LABEL
DIRECTIONS

PRINCIPAL DISPLAY PANEL - 60 mL Bottle Label

**DOCTORS
FOSTERandSMITH**

**PRO-Wormer•2
(pyrantel pamoate)
CANINE ANTHELMINTIC SUSPENSION**

Removes Large Roundworms
and Hookworms

For Animal Use Only Keep Out of Reach of Children

ANADA 200-007, Approved by FDA

Net Contents: 2 fl oz (60 mL)

ACTIVE INGREDIENT: 4.54 mg of pyrantel base as pyrantel pamoate per mL.
IMPORTANT: SHAKE WELL BEFORE USE.
REFER TO PACKAGE INSERT FOR COMPLETE DIRECTIONS

STORAGE: Store at controlled room temperature between 15° - 30 °C (59° - 86 °F). Keep container tightly closed when not in use.

FOR QUESTIONS OR TO REORDER:
 1-800-562-7169
 DrsFosterSmith.com
 Manufactured for:
 Foster and Smith, Inc.
 2253 Air Park Road
 Rhinelander, WI 54501
 Item #30311 Iss. 05-12

**DOCTORS
FOSTERandSMITH
PRO-Wormer•2[®]
(pyrantel pamoate)
CANINE ANTHELMINTIC SUSPENSION
Removes Large Roundworms
and Hookworms
For Animal Use Only • Keep Out of Reach of Children
ANADA 200-007, Approved by FDA
Net Contents: 2 fl oz (60 mL)**

INDICATIONS: For the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies. The presence of these parasites should be confirmed by laboratory fecal exam. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism. PRO-Wormer•2[®] may also be used to prevent reinestation of *Toxocara canis* in puppies and adult dogs and in lactating bitches after whelping.

DIRECTIONS FOR USE: Administer 1 full teaspoon (5 mL) for each 10 lb of body weight. To assure proper dosage, weigh animal prior to treatment. It is not necessary to withhold food prior to or after treatment. Dogs usually find this dewormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2-4 weeks after first treatment. If your dog looks or acts sick, consult your veterinarian before treatment. For maximum control and prevention of reinestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8 and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *Toxocara canis* reinestation.

8 25141 20270 6

1985 1984 DOCTORS LABEL
FOSTERandSMITH

PRO-WORMER 2

pyrantel pamoate suspension

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:65713-303
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
pyrantel pamoate (UNII: 81BK194Z5M) (pyrantel - UNII:4QIH0N49E7)	pyrantel	4.54 mg in 1 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65713-303-02	60 mL in 1 BOTTLE, PLASTIC		
2	NDC:65713-303-16	473 mL in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200007	08/30/2012	

Labeler - Foster and Smith Inc (122135148)**Registrant** - Happy Jack Inc (003189354)**Establishment**

Name	Address	ID/FEI	Business Operations
ABS Corporation		048022727	MANUFACTURE

Establishment

Name	Address	ID/FEI	Business Operations
COSMA Spa		428655732	API MANUFACTURE

Revised: 7/2013

Foster and Smith Inc