

## **RITE AID DOUBLE ANTIBIOTIC- double antibiotic ointment**

### **RITE AID**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **Drug Facts**

#### **Active ingredients (In each gram)**

Bacitracin 500 units

Polymyxin B Sulfate 10,000 units

#### **Purpose**

First aid antibiotic

#### **Uses**

First aid to help prevent infection in minor:

- cuts
- scrapes
- burns

#### **Warnings**

**For external use only**

#### **Allergy alert:**

- do not use if you are allergic to any of the ingredients

#### **Do not use**

- in or near the eyes
- on large areas of the body

#### **Consult a doctor before use if you have**

deep or puncture wounds  
animal bites  
serious burns

#### **When using this product**

- do not use longer than 1 week

**Keep out of the reach of children.**

If swallowed get medical help or contact a Poison Control Center immediately.

**Directions**

- clean the affected area
- apply a small amount (equal to surface area of tip of finger) on the area 1 to 3 times daily.
- may be covered with a sterile bandage

**Inactive ingredient**

Petrolatum

**Store at a controlled room temperature 20°-25°C (68° to 77°F)**

**Questions?Call; 1-800-719-9260**

**Principal display panel -Tube**

Double Antibiotic Ointment      NDC 11822-0093-5

Bacitracin Zinc - 500 Bacitracin units

Polymixin B Sulfate - 10,000 units

First Aid Antibiotic

NET WT 0.5oz (14g)

NDC 11822-0093-5

# DOUBLE ANTIBIOTIC OINTMENT

BACITRACIN ZINC AND POLYMYXIN  
B SULFATE OINTMENT

## FIRST AID ANTIBIOTIC

Helps prevent infection in  
minor cuts, scrapes & burns



NET WT 0.5 OZ (14 g)

### Active ingredients (in each gram)

Bacitracin 500 units, Polymyxin B 10,000 units.....

### Purpose

First aid antibiotic

**Uses** first aid to help prevent infection in: • minor cuts • scrapes • burns.

**Warnings** • For external use only. **Allergy alert:** • do not use if allergic to any of the ingredients.

**Do not use** • in or near the eyes • on large areas of the body. **Ask a doctor before use if you have**

• deep or puncture wounds • animal bites • serious burns. **When using this product** • do not use

longer than 1 week. **Stop use and ask a doctor if** • condition lasts or gets worse • symptoms last

for more than 7 days or clear up and come back within a few days • a rash or other allergic reaction

occurs. **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control

Center immediately. 1-800-222-1222 **Directions** • clean affected area • apply a small amount (equal

to the surface area of tip of finger) on area 1 to 3 times daily • may be covered with a sterile bandage.

**Other information** • Store at controlled room temperature 68°-77°F (20°-25°C).

**Questions?** Call 1-800-719-9260 \*This product is not manufactured or distributed by ©Johnson & Johnson

Consumer Products Company - A Division of Johnson & Johnson Consumer Companies, Inc., owner of the

registered trademark Polysporin®

DISTRIBUTED BY: RITE AID, 30 HUNTER LANE, CAMP HILL, PA 17011 www.riteaid.com MADE IN THE USA OF U.S. & IMPORTED INGREDIENTS AND COMPONENTS

255974

## Principal Display Panel -Carton

Double Antibiotic Ointment NDC 11822-0093-5

Bacitracin Zinc - 500 Bacitracin units

Polymyxin B Sulfate - 10,000 units

First Aid Antibiotic

NET WT 0.5oz (14g)



**Inactive Ingredients**

Ingredient Name	Strength
PETROLATUM (UNII: 4T6H12BN9U)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0093-5	1 in 1 CARTON	10/01/2022	
1		14 g in 1 TUBE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333B	10/01/2022	

**Labeler** - RITE AID (014578892)

Revised: 10/2022

RITE AID