PENAQUA SOL-G- penicillin g potassium powder, for solution Bimeda Inc.

PenAqua Sol-G® (penicillin G potassium, USP) FOR ORAL USE IN TURKEYS ONLY ANTIBIOTIC for drinking water

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Nonsterile

ACTIVE INGREDIENT:

0.500 billion units penicillin G potassium

NOT FOR USE IN HUMANS

KEEP OUT OF REACH OF CHILDREN

Restricted Drug (California) - Use Only as Directed

Approved by FDA under ANADA # 200-103

Net Contents: 0.500 billion I.U.

INDICATIONS: For treatment of erysipelas in turkeys caused by *Erysipelothrix* rhusiopathiae.

DOSAGE AND ADMINISTRATION: Administer orally at a dosage of 1,500,000 units of penicillin per gallon (3.8 liters) of drinking water for 5 consecutive days.

DIRECTIONS: Dissolve contents of pouch in 2 pints (946 mL) of water in glass or plastic container. Shake well to dissolve. Allow the concentrated solution to stand until the foam disappears. The concentrated solution should be used up or discarded within one hour of preparation.

Automatic Watering Systems: Pour the concentrated solution into a glass or plastic container, then add enough water to make 2.6 gallons (9.9 liters) of stock solution. [This amount of solution will medicate 333 gallons (1260 liters) of drinking water]. The automatic waterer should be adjusted to deliver 1 ounce (30 mL) of stock solution per gallon (3.8 liters) of drinking water. Prepare fresh stock solutions and medicated drinking water solutions every twelve (12) hours. All solutions in contact with galvanized metal should be changed every three (3) hours.

Gravity Flow Watering Systems: Pour the concentrated solution into enough water to make 333 gallons (1260 liters) of drinking water. In gravity flow watering systems, prepare fresh solutions every twelve (12) hours. All solutions in contact with galvanized metal should be changed every three (3) hours.

Drinking water prepared as directed above will contain 1,500,000 units of penicillin G potassium per gallon (3.8 liters).


WARNINGS: Treated turkeys must not be slaughtered for food during treatment and for one day after last treatment.

Do not use in turkeys producing eggs for human consumption.

PRECAUTIONS: For best results, the treatment should be started at the first sign of infection. If improvement is not noted after 3 to 4 days of treatment, consult a poultry pathologist or veterinarian. Use only as directed.

Store at or below 25°C (77°F). Protect from excessive heat, 40°C (104°F), and moisture. Label recommendations for storage and replacement of stock and medicated water solutions must be followed to assure the performance of this drug product.

The Safety Data Sheet (SDS) contains more detailed occupational safety information. To report suspected adverse drug events, for technical assistance or to obtain a copy of the SDS, contact Bimeda, Inc. at 1-888-524-6332. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at www.fda.gov/reportanimalae.





PenAqua Sol-G[®]

(penicillin G potassium, USP) (penicilina G potasio, USP)

FOR ORAL USE IN TURKEYS ONLY/PARA USO ORAL SOLO EN PAVOS

ANTIBIOTIC for drinking water

ANTIBIÓTICO para agua de consumo

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

PRECAUCIÓN: Las leyes federales restringen este fármaco al uso por o bajo la orden de un veterinario con licencia,

Nonsterile / No estéril

ACTIVE INGREDIENT:

1.000 billion units penicillin G potassium

INGREDIENTE ACTIVO:

1.00 billón de unidades de penicilina G potasio

NOT FOR USE IN HUMANS / NO ADMINISTRAR A HUMANOS

KEEP OUT OF REACH OF CHILDREN / MANTENER FUERA DEL ALCANCE DE LOS NIÑOS

Restricted Drug (California) – Use Only as Directed / Fármaco Restringido (California) – Solo Usar Como Se Indica Approved by FDA under ANADA # 200-103 / Aprobado por la FDA bajo ANADA # 200-103



Net

Contents: 1.000 billion I.U.

Contenido Neto: 1.000 billón de I.U.

PENAQUA SOL-G

penicillin g potassium powder, for solution

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:61133-5448
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PENICILLIN G POTASSIUM (UNII: VL775ZTH4C) (PENICILLIN G - UNII:Q42T66VG0C)	PENICILLIN G	500000000 [iU] in 500000000 [iU]

Pac	

#	Item Code	Package Description	Marketing Start Date	Marketing End Date

 1 NDC:61133-5448-3
 500000000 [iU] in 1 POUCH

 2 NDC:61133-5448-1
 1000000000 [iU] in 1 POUCH

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANADA	ANADA200103	09/12/2007		

PENAQUA SOL-G

penicillin g potassium powder, for solution

Product Information				
Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:61133-5449	
Route of Administration	ORAL			

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
PENICILLIN G POTASSIUM (UNII: VL775ZTH4C) (PENICILLIN G - UNII:Q42T66VG0C)	PENICILLIN G	1000000000 in 1000000000 [iU]	

P	ackaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61133-5449-1	1000000000 [iU] in 1 POUCH		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200103	08/02/2013	10/09/2015

Labeler - Bimeda Inc. (060492923)

Registrant - Bimeda Inc. (060492923)

Establishment			
Name	Address	ID/FEI	Business Operations
Constant Irwindale		010569463	manufacture

Revised: 1/2024 Bimeda Inc.