

**8 HR ARTHRITIS PAIN RELIEF- acetaminophen tablet, extended release  
DOLGENCORP, INC.**

-----  
**699R Dollar General 55910-892 Acetaminophen Extended Release Tablets  
650 mg Arthritis Pain**

***Drug Facts***

***Active ingredient (in each caplet)***

Acetaminophen 650 mg

***Purpose***

Pain reliever/fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:
  - o minor pain of arthritis
  - o muscular aches
  - o backache
  - o premenstrual and menstrual cramps
  - o the common cold
  - o headache
  - o toothache
- temporarily reduces fever

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have** liver disease

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug

warfarin

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms

**Directions**

**Do not take more than directed.**

See overdose warning

adults

- take 2 caplets every 8 hours with water
  - swallow whole; do not crush, chew, split, or dissolve
  - do not take more than 6 caplets in 24 hours
  - do not use for more than 10 days unless directed by a doctor
- under 18 years of age
- ask a doctor

**Other information**

- store between 20-25°C (68-77°F)
- The FDA approved Dissolution methods differ from USP

**Inactive ingredients** carnauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

**Questions or comments?** Call **1-877-290-4008**

**DG™** health

NDC 55910-892-26

# 8 hr Arthritis Pain Relief

**Acetaminophen Extended-Release Tablets USP, 650 mg Pain Reliever/Fever Reducer**  
For the Temporary Relief of Minor Arthritis Pain

**DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN**

**225 Caplets\*\* -650 mg Each**  
(\*\*Capsule Shaped Bi-Layer Tablets)

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**  
**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

**Uses**

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - muscular aches
  - backache
  - premenstrual and menstrual cramps
  - the common cold
  - headache
  - toothache
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take more than 6 caplets in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen. 3 or more alcoholic drinks every day while using this product may include: ■ skin reddening ■ blisters ■ rash ■ a skin reaction occurs, stop use and seek medical help right away. **Stop using and ask a doctor if:**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present
- These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

**Do not take more than directed.** See overdose warning adults ■ take 2 caplets every 8 hours with water ■ swallow whole; do not crush, chew, split or dissolve ■ do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor under 18 years of age ■ ask a doctor

**Other information**

■ store between 20-25°C (68-77°F)  
■ The FDA approved Dissolution methods differ from USP **Questions or comments?** Call 1-877-290-4008  
DISTRIBUTED BY: OLD EAST MAIN CO.  
100 MISSION RIDGE, GOODLETTSVILLE, TN 37072  
699R 0123 Made in India

Lot No.:

Exp. Date:

**Varnish Omit Area**

Varnish Omit

Exp Date:  
Lot No.:



TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**

**Active ingredient (in each caplet)** **Purpose**  
Acetaminophen 650 mg.....Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - muscular aches
  - backache
  - premenstrual and menstrual cramps
  - the common cold
  - headache
  - toothache
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take more than 6 caplets in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen. 3 or more alcoholic drinks every day while using this product. **Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Drug Facts (continued)**

**Ask a doctor before use if you have** liver disease  
**Ask a doctor or pharmacist before use if you are taking** the blood thinning drug warfarin  
**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

**Do not take more than directed.** see overdose warning

<b>adults</b>	<ul style="list-style-type: none"> <li>■ take 2 caplets every 8 hours with water</li> <li>■ swallow whole; do not crush, chew, split or dissolve</li> <li>■ do not take more than 6 caplets in 24 hours</li> <li>■ do not use for more than 10 days unless directed by a doctor</li> </ul>
<b>under 18 years of age</b>	<ul style="list-style-type: none"> <li>■ ask a doctor</li> </ul>

**Drug Facts (continued)**

**Other information**

- store between 20-25°C (68-77°F)
- The FDA approved Dissolution methods differ from USP

**Inactive ingredients** carnauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

**Questions or comments?**  
Call 1-877-290-4008

**Contains no aspirin**

DISTRIBUTED BY OLD EAST MAIN CO.  
100 MISSION RIDGE, GOODLETTSVILLE, TN 37072  
699R 0123  
MADE IN INDIA

\*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare division, owner of the registered trademark Tylenol®

NDC 55910-892-26  
Compare to the active ingredient of Tylenol® 8-hr Arthritis Pain\*

**DG™** health

**8 hr Arthritis Pain Relief**  
**Acetaminophen Extended-Release Tablets USP, 650 mg Pain Reliever/Fever Reducer**  
For the Temporary Relief of Minor Arthritis Pain

**225 Caplets\*\***  
**650 mg Each**  
(\*\*Capsule Shaped Bi-Layer Tablets)

Actual Size

**DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN**

**DG™** health  
**8 hr Arthritis Pain Relief**  
Acetaminophen Extended-Release Tablets, 650 mg  
Pain Reliever/Fever Reducer  
For the Temporary Relief of Minor Arthritis Pain

health DG™

NDC 55910-892-05

# 8 hr Arthritis Pain Relief

Acetaminophen Extended-Release Tablets USP, 650 mg  
Pain Reliever/Fever Reducer

For the Temporary Relief of Minor Arthritis Pain

**DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN**

**50 Caplets\*\* - 650 mg Each**

(\*\*Capsule Shaped Bi-Layer Tablets)

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

**Uses** ■ temporarily relieves minor aches and pains due to:  
 ● minor pain of arthritis ● muscular aches ● backache  
 ● premenstrual and menstrual cramps ● the common cold  
 ● headache ● toothache ■ temporarily reduces fever

**Warnings** **Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 6 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. **Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:  
 ■ skin reddening ■ blisters ■ rash ■ a skin reaction occurs, stop use and seek medical help right away. **Stop use and ask a doctor if** ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present These could be signs of a serious condition. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions** ■ Do not take more than directed. See overdose warning. ■ Take 2 caplets every 8 hours with water ■ swallow whole; do not crush, chew, split or dissolve ■ do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor ■ ask a doctor

**Other information** ■ store between 20-25°C (68-77°F)  
 ■ The FDA approved Dissolution methods differ from USP **Questions or comments?** Call 1-877-290-4008

DISTRIBUTED BY: OLD EAST MAIN CO., 100 MISSION RIDGE, GOODLETTSVILLE, TN 37072  
©99R 0123 Made in India

Varnish Omit Area

Lot No.:  
Exp. Date:

Varnish Omit

Exp Date:  
Lot No.:



699R 0123 MADE IN INDIA  
100 MISSION RIDGE, GOODLETTSVILLE, TN 37072  
DISTRIBUTED BY OLD EAST MAIN CO.  
Healthcare Division, a name of the registered trademark Tylenol®  
Johnson & Johnson Consumer Inc., McKel Consumer  
\*This product is not manufactured or distributed by  
THE BOTTLE IS CUT, TORN, BROKEN OR MISSING  
TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

## Drug Facts

Active ingredient (in each caplet)	Purpose
Acetaminophen 650 mg.....	Pain reliever/fever reducer

## Uses

■ temporarily relieves minor aches and pains due to:  
 ■ minor pain of arthritis  
 ■ muscular aches ■ backache  
 ■ premenstrual and menstrual cramps  
 ■ the common cold ■ headache  
 ■ toothache  
 ■ temporarily reduces fever

## Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take  
 ■ more than 6 caplets in 24 hours, which is the maximum daily amount  
 ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:  
 ■ skin reddening ■ blisters ■ rash  
 If a skin reaction occurs, stop use and seek medical help right away.

## Drug Facts (continued)

### Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have liver disease**

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

### Stop use and ask a doctor if

■ pain gets worse or lasts more than 10 days  
 ■ fever gets worse or lasts more than 3 days  
 ■ new symptoms occur  
 ■ redness or swelling is present  
 These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Drug Facts (continued)

**Directions** Do not take more than directed. See overdose warning

### adults

■ take 2 caplets every 8 hours with water  
 ■ swallow whole; do not crush, chew, split or dissolve  
 ■ do not take more than 6 caplets in 24 hours  
 ■ do not use for more than 10 days unless directed by a doctor

**under 18 years of age**

■ ask a doctor

### Other information

■ store between 20-25°C (68-77°F)  
 ■ The FDA approved Dissolution methods differ from USP

**Inactive ingredients** carnauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

### Questions or comments?

Call 1-877-290-4008

**Contains No Aspirin**

health DG™

NDC 55910-892-05  
Compare to the active ingredient of Tylenol® 8-hr Arthritis Pain™

# 8 hr Arthritis Pain Relief

Acetaminophen Extended-Release Tablets USP, 650 mg

Pain Reliever/Fever Reducer

For the Temporary Relief of

Minor Arthritis Pain

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN

**50 Caplets\*\*  
650 mg Each**

(\*\*Capsule Shaped Bi-Layer Tablets)

Actual Size

E827

Varnish/Print Omit Area

health DG™  
**8 hr Arthritis Pain Relief**  
Acetaminophen Extended-Release Tablets USP, 650 mg  
Pain Reliever/Fever Reducer  
For the Temporary Relief of Minor Arthritis Pain



7100C 55910-892-42  
**DG™** health

# 8 hr Arthritis Pain Relief

Acetaminophen Extended-Release Tablets USP, 650 mg  
 Pain Reliever/Fever Reducer  
 For the Temporary Relief of Minor Arthritis Pain

**DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN**

**24 Caplets\*\* - 650 mg Each**

(\*Capsule Shaped Bi-Layer Tablets)

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

**Uses** ■ temporarily relieves minor aches and pains due to: ■ minor pain of arthritis ■ muscular aches ■ backache ■ premenstrual and menstrual cramps ■ the common cold ■ headache ■ toothache ■ temporarily reduces fever

**Warnings Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 6 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. **Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Stop use and ask a doctor if** ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** **Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions Do not take more than directed.** See overdose warning adults: ■ take 2 caplets every 8 hours with water ■ swallow whole; do not crush, chew, split or dissolve ■ do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor **under 18 years of age:** ■ ask a doctor

**Other information** ■ store between 20-25°C (68-77°F) ■ The FDA approved Dissolution methods differ from USP

**Questions or comments?** Call 1-877-290-4008  
 DISTRIBUTED BY OLD EAST MAIN CO.  
 100 MISSION RIDGE, GOODLETTSVILLE, TN 37072  
 699R 0123 Made in India

Lot No.:

Exp. Date:

**Varnish Omit Area**

Exp Date: Varnish Omit  
 Lot No.:  
 5591089242  
 3  
 0  
 A1314  
 699R 0123 MADE IN INDIA  
 DISTRIBUTED BY OLD EAST MAIN CO.  
 100 MISSION RIDGE, GOODLETTSVILLE, TN 37072  
 Headcare division, owner of the registered trademark "Varnish Omit" is not responsible or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer  
 TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

### Drug Facts

**Active ingredient (in each caplet)**  
 Acetaminophen 650 mg.....Pain reliever/fever reducer

**Uses** ■ temporarily relieves minor aches and pains due to: ■ minor pain of arthritis ■ muscular aches ■ backache ■ premenstrual and menstrual cramps ■ the common cold ■ headache ■ toothache ■ temporarily reduces fever

**Warnings Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

■ more than 6 caplets in 24 hours, which is the maximum daily amount  
 ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:  
 ■ skin reddening ■ blisters ■ rash

### Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**  
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have liver disease**

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**Stop use and ask a doctor if**  
 ■ pain gets worse or lasts more than 10 days  
 ■ fever gets worse or lasts more than 3 days  
 ■ new symptoms occur  
 ■ redness or swelling is present  
 These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Drug Facts (continued)

**Directions Do not take more than directed.** See overdose warning

**adults**  
 ■ take 2 caplets every 8 hours with water  
 ■ swallow whole; do not crush, chew, split or dissolve  
 ■ do not take more than 6 caplets in 24 hours  
 ■ do not use for more than 10 days unless directed by a doctor

**under 18 years of age**  
 ■ ask a doctor

### Other information

■ store between 20-25°C (68-77°F)  
 ■ The FDA approved Dissolution methods differ from USP

**Inactive ingredients** carnauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

### Questions or comments?

Call 1-877-290-4008

**Contains No Aspirin**

NDC 55910-892-42  
 Compare to the active ingredient of Tylenol® 8 hr Arthritis Pain\*

**8 hr Arthritis Pain Relief**  
 Acetaminophen Extended-Release Tablets USP, 650 mg  
 Pain Reliever/Fever Reducer  
 For the Temporary Relief of Minor Arthritis Pain

**24 Caplets\*\* 650 mg Each**

(\* Capsule Shaped Bi-Layer Tablets)  
 Actual Size

**DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN**

**Actual Size**

**8 hr Arthritis Pain Relief**  
 Acetaminophen Extended-Release Tablets USP, 650 mg  
 Pain Reliever/Fever Reducer  
 For the Temporary Relief of Minor Arthritis Pain

**DG™** health

# 8 HR ARTHRITIS PAIN RELIEF

acetaminophen tablet, extended release

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55910-892
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>SODIUM STARCH GLYCOLATE TYPE A</b> (UNII: H8AV0SQX4D)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I3O)	
<b>HYDROXYETHYL CELLULOSE (140 MPa.S AT 5%)</b> (UNII: 8136Y38GY5)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	white (White to off white)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	71
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-892-42	1 in 1 CARTON	04/29/2023	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:55910-892-05	1 in 1 CARTON	04/29/2023	
2		50 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:55910-892-26	1 in 1 CARTON	04/29/2023	
3		225 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

ANDA	ANDA215486	04/29/2023	
------	------------	------------	--

**Labeler -** DOLGENCORP, INC. (068331990)

**Registrant -** TIME CAP LABORATORIES, INC. (037052099)

**Establishment**

Name	Address	ID/FEI	Business Operations
Marksans Pharma Ltd		925822975	manufacture(55910-892)

Revised: 4/2023

DOLGENCORP, INC.