LIDONEX- lidocaine hcl liquid Newpharma, Inc.

Active Ingredient

Lidocaine HCL, USP 4%

Purpose

Topical Anesthetic

Use

For temporarily relief of pain

Warnings

For external use only

Flammable: Keep away from fire or flame

Do not use

- on wounds or damaged skin
- in large quantaties
- with a heating pad
- if you are allergic to any ingredients of this product

When using this product

- use only as directed.
- avoid contact with the eyes, mucous membranes or rashes
- do not bandage tightly

Stop use and ask doctor if

- skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

If pregnant or breast-feeding. Ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Adults and children over 12 years of age and over

- clean and dry affected area
- apply to affected area not more than 3 to 4 times daily

Children under 12 years of age: Consult a doctor

Inactive ingredients

glycerin, isopropyl alcohol, menthol, methylparaben, propylparaben, purified water,

xanthan gum

- avoid storing prouct in direct sunlight
- prodect product from excessive moisture
- store with lid closed tightly

Questions or comments? (305) 592 - 9216



DIST, BY: NEWPHARMA, INC. MIAMI, FL 33122

DRUG FACTS

Active Ingredients Lidocaine HCI., USP 4%. Purpose Topical Anesthetic

Uses For temporary relief of pain

Warnings

For external use only

Flammable: Keep away from fire or flame

Do not use

- on wounds or damaged skin in large quantities with a heating pad
- if you are allergic to any ingredients of this product

When using this product

■ use only as directed ■ avoid contact with the eyes, mucous membranes or rashes ■ do not use bandage tightly

Stop use and ask a doctor if

- skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering ■ condition worsens ■ symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

If pregnant or breast-feeding. Ask a health professional before use. Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Adults and children 12 years of age and over

■ clean and dry affected area ■ apply to affected area not more than 3 to 4 times daily

Children under 12 years of age: Consult a doctor

Other information

■ Avoid storing product in direct sunlight ■ Protect product from excessive moisture ■ Store with lid closed tightly

Inactive Ingredients

glycerin, isopropyl alcohol, menthol, methylparaben, propylparaben, purified water, xanthan gum

Questions or comments? (305)592-9216

CODE L-003 Rev: 09/22

LIDONEX

lidocaine hcl liquid

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:73564-800

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name
Basis of Strength
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE UNII:98PI200987)
LIDOCAINE HYDROCHLORIDE
ANHYDROUS
4 g
in 100 mL

Inactive Ingredients					
Ingredient Name	Strength				
GLYCERIN (UNII: PDC6A3C0OX)					
ISOPROPYL ALCOHOL (UNII: ND2M416302)					
MENTHOL (UNII: L7T10EIP3A)					
METHYLPARABEN (UNII: A2I8C7HI9T)					
PROPYLPARABEN (UNII: Z8IX2SC10H)					
WATER (UNII: 059QF0KO0R)					
XANTHAN GUM (UNII: TTV12P4NEE)					

l	P	Packaging Packaging			
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:73564-800- 02	60 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/04/2022	

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC Monograph Drug	M017	10/04/2022			

Labeler - Newpharma, Inc. (019170437)

Revised: 2/2024 Newpharma, Inc.