

**LEADER ASPIRIN- aspirin tablet, film coated**  
**Cardinal Health 110, LLC. dba Leader**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Cardinal Health Aspirin Drug Facts**

**Active ingredient (in each tablet)**

Aspirin 325 mg (NSAID\*)

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- colds
- minor pain of arthritis
- headache
- muscle pain
- toothache
- menstrual pain
- temporarily reduces fever

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Do not use**

if you have ever had an allergic reaction to any other pain reliever/fever reducer

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

### **Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

## Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- children under 12 years: consult a doctor

## Other information

- store at 20-25°C (68-77°F)

## Inactive ingredients

corn starch, dibasic calcium phosphate dihydrate, hypromellose, talc, triacetin

## Questions or comments?

**1-800-719-9260**

## Principal Display Panel

LEADER™

Genuine | Adult

Aspirin

Tablets, 325 mg | Pain Reliever

Fever Reducer (NSAID)

Safe Pain Relief Plus Lifesaving Benefits

Talk to Your Doctor or Other Healthcare Provider Before Using This Product for Your Heart.

COMPARE TO BAYER® ASPIRIN

active ingredient

300 COATED TABLETS

100% Money Back Guarantee

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

**Drug Facts (continued)**

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn

◀ PEEL BACK HERE

LEADER<sup>™</sup>

NDC 70000-0507-2

**Genuine | Adult Aspirin**

**Tablets, 325 mg | Pain Reliever  
Fever Reducer (NSAID)**

Safe Pain Relief Plus  
Lifesaving Benefits

Talk to Your Doctor  
or Other Healthcare  
Provider Before  
Using This Product  
for Your Heart.

**300 COATED TABLETS**

COMPARE TO  
BAYER® ASPIRIN  
active ingredient\*\*

100% Money  
Back Guarantee

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING

**Drug Facts**

**Active ingredient (in each tablet) Purposes**  
Aspirin 325 mg (NSAID)\*.....Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses** ■ temporarily relieves minor aches and pains due to: ■ colds ■ minor pain of arthritis ■ headache ■ toothache ■ muscle pain ■ menstrual pain ■ temporarily reduces fever

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

REV. 11/19  
CIN 5515671



: 27087 E9 F2

**Drug Facts (continued)**

- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic ■ you have asthma

**Ask a doctor or pharmacist before use** if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint ■ vomit blood ■ have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present ■ new symptoms occur
- ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- children under 12 years: consult a doctor

**Other information**

- store at 20-25°C (68-77°F)

**Inactive ingredients**

corn starch, dibasic calcium phosphate dihydrate, hypromellose, talc, triacetin

**Questions or comments?** 1-800-719-9260

\*\*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Bayer® Aspirin. ©2019 Cardinal Health. All Rights Reserved. CARDINAL HEALTH, the Cardinal Health LOGO, LEADER, and the Leader LOGO are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners. DISTRIBUTED BY CARDINAL HEALTH, DUBLIN, OHIO 43017 www.myleader.com 1-800-200-6313

**100% Money Back Guarantee**  
Return to place of purchase if not satisfied.

ADHESIVE AREA  
NO COPY



## LEADER ASPIRIN

aspirin tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:70000-0507
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

### Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
TALC (UNII: 7SEV7J4R1U)	
TRIACETIN (UNII: XHX3C3X673)	

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	14mm
<b>Flavor</b>		<b>Imprint Code</b>	Aspirin;L
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0507-2	300 in 1 BOTTLE; Type 0: Not a Combination Product	01/21/2020	
2	NDC:70000-0507-1	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/20/2020	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	01/21/2020	

**Labeler** - Cardinal Health 110, LLC. dba Leader (063997360)

Revised: 11/2022

Cardinal Health 110, LLC. dba Leader