

**NIGHTTIME COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled
Meijer, Inc.**

Nighttime Cold and Flu

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethorphan hydrobromide 15 mg

Doxylamine succinate 6.25 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat n headache n minor aches and pains
- fever
- runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen.

Severe liver damage may occur if you take:

- more than 4 doses in 24 hours, which is the maximum daily amount for this product.
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, or occurs with or is followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin.

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur n avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults and for children even if you do not notice any signs or symptoms.

Directions

- take only as directed - **see Overdose warning.**
- do not exceed 4 doses per 24 hours

adults & children 12 years & over

2 softgels with water every 6 hours

children 4 to under 12 years

ask a doctor

children under 4 years

do not use

When using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information

- store between 15°C to 30°C (59°F to 86°F)
- avoid excessive heat

Inactive Ingredients

D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, lecithin, myglyol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan, titanium dioxide

Questions or comments?

1-888-333-9792

Principal Display Carton

EXP. DATE:
LOT NO.:

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION
TAMPER EVIDENT- DO NOT USE IF PACKAGING IS OPENED OR IF
BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

NDC 79481-002-7

meijer

COMPARE TO VICKS®
NYQUIL® COLD & FLU
LIQUICAPS®
ACTIVE INGREDIENT*

Nighttime Cold & Flu

PAIN RELIEVER/FEVER REDUCER,
COUGH SUPPRESSANT
ANTIHISTAMINE &
NASAL DECONGESTANT

Acetaminophen | 325 mg
Dextromethorphan HBr | 15 mg
Doxylamine Succinate | 6.25 mg

24
SOFTGELS**
(**Liquid-filled Capsules)

RELIEVES: ACHEs, FEVER,
SORE THROAT, COUGH,
RUNNY NOSE, SNEEZING

Actual Size

meijer
Nighttime Cold & Flu

Drug Facts

Active ingredients (in each softgel)

Acetaminophen	Pain reliever/fever reducer
325 mg	
Dextromethorphan HBr	Cough suppressant
15 mg	
Doxylamine succinate	Antihistamine
6.25 mg	

Uses temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4 doses in 24 hours, which is the maximum daily amount for this product.
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin redness
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, or occurs with or is followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma

Purpose

- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are:

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin.

When using this product

- do not use more than directed
- drowsiness may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults and for children even if you do not notice any signs or symptoms.

Directions

- take only as directed - see **Overdose warning**.
- do not exceed 4 doses per 24 hours

Adults and children 12 years and over: 2 softgels with water every 6 hours

Children 4 to under 12 years: ask a doctor

Children under 4 years: do not use

When using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Drug Facts (continued)

Other information

■ see below to see USP (597 to 598 F) ■ need assessment

Inactive ingredients (D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, lecithin, myglyol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan, titanium dioxide)

Questions or comments? 1-888-333-9792

This product is not manufactured or distributed by the Procter & Gamble Company, owner of the registered trademarks Vicks®, NyQuil®, and LiquiCaps®

how2recycle.info
PAPER
MULTILAYER
BOX
TRAY

Made in India

DIST. BY MEIJER DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544
www.meijer.com
ORIG 0623

OUR QUALITY GUARANTEE
The Meijer Family

PID 467803

7 60236 30881 2

EXP. DATE:
LOT NO.:

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION
TAMPER EVIDENT- DO NOT USE IF PACKAGING IS OPENED OR IF
BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

NDC 79481-002-7

meijer

COMPARE TO VICKS®
NYQUIL® COLD & FLU
LIQUICAPS®
ACTIVE INGREDIENT*

Nighttime Cold & Flu

PAIN RELIEVER/FEVER REDUCER,
COUGH SUPPRESSANT
ANTIHISTAMINE &
NASAL DECONGESTANT

Acetaminophen | 325 mg
Dextromethorphan HBr | 15 mg
Doxylamine Succinate | 6.25 mg

24
SOFTGELS**
(**Liquid-filled Capsules)

RELIEVES: ACHEs, FEVER,
SORE THROAT, COUGH,
RUNNY NOSE, SNEEZING

Actual Size

meijer
Nighttime Cold & Flu

NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79481-0027
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SORBITOL (UNII: 506T60A25R)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	

Product Characteristics

Color	green	Score	no score
Shape	OVAL	Size	21mm
Flavor		Imprint Code	902;215;AP02
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79481-0027-2	2 in 1 CARTON	09/15/2022	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/15/2022	

Labeler - Meijer, Inc. (006959555)

Revised: 12/2023

Meijer, Inc.