

# **ASPIRIN 81MG ENTERIC COATED DELAYED RELEASE- aspirin tablet, coated** **Dr. Wayne's**

-----

## **Aspirin 81mg**

### **Active ingredient(s)**

Aspirin 81mg (NSAID)\*

### **Purpose**

Pain reliever

### **Use(s)**

- for the temporary relief of minor aches and pain or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.
- ask your doctor about other uses for this product

### **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms

should not use this product. When using this product, if changes in behavior with nausea and

vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome,

a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or non prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you are allergic to aspirin or other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if**

you are taking a prescription drug for

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - redness or swelling is present
  - new symptoms occur
  - ringing in the ears or loss of hearing occurs
- You may report serious side effects to 1-888-952-0050

**Pregnancy/Breastfeeding**

If pregnant or breast-feeding, ask a health professional before use.

It is especially important not to use aspirin during the last 3 months of pregnancy unless specifically directed to do

so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away

**Directions**

- drink a full glass of water with each dose
- **adults and children 12 years and over:** take 4 to 8 tablets every 4 hours not to

exceed 48 tablets in 24 hours unless directed by a doctor

- **children under 12 years of age:** consult a doctor

### **Other information**

- store at room temperature
- do not use if imprinted safety seal under cap is broken or missing

### **Inactive ingredients**

anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow

#10 aluminum lake, iron oxide ochre, methacrylic acid copolymer, microcrystalline cellulose, polysorbate 80,

simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

### **Questions**

1-229-417-5113, Monday Through Friday 9am-5pm

### **Principal Display Panel**

[illegible]

mean eye eye, any? ■ You experience any or more troubling signs of stomach bleeding ■ feel hot ■ mouth hoarse ■ three bloody or black stools ■ three stomach pains that don't go better ■ any pain worse or lasts more than 10 days ■ redness or swelling is present ■ new symptoms occur ■ ringing in the ears or loss of hearing occurs ■ two major weight losses since effects to 1,000-502-4930

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center (1-800-222-1222) right away.

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 6 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years of age: consult a doctor

**Other information** ■ store at room temperature ■ do not use if imprint(s) don't seal under cap is broken or missing

**Inactive ingredients** anhydrous benzocaine, camphor wax, colloidal silicon oxide, croscarmellose sodium, DVE, gelatin #10, aluminum hydroxide, iron oxide yellow, methacrylic acid copolymer, microcrystalline cellulose, polyethylene glycol, polyethylene glycol succinate, sodium lauryl sulfate, sodium tartrate, titanium dioxide, white talc

**Questions or comments?**  
1-224-417-5113, Monday Through Friday, 9am-5pm

# ASPIRIN 81MG ENTERIC COATED DELAYED RELEASE

aspirin tablet, coated

## Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72714-405
Route of Administration	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

## Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
BROWN IRON OXIDE (UNII: 1N032N7MFO)	
METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
SODIUM HYDRIDE (UNII: 23J3BHR95O)	
SODIUM LAURETH SULFATE (UNII: BPV390UAP0)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

## Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	HEART
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72714-405-17	300 in 1 BOTTLE; Type 0: Not a Combination Product	12/15/2021	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	12/15/2021	

**Labeler -** Dr. Wayne's (831596312)

Revised: 11/2021

Dr. Wayne's