

**ASPIRIN 81 MG LOW DOSE- aspirin tablet, delayed release**  
**Meijer Distribution Inc**

-----  
**Meijer 44-600A-Delisted**

***Active ingredient (in each tablet)***

Aspirin 81 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever

***Uses***

for the temporary relief of minor aches and pains or as recommended by your doctor.  
**Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - vomit blood
  - have bloody or black stools
  - feel faint
  - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

***Other information***

- use by expiration date on package
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

## ***Inactive ingredients***

corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

## ***Questions or comments?***

**1-800-426-9391**

## **Principal Display Panel**

NDC 41250-600-14

**meijer®**

**Compare to Bayer®  
Low Dose Aspirin  
active ingredient†**

**aspirin 81 mg**  
Pain Reliever (**NSAID**)

**Low  
Dose**

Safety Coated | Aspirin Regimen

500 Enteric Coated Tablets

Actual  
Size


Talk to your doctor or other healthcare provider  
before using this product for your heart.

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY  
SEAL UNDER CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed by Bayer AG, owner of the registered  
trademark Bayer® Low Dose Aspirin. 50844 REV0122C60014

**DIST. BY MEIJER  
DISTRIBUTION, INC.  
GRAND RAPIDS, MI 49544  
www.meijer.com**

NDC 41250-600-14




Compare to Bayer® Low Dose Aspirin active ingredient\*

# aspirin 81 mg

Pain Reliever (NSAID)

**LOW DOSE** Safety Coated | Aspirin Regimen

500 Enteric Coated Tablets  Actual Size

Talk to your doctor or other healthcare provider before using this product for your heart.

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**Drug Facts**


**Active ingredient (in each tablet)**  
Aspirin 81 mg (NSAID)\* ..... Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Uses** For the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:  
■ hives ■ facial swelling ■ shock ■ asthma (wheezing)  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you  
■ are age 60 or older ■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticoagulant) or steroid drug  
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
■ have 3 or more alcoholic drinks every day while using this product  
■ take more or for a longer time than directed

**Do not use**  
■ if you are allergic to aspirin or any other pain reliever/fever reducer  
■ if you have ever had an allergic reaction to this product or any of its ingredients

\*This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Low Dose Aspirin.  
50844 REV1012200014  
PID 466056



7 08820 87719 1

DIST. BY MEIJER  
GRAND RAPIDS, MI 49544  
www.meijer.com

**PEEL HERE FOR MORE DRUG FACTS**

**Adhesive Area**

**STOP PEELING**

**Drug Facts (continued)**

**Ask a doctor before use if**  
■ stomach bleeding warning applies to you  
■ you have a history of stomach problems, such as heartburn  
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease  
■ you have asthma  
■ you are taking a diuretic drug for  
■ gout  
■ diabetes  
■ arthritis

**Ask a doctor or pharmacist before use if you are taking a prescription drug for**  
■ dizziness  
■ drowsiness  
■ drowsiness or lightheadedness  
■ drowsiness or lightheadedness  
■ drowsiness or lightheadedness

**Stop use and ask a doctor if**  
■ an allergic reaction occurs. Seek medical help right away.  
■ you experience any of the following signs of stomach bleeding:  
■ vomit blood  
■ have bloody or black stools  
■ feel faint  
■ have stomach pain that does not get better  
■ ringing in the ears or a loss of hearing occurs  
■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present  
■ new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
■ do not take more than directed  
■ drink a full glass of water with each dose  
■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor  
■ children under 12 years: do not use unless directed by a doctor

**Other information**  
■ use by expiration date on package  
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

**Inactive ingredients** corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

**(Questions or comments?) 1-800-426-9391**

Meijer 44-600A

## ASPIRIN 81 MG LOW DOSE

aspirin tablet, delayed release

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-600
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

### Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	

<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)
<b>METHACRYLIC ACID</b> (UNII: 1CS02G8656)
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ05DW1A)
<b>SHELLAC</b> (UNII: 46N107B71O)
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)
<b>WATER</b> (UNII: 059QF0KO0R)
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)
<b>TALC</b> (UNII: 7SEV7J4R1U)
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)
<b>TRIACETIN</b> (UNII: XHX3C3X673)
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)

### Product Characteristics

<b>Color</b>	yellow	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	6mm
<b>Flavor</b>		<b>Imprint Code</b>	L
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-600-98	2 in 1 CARTON	05/01/2011	11/02/2026
1		365 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:41250-600-14	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/01/2011	11/02/2026
3	NDC:41250-600-32	1 in 1 CARTON	05/01/2011	03/29/2019
3		120 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/01/2011	11/02/2026

**Labeler** - Meijer Distribution Inc (006959555)

### Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

LNK International, Inc.		832867837	manufacture(41250-600)
-------------------------	--	-----------	------------------------

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(41250-600)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(41250-600) , pack(41250-600)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(41250-600)

Revised: 1/2025

Meijer Distribution Inc