

PAIN RELIEF- naproxen sodium tablet
BETTER LIVING BRANDS, LLC

608R_Albertsons_21130-608_Naproxen sodium tablets 220 mg

Drug Facts

Active ingredient (in each caplet)

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains due to:

- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold

temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 1. feel faint
 2. vomit blood
 3. have bloody or black stools
 4. have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 1. chest pain
 2. trouble breathing
 3. weakness in one part or side of body
 4. slurred speech
 5. leg swelling
- pain gets worse or lasts more than 10 days

- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

Adults and children 12 years and older

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12-hour period
- do not exceed 3 caplets in a 24-hour period

Children under 12 years

- ask a doctor

Other information

- **each caplet contains:** sodium 20 mg
- store at 20-25°C (68-77°F).
- avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, FD&C blue #2 Lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, titanium dioxide

Questions or comments? 1-877-290-4008

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except a salicylate, increase the risk of heart attack, or stroke, and stroke, can occur while you are taking the drug.

Do not use: ■ if you have ever had an allergic reaction to any other nonsteroidal anti-inflammatory drug (NSAID).

Ask a doctor before you take: ■ if you are taking warfarin (a blood thinner) ■ if you have a history of stomach problems such as ulcers, heartburn, or bleeding from your rectum or intestines; ■ if you have high blood pressure; ■ if you have diabetes, heart disease, or a stroke; ■ if you are pregnant or nursing; ■ if you have side effects from taking pain in the liver or liver problems.

Ask a doctor or pharmacist before you use: ■ if you are under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke; ■ taking naproxen may decrease this benefit of aspirin ■ taking another drug.

When using this medicine:
■ take with food or milk if stomach upset occurs.

Stop using and call a doctor: ■ if you experience any of the following signs of stomach bleeding:
■ black, tarry, or vomit blood ■ new bloody or black stools
■ have stomach pain that does not get better ■ chest pain
■ trouble breathing ■ weakness in one part or side of body
■ slurred speech ■ eye swelling
■ pain lasts longer or worse than 10 days ■ fever gets worse or lasts more than 3 days ■ you have difficulty swallowing ■ it feels like the pill is stuck in your throat ■ swelling of the mouth or throat is present in the painful area ■ any new symptoms appear.

If pregnant or breast-feeding: ask a health professional before use. It is especially important not to take naproxen during the last weeks of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep up to date with vaccination: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-2222).

Directions: ■ do not take more than directed
■ the smallest effective dose should be used

Directions: ■ drink at least 8 fluid ounces of water with each dose.

Adults and children 12 years and older: ■ take 1 capsule every 8 to 12 hours while symptoms last ■ if you take the first dose, you may take 2 capsules in the first hour ■ do not exceed 2 capsules in any 1- to 12-hour period ■ do not exceed 3 capsules in a 24-hour period
■ ask a doctor

Other Information: ■ each capsule contains: sodium 20 mg, potassium 10 mg, magnesium 10 mg, calcium 10 mg, zinc 10 mg, and copper 10 mg.

inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, D&C Blue #2 Lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, port wine, titanium dioxide

Questions or comments? 1-877-296-408



Adhesive Area



NDC 21130-608-05
Signature care
Quality
Guaranteed
Pain Relief
NAPROXEN SODIUM
TABLETS USP, 220 mg
Pain Reliever/Fever Reducer
(NSAID)
Strength to last 12 hours 50 CAPLETS*
*Capsule-shaped tablets
RE-ENTER CARTON FOR COMPLETE
WARNINGS AND PRODUCT INFORMATION
WARNING
IF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING
THE WRITING, DO NOT USE THIS PRODUCT.
TAMPER EVIDENCE: DO NOT OPEN THIS PRODUCT.
DO NOT USE IF THE WRITING IS FADING, BLURRED
OR UNREADABLE.

Dosage

- take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 - have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed
- Heart attack and stroke warning:** If pregnant or breast-feeding, ask a health professional before use.
- Keep out of reach of children.**
- Directions**
- do not take more than directed
 - the smallest effective dose should be used
 - drink a full glass of water with each dose
 - Adults and children 12 years and older: take 1 caplet every 8 to 12 hours while symptoms last
 - for the first dose you may take 2 caplets within the first hour
 - do not exceed 2 caplets in any 8- to 12-hour period
 - do not exceed 3 caplets in a 24-hour period

Children under 12 years: ask a doctor
Questions or comments?
1-877-290-4008

DISTRIBUTED BY: BETTER LIVING BRANDS LLC
P.O. BOX 99, PLEASANTON, CA 94566-0009
41-889-725-3929
Made in India
608R 0228 55934

- take offter drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 - have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed
- Heart attack and stroke warning:** These can be fatal. The risk is higher if you use more than directed or for longer than directed.
- If pregnant or breast-feeding:** ask a health professional before use.
- Keep out of reach of children.**
- Directions**
- do not take more than directed
 - the smallest effective dose should be used
 - drink a full glass of water with each dose
 - Adults and children 12 years and older: take 1 caplet every 8 to 12 hours while symptoms last
 - for the first dose you may take 2 caplets within the first hour
 - do not exceed 2 caplets in any 8- to 12-hour period
 - do not exceed 3 caplets in a 24-hour period

■ take a blood thinning (anticoagulant) or steroid drug ■ take offter drugs

containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

■ have 3 or more alcoholic drinks every

day while using this product ■ take

more or for a longer time than directed

Heart attack and stroke warning:

If pregnant or breast-feeding, ask a

health professional before use.

Keep out of reach of children.

Directions

■ do not take more than directed

■ the smallest effective dose should

be used

■ drink a full glass of water with each dose

■ Adults and children 12 years and older:

take 1 caplet every 8 to 12 hours

while symptoms last

■ for the first dose you may take 2

caplets within the first hour

■ do not exceed 2 caplets in any 8- to

12-hour period

■ do not exceed 3 caplets in a 24-hour

period

■ Children under 12 years: ask a doctor

Questions or comments?
1-877-290-4008

DISTRIBUTED BY: BETTER LIVING BRANDS LLC
P.O. BOX 99, PLEASANTON, CA 94566-0009
41-889-725-3929
Made in India
608R 0228 55934

Warning Only Area

Do not eat (containing aspirin label)

Take

With

Water

Or

With

Food

Or

With

Water

Or

With

Food



Adhesive Area



NDC 21130-608-42

Pain Relief

NAPROXEN SODIUM
TABLETS USP, 220 mg
Pain Reliever/Fever Reducer
(NSAID)

Strength to last 12 hours 24 CAPLETS**
**Capsule-shaped tablets

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10


10

10

10

10

10



PAIN RELIEF

naproxen sodium tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:21130-608
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

POVIDONE (UNII: FZ989GH94E)

SILICON DIOXIDE (UNII: ETJ7Z6XBU4)

FD&C BLUE NO. 2 (UNII: L06K8R7DQK)

HYPROMELLOSES (UNII: 3NXW29V3WO)

MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

Product Characteristics

Color	blue	Score	no score
Shape	OVAL	Size	12mm
Flavor		Imprint Code	144
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-608-42	1 in 1 CARTON	06/13/2023	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:21130-608-05	1 in 1 CARTON	06/13/2023	
2		50 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:21130-608-20	200 in 1 BOTTLE; Type 0: Not a Combination Product	06/13/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090545	06/13/2023	

Labeler - BETTER LIVING BRANDS, LLC (009137209)

Registrant - TIME CAP LABORATORIES, INC (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	manufacture(21130-608)

Revised: 6/2023

BETTER LIVING BRANDS, LLC