

**ACETAMINOPHEN- acetaminophen tablet, coated**  
**RETAIL BUSINESS SERVICES , LLC.**

-----  
**1004-CAR-2023-1102**

***Drug Facts***

**Active ingredient (in each caplet)**

Acetaminophen 500 mg

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - the common cold
  - headache
  - backache
  - minor pain of arthritis
  - toothache
  - muscular aches
  - premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert**

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or

pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have** liver disease.

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"><li>▪ take 2 caplets every 6 hours while symptoms last</li><li>▪ do not take more than 6 caplets in 24 hours, unless directed by a doctor</li><li>▪ do not use for more than 10 days unless directed by a doctor</li></ul>
children under 12 years	<ul style="list-style-type: none"><li>▪ ask a doctor</li></ul>

**Other information**

- store between 20-25°C (68-77°F)
- retain carton for complete product information

**Inactive ingredients**

hypromellose, mineral oil, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

# PRINCIPAL DISPLAY PANEL

CAREone®

NDC 72476-104-01

Compare to the active ingredient in Tylenol® Extra Strength Caplets†

Extra Strength

ACETAMINOPHEN

Pain Reliever / Fever Reducer

For Adults

Actual Size

24 CAPLETS

500 mg EACH



NC

CAREone®

NDC 72476-104-01

Compare to the active ingredient in Tylenol® Extra Strength Caplets†

Extra Strength

ACETAMINOPHEN

Pain Reliever / Fever Reducer

For Adults



Actual Size

24 CAPLETS  
500 mg EACH

NC

For Adults

Pain Reliever / Fever Reducer

ACETAMINOPHEN

Extra Strength

CAREone®

### Drug Facts

**Active ingredient (in each caplet)**

Acetaminophen 500 mg...Pain reliever/fever reducer

### Uses

- temporarily relieves minor aches and pains due to:
  - headache
  - backache
  - toothache
  - premenstrual and menstrual cramps
  - temporarily reduces fever
- the common cold
- minor pain of arthritis
- muscular aches

### Warnings

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take
- more than 4,000 mg of acetaminophen in 24 hours
  - with other drugs containing acetaminophen
  - 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
  - blisters
  - rash
- If a skin reaction occurs, stop use and seek medical help right away.

### Drug Facts (continued)

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have liver disease.**

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

#### Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Drug Facts (continued)

**Directions** ■ do not take more than directed (see overdose warning)

- |                                       |  |
|---------------------------------------|--|
| adults and children 12 years and over | ■ take 2 caplets every 6 hours while symptoms last                         |
|                                       | ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor |
|                                       | ■ do not use for more than 10 days unless directed by a doctor             |

- |                         |                |
|-------------------------|----------------|
| children under 12 years | ■ ask a doctor |
|-------------------------|----------------|

### Other information

- store between 20-25°C (68-77°F)
- retain carton for complete product information

**Inactive ingredients** hypromellose, mineral oil, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

DISTRIBUTED BY:  
ADUSA DISTRIBUTION, LLC  
SALISBURY, NC 28147

For product questions or concerns,  
contact us at 1-833-992-3872

Quality guaranteed or your money back.

†This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® Extra Strength Caplets.



DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING



F100401CAR\_R0

T-0007 0223



3 41520 31942 6

INK AND COATING FREE FOR LOT AND EXPIRATION STAMPING

# ACETAMINOPHEN

acetaminophen tablet, coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:72476-104
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

## Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MINERAL OIL (UNII: T5L8T28FGP)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	M2A4;57344
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72476-104-01	1 in 1 CARTON	01/20/2023	
1		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/20/2023	

Revised: 11/2023

RETAIL BUSINESS SERVICES , LLC.