

G-CON-X- dexbrompheniramine maleate and pseudoephedrine hydrochloride liquid **McLaren Medical**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

G-Con-X

Drug Facts

<i>Active Ingredients (in each 5 mL, teaspoonful)</i>	<i>Purpose</i>
Pseudoephedrine HCl 30 mg	Nasal Decongestant
Dexbrompheniramine Maleate 1 mg	Antihistamine

Indications

- For the temporary relief of nasal congestion due to the common cold, hay fever or other upper respiratory allergies (allergic rhinitis).
- Helps decongest sinus openings and passages; temporarily relieves sinus congestion and pressure.
- For the temporary relief of runny nose, sneezing, itching of the nose or throat and itchy, watery eyes due to hay fever or other respiratory allergies.

Warnings

- **Do not exceed recommended dosage.**
If nervousness, dizziness or sleeplessness occurs, discontinue use and consult a doctor.
- If symptoms do not improve within 7 days or are accompanied by fever, consult a doctor.
- Do not take this product if you have heart disease, high blood pressure, thyroid disease, diabetes or difficulty in urination due to enlargement of the prostate gland unless directed by a doctor.
- Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- May cause excitability especially in children.
- Do not take this product, unless directed by a doctor, if you have a breathing problem such as emphysema or chronic bronchitis or if you have glaucoma or difficulty in urination due to the enlargement of the prostate gland.
- May cause drowsiness; alcohol, sedatives and tranquilizers may increase the drowsiness effect.

Avoid alcoholic beverages while taking this product.

Do not take this product if you are taking sedatives or tranquilizers, without first consulting your doctor. Use caution when driving a motor vehicle or operating machinery.

- **If pregnant or breast-feeding, ask a health professional before use.**
- **In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.**
- **Keep this and all drugs out of the reach of children.**

Directions

Do not take more than 6 doses in any 24-hour period.

Adults and children 12 years of age and over	2 teaspoonfuls (10 mL) every 4 to 6 hours
Children 6 to under 12 years of age	1 teaspoonful (5 mL) every 4 to 6 hours
Children under 6 years of age	Consult a doctor

Inactive Ingredients

Citric Acid, Grape Flavor, Propylene Glycol, Purified Water, Saccharine Sodium, Sodium Benzoate, Sorbitol, Sucralose.

Other Information

Store at 20°-25°C (68°-77°F)

Tamper evident by seal under cap. Do not use if the seal is broken or missing.

PRINCIPAL DISPLAY PANEL - 473 mL Bottle Label

NDC-43913-405-16

G-Con-X**Antihistamine & Nasal Decongestant**

Sugar Free • Dye Free • Alcohol Free • Phenylalanine Free

Grape Flavor

16 FL OZ (473 mL)

Multiple Dose Unit Package

For Dispensing Under Pharmaceutical Supervision Only

McLaren Medical

NDC-43913-405-16

G-Con-X

Antihistamine & Nasal Decongestant

Sugar Free • Dye Free • Alcohol Free • Phenylalanine Free

Grape Flavor

16 FL OZ (473 mL)

Multiple Dose Unit Package

For Dispensing Under Pharmaceutical Supervision Only



Drug Facts (continued)

Do not take this product if you are taking sedatives or tranquilizers, without first consulting your doctor. Use caution when driving a motor vehicle or operating machinery.

- If pregnant or breast-feeding, ask a health professional before use.
- In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.
- Keep this and all drugs out of the reach of children.

Directions

Do not take more than 6 doses in any 24-hour period.

Adults and children 12 years of age and over	2 teaspoonfuls (10 mL) every 4 to 6 hours
Children 6 to under 12 years of age	1 teaspoonful (5 mL) every 4 to 6 hours
Children under 6 years of age	Consult a doctor

Inactive Ingredients

Citric Acid, Grape Flavor, Propylene Glycol, Purified Water, Saccharine Sodium, Sodium Benzoate, Sorbitol, Sucralose.

Other Information

Store at 20°-25°C (68°-77°F)
Tamper evident by seal under cap. Do not use if the seal is broken or missing.

Manufactured in the USA for: McLaren Medical Inc
4070 Laguna St, Coral Gables, FL 33146

LOT #:

EXP DATE:



Drug Facts

Active Ingredients	Purpose
(In each 5 mL, teaspoonful) Pseudoephedrine HCl 30 mg	Nasal Decongestant
Dexbrompheniramine Maleate 1 mg	Antihistamine

Indications

- For the temporary relief of nasal congestion due to the common cold, hay fever or other upper respiratory allergies (allergic rhinitis).
- Helps decongest sinus openings and passages; temporarily relieves sinus congestion and pressure.
- For the temporary relief of runny nose, sneezing, itching of the nose or throat and itchy, watery eyes due to hay fever or other respiratory allergies.

Warnings

- Do not exceed recommended dosage.
- If nervousness, dizziness or sleeplessness occurs, discontinue use and consult a doctor.
- If symptoms do not improve within 7 days or are accompanied by fever, consult a doctor.
- Do not take this product if you have heart disease, high blood pressure, thyroid disease, diabetes or difficulty in urination due to enlargement of the prostate gland unless directed by a doctor.
- Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- May cause excitability especially in children.
- Do not take this product, unless directed by a doctor, if you have a breathing problem such as emphysema or chronic bronchitis or if you have glaucoma or difficulty in urination due to the enlargement of the prostate gland.
- May cause drowsiness; alcohol, sedatives and tranquilizers may increase the drowsiness effect. Avoid alcoholic beverages while taking this product.

G-CON-X

dexbrompheniramine maleate and pseudoephedrine hydrochloride liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:43913-405
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Dexbrompheniramine Maleate (UNII: BPA9UT29BS) (Dexbrompheniramine - UNII:75T64B71RP)	Dexbrompheniramine Maleate	1 mg in 5 mL
Pseudoephedrine Hydrochloride (UNII: 6V9V2RYJ8N) (Pseudoephedrine - UNII:7CUC9DDI9F)	Pseudoephedrine Hydrochloride	30 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
Sodium Benzoate (UNII: OJ245FE5EU)	
Citric Acid Monohydrate (UNII: 2968PHW8QP)	
Sorbitol (UNII: 506T60A25R)	
Saccharin Sodium (UNII: SB8ZUX40TY)	
Sucralose (UNII: 96K6UQ3ZD4)	
Propylene Glycol (UNII: 6DC9Q167V3)	
Water (UNII: 059QF0KO0R)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:439 13-405-16	473 mL in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	02/11/2014	

Labeler - McLaren Medical (013770591)

Registrant - davAgen Pharmaceutical, LLC (967545935)

Establishment

Name	Address	ID/FEI	Business Operations
davAgen Pharmaceutical, LLC		967545935	MANUFACTURE(439 13-405) , PACK(439 13-405) , LABEL(439 13-405) , ANALYSIS(439 13-405)

Revised: 4/2014

McLaren Medical