

SODIUM CHLORIDE - sodium chloride injection, solution
Becton Dickinson and Company

0.9% Sodium Chloride Injection, USP

Rx only

DESCRIPTION

0.9% Sodium Chloride Injection, USP solution is sterile and nonpyrogenic. It is a parenteral solution containing sodium chloride in water for injection intended for intravenous administration.

For 0.9% Sodium Chloride Injection, USP, each 100 mL contains 900 mg sodium chloride in water for injection. Electrolytes per 1,000 mL: sodium 154 mEq; chloride 154 mEq. The osmolarity is 308 mOsmol/L (calc.).

The pH in the 100 mL and smaller containers is 6.0; for the 250 mL and larger containers, the pH is 5.6.

The pH range is 4.5 to 7.0 for all containers.

The solution contains no bacteriostat, antimicrobial agent or added buffer and is intended only as a single-dose injection. When smaller doses are required the unused portion should be discarded.

The solution is a parenteral fluid and electrolyte replenisher.

Sodium Chloride, USP is chemically designated NaCl, a white crystalline powder freely soluble in water.

Water for injection, USP is chemically designated H₂O.

The flexible container is fabricated from a specially formulated non-plasticized, film containing polypropylene and thermoplastic elastomers (**freeflex**[®] bag). The amount of water that can permeate from the container into the overwrap is insufficient to affect the solution significantly. Solutions in contact with the flexible container can leach out certain of the container's chemical components in very small amounts within the expiration period. The suitability of the container material has been confirmed by tests in animals according to USP biological tests for plastic containers.

CLINICAL PHARMACOLOGY

When administered intravenously, the solution provides a source of water and electrolytes.

Solutions which provide combinations of hypotonic or isotonic concentrations of sodium chloride are suitable for parenteral maintenance or replacement of water and electrolyte requirements.

Isotonic concentrations of sodium chloride are suitable for parenteral replacement of

chloride losses that exceed or equal the sodium loss. Hypotonic concentrations of sodium chloride are suited for parenteral maintenance of water requirements when only small quantities of salt are desired. A hypertonic concentration of sodium chloride may be used to repair severe salt depletion syndrome.

Sodium chloride in water dissociates to provide sodium (Na^+) and chloride (Cl^-) ions. Sodium (Na^+) is the principal cation of the extracellular fluid and plays a large part in the therapy of fluid and electrolyte disturbances. Chloride (Cl^-) has an integral role in buffering action when oxygen and carbon dioxide exchange occurs in the red blood cells. The distribution and excretion of sodium (Na^+) and chloride (Cl^-) are largely under the control of the kidney which maintains a balance between intake and output.

Water is an essential constituent of all body tissues and accounts for approximately 70% of total body weight. Average normal adult daily requirements range from two to three liters (1.0 to 1.5 liters each for insensible water loss by perspiration and urine production).

Water balance is maintained by various regulatory mechanisms. Water distribution depends primarily on the concentration of electrolytes in the body compartments and sodium (Na^+) plays a major role in maintaining physiologic equilibrium.

INDICATIONS AND USAGE

Intravenous solutions containing sodium chloride are indicated for parenteral replenishment of fluid and sodium chloride as required by the clinical condition of the patient.

CONTRAINDICATIONS

None known.

WARNINGS

Sodium Chloride Injection, USP should be used with great care, if at all, in patients with congestive heart failure, severe renal insufficiency and in clinical states in which there exists edema with sodium retention.

The intravenous administration of Sodium Chloride Injection, USP can cause fluid and/or solute overloading resulting in dilution of serum electrolyte concentrations, overhydration, congested states or pulmonary edema.

The risk of dilutive states is inversely proportional to the electrolyte concentration of the injections. The risk of solute overload causing congested states with peripheral and pulmonary edema is directly proportional to the electrolyte concentrations of the injections.

In patients with diminished renal function, administration of Sodium Chloride Injection, USP may result in sodium retention.

PRECAUTIONS

General

Do not use plastic containers in series connections. Such use could result in air embolism due to residual air being drawn from the primary container before administration of the fluid from the secondary container is completed.

Pressurizing intravenous solutions contained in flexible plastic containers to increase flow rates can result in air embolism if the residual air in the container is not fully evacuated prior to administration.

Use of a vented intravenous administration set with the vent in the open position could result in air embolism. Vented intravenous administration sets with the vent in the open position should not be used with flexible plastic containers.

Laboratory Tests

Clinical evaluation and periodic laboratory determinations are necessary to monitor changes in fluid balance, electrolyte concentrations and acid-base balance during prolonged parenteral therapy or whenever the condition of the patient warrants such evaluation.

DRUG INTERACTIONS

Caution must be exercised in the administration of Sodium Chloride Injection, USP to patients receiving corticosteroids or corticotropin.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Studies have not been performed with Sodium Chloride Injection, USP to evaluate the potential for carcinogenesis, mutagenesis or impairment of fertility.

Pregnancy:

Teratogenic Effects

Animal reproduction studies have not been conducted with Sodium Chloride Injection, USP. It is also not known whether Sodium Chloride Injection, USP can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Sodium Chloride Injection, USP should be given to a pregnant woman only if clearly needed.

Labor and Delivery

Studies have not been conducted to evaluate the effects of Sodium Chloride Injection, USP on labor and delivery. Caution should be exercised when administering this drug during labor and delivery.

Nursing Mothers

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when Sodium Chloride Injection, USP is administered to a nursing mother.

Pediatric Use

The use of Sodium Chloride Injection, USP in pediatric patients is based on clinical practice.

Plasma electrolyte concentrations should be closely monitored in the pediatric population as this population may have impaired ability to regulate fluids and electrolytes.

Geriatric Use

Clinical studies of Sodium Chloride Injection, USP did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or drug therapy.

This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

Do not administer unless solution is clear and container is undamaged. Discard unused portion.

ADVERSE REACTIONS

Reactions which may occur because of the solution or the technique of administration include febrile response, infection at the site of injection, venous thrombosis or phlebitis extending from the site of injection, extravasation and hypervolemia.

If an adverse reaction does occur, discontinue the infusion, evaluate the patient, institute appropriate therapeutic countermeasures and save the remainder of the fluid for examination if deemed necessary.

OVERDOSAGE

In the event of overhydration or solute overload, re-evaluate the patient and institute appropriate corrective measures (see **WARNINGS, PRECAUTIONS, and ADVERSE REACTIONS**).

DOSAGE AND ADMINISTRATION

The dose is dependent upon the age, weight and clinical condition of the patient.

Additives may be incompatible. Consult with pharmacist, if available. When introducing additives, use aseptic technique, mix thoroughly and do not store.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit (see **PRECAUTIONS**).

INSTRUCTIONS FOR USE

Check flexible container solution composition, lot number, and expiry date.

Do not remove solution container from its overwrap until immediately before use.

Use sterile equipment and aseptic technique.

To Open

1. Turn solution container over so that the text is face down. Using the pre-cut corner tabs, peel open the overwrap and remove solution container.
2. Check the solution container for leaks by squeezing firmly. If leaks are found, or if the seal is not intact, discard the solution.
3. Do not use if the solution is cloudy or a precipitate is present.

To Add Medication

1. Identify WHITE Additive Port with arrow pointing toward container.
2. Immediately before injecting additives, break off WHITE Additive Port Cap with the arrow pointing toward container.
3. Hold base of WHITE Additive Port horizontally.
4. Insert needle horizontally through the center of WHITE Additive Port's septum and inject additives.
5. Mix container contents thoroughly.

Preparation for Administration

1. Immediately before inserting the infusion set, break off BLUE Infusion Port Cap with the arrow pointing away from container.
2. Use a non-vented infusion set or close the air-inlet on a vented set.
3. Close the roller clamp of the infusion set.
4. Hold the base of BLUE Infusion Port.
5. Insert spike through BLUE Infusion Port by rotating wrist slightly until the spike is inserted.

NOTE: See full directions accompanying administration set.

WARNING: Do not use flexible container in series connections.

HOW SUPPLIED

0.9% Sodium Chloride Injection, USP is supplied in single-dose flexible plastic containers as follows:

Product No.	NDC No.	Strength	Fill Sizes	Bag Size
1727170102	17271-701-02	0.9% (9 mg/mL)	50 mL	100 mL
1727170103	17271-701-03	0.9% (9 mg/mL)	100 mL	100 mL
1727170105	17271-701-05	0.9% (9 mg/mL)	250 mL	250 mL
1727170106	17271-701-06	0.9% (9 mg/mL)	500 mL	500 mL
1727170107	17271-701-07	0.9% (9 mg/mL)	1,000 mL	1,000 mL

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from freezing. The container closure is not made with natural rubber latex. Non-PVC, Non-DEHP, Sterile.



Becton, Dickinson and Company

1 Becton Drive

Franklin Lakes, NJ 07417 USA

For product inquiry: 1-800-523-0502

Distributed by BD. Manufactured by Fresenius Kabi.

Revised: October 2022

PACKAGE LABEL - PRINCIPAL DISPLAY PANEL - Sodium Chloride 50 mL Bag Label

freeflex® NDC 17271-701-02 **50 mL**

0.9% Sodium Chloride Injection, USP

For intravenous use.

Rx only

free *flex*[®]

NDC 17271-701-02

50 mL

0.9% Sodium Chloride Injection, USP

For intravenous use.

Rx only

Each 100 mL contains: Sodium Chloride 900 mg in water for injection.

Electrolytes per 1,000 mL:

Sodium 154 mEq

Chloride 154 mEq

308 mOsmol/LITER (CALC.) pH 4.5 to 7.0

Single Dose Only. Discard Unused Portion.

Additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic technique, mix thoroughly and do not store. Use only if solution is clear and container is undamaged. Must not be used in series connections.

Usual dosage: See package insert.

STORE AT: 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from freezing.

The container closure is not made with natural rubber latex.

Non-PVC,
Non-DEHP,
Sterile.

R2



(01)00317271701025



Becton,
Dickinson
and
Company
1 Becton Drive

Franklin Lakes, NJ 07417 USA

For product inquiry: 1-800-523-0502

Distributed by BD. Manufactured by Fresenius Kabi.



EXP: MMYYYY

LOT: 14XXZZZZ

403919

1727170102

NDC 17271-701-02

0.9% Sodium Chloride Injection, USP

50 mL x 60

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from freezing.

NDC 17271-701-02

1727170102

0.9% Sodium Chloride Injection, USP

50 mL x 60

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from freezing.

LOT

EXP



Becton, Dickinson and Company
1 Becton Drive
Franklin Lakes, NJ 07417 USA

For product inquiry:

1-800-523-0502

Distributed by BD.

Manufactured by Fresenius Kabi.

QTY 60



(01)50317271701020

63940

PACKAGE LABEL - PRINCIPAL DISPLAY PANEL - Sodium Chloride 100 mL Bag Label

freeflex® NDC 17271-701-03 **100 mL**

0.9% Sodium Chloride Injection, USP

For intravenous use. Rx only

free flex®

NDC 17271-701-03

100 mL

0.9% Sodium Chloride Injection, USP

For intravenous use.

Rx only

Each 100 mL contains: Sodium Chloride 900 mg in water for injection.

Electrolytes per 1,000 mL:

Sodium 154 mEq

Chloride 154 mEq

308 mOsmol/LITER (CALC.) pH 4.5 to 7.0

Single Dose Only. Discard Unused Portion.

Additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic technique, mix thoroughly and do not store. Use only if solution is clear and container is undamaged. Must not be used in series connections.

Usual dosage: See package insert.

STORE AT: 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from freezing.

The container closure is not made with natural rubber latex.

Non-PVC,
Non-DEHP,
Sterile.

R2



(01)00317271701032



Becton,
Dickinson
and
Company
1 Becton Drive

Franklin Lakes, NJ 07417 USA

For product inquiry: 1-800-523-0502

Distributed by BD. Manufactured by Fresenius Kabi.



EXP: MMYYYY

LOT: 14XXZZZ

403921

1727170103

NDC 17271-701-03

0.9% Sodium Chloride Injection, USP

100 mL x 50

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from freezing.

NDC 17271-701-03

1727170103

0.9% Sodium Chloride Injection, USP

100 mL x 50

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from freezing.

LOT

EXP



Becton, Dickinson and Company
1 Becton Drive
Franklin Lakes, NJ 07417 USA

For product inquiry:

1-800-523-0502

Distributed by BD.

Manufactured by Fresenius Kabi.

QTY 50



(01)50317271701037

63941

**PACKAGE LABEL - PRINCIPAL DISPLAY PANEL - Sodium Chloride 250 mL Bag
Label**

freeflex[®] NDC 17271-701-05 **250 mL**

0.9% Sodium Chloride Injection, USP

For intravenous use.

Rx only

free flex®

NDC 17271-701-05

250 mL

0.9% Sodium Chloride Injection, USP

For intravenous use.

Rx only

Each 100 mL contains: Sodium Chloride
900 mg in water for injection.

Electrolytes per 1,000 mL:

Sodium 154 mEq

Chloride 154 mEq

308 mOsmol/LITER (CALC.) pH 4.5 to 7.0

50

Single Dose Only. Discard Unused Portion.

Additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic technique, mix thoroughly and do not store. Use only if solution is clear and container is undamaged. Must not be used in series connections.

100

Usual dosage: See package insert.

STORE AT: 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from freezing.

150

The container closure is not made with natural rubber latex.

Non-PVC, Non-DEHP, Sterile.

R2 →

200





(01)00317271701056



EXP: MMYYYY

LOT: 14XXZZZZ



403922

mL

Becton, Dickinson and Company

1727170105

1 Becton Drive

Franklin Lakes, NJ 07417 USA

For product inquiry: 1-800-523-0502

Distributed by BD. Manufactured by Fresenius Kabi.

PACKAGE LABEL - PRINCIPAL DISPLAY PANEL - Sodium Chloride 250 mL Case Label

NDC 17271-701-05

0.9% Sodium Chloride Injection, USP

250 mL x 30

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from freezing.

NDC 17271-701-05

1727170105

0.9% Sodium Chloride Injection, USP
250 mL x 30

QTY 30

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from freezing.

LOT

EXP



Becton, Dickinson and Company
1 Becton Drive
Franklin Lakes, NJ 07417 USA

For product inquiry:

1-800-523-0502

Distributed by BD.

Manufactured by Fresenius Kabi.



(01)50317271701051

63942

PACKAGE LABEL - PRINCIPAL DISPLAY PANEL - Sodium Chloride 500 mL Bag Label

freeflex® NDC 17271-701-06 **500 mL**

0.9% Sodium Chloride Injection, USP

For intravenous use.

Rx only

freeflex®

NDC 17271-701-06

500 mL

0.9% Sodium Chloride Injection, USP

For intravenous use.

Rx only

100

Each 100 mL contains: Sodium Chloride 900 mg
in water for injection.

Electrolytes per 1,000 mL:

Sodium 154 mEq

Chloride 154 mEq

308 mOsmol/LITER (CALC.) pH 4.5 to 7.0

200

Single Dose Only. Discard Unused Portion.

Additives may be incompatible. Consult with pharmacist.
When introducing additives, use aseptic technique, mix
thoroughly and do not store. Use only if solution is clear and
container is undamaged. Must not be used in series
connections.

300

Usual dosage: See package insert.

STORE AT: 20° to 25°C (68° to 77°F) [see USP Controlled
Room Temperature]. Protect from freezing.

The container closure is not made with natural rubber latex.
Non-PVC, Non-DEHP, Sterile.

400

R2



(01)00317271701063



EXP: MMYYYY

LOT: 14XXZZZZ

mL



Becton, Dickinson and Company

1 Becton Drive

Franklin Lakes, NJ 07417 USA

For product inquiry: 1-800-523-0502

403923

Distributed by BD. Manufactured by Fresenius Kabi.

1727170106

PACKAGE LABEL - PRINCIPAL DISPLAY PANEL - Sodium Chloride 500 mL Case Label

NDC 17271-701-06

0.9% Sodium Chloride Injection, USP

500 mL x 20

Store at 20° to 25°C (68° to

77°F) [see USP Controlled

Room Temperature]. Protect

from freezing.

NDC 17271-701-06

1727170106

0.9% Sodium Chloride Injection, USP
500 mL x 20

QTY 20

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from freezing.

LOT

EXP



Becton, Dickinson and Company
1 Becton Drive
Franklin Lakes, NJ 07417 USA

For product inquiry:

1-800-523-0502

Distributed by BD.

Manufactured by Fresenius Kabi.



(01)50317271701068

63943

PACKAGE LABEL - PRINCIPAL DISPLAY PANEL - Sodium Chloride 1,000 mL Bag Label

freeflex® NDC 17271-701-07 **1,000 mL**

0.9% Sodium Chloride Injection, USP

For intravenous use.

Rx only

100

freeflex[®]

NDC 17271-701-07

1,000 mL

0.9% Sodium Chloride Injection, USP

For intravenous use.

Rx only

200

Each 100 mL contains: Sodium Chloride 900 mg
in water for injection.

Electrolytes per 1,000 mL:

Sodium 154 mEq

Chloride 154 mEq

308 mOsmol/LITER (CALC.) pH 4.5 to 7.0

300

Single Dose Only. Discard Unused Portion.

Additives may be incompatible. Consult with pharmacist.
When introducing additives, use aseptic technique, mix
thoroughly and do not store. Use only if solution is clear and
container is undamaged. Must not be used in series
connections.

400

Usual dosage: See package insert.

STORE AT: 20° to 25°C (68° to 77°F) [see USP Controlled
Room Temperature]. Protect from freezing.

500

The container closure is not made with natural rubber latex.
Non-PVC, Non-DEHP, Sterile.

600

R2



(01)00317271701070



EXP: MMYYYY

LOT: 14XXZZZZ

700



Becton, Dickinson and Company
1 Becton Drive
Franklin Lakes, NJ 07417 USA

403924
1727170107

mL For product inquiry: 1-800-523-0502

Distributed by BD. Manufactured by Fresenius Kabi.

**PACKAGE LABEL - PRINCIPAL DISPLAY PANEL - Sodium Chloride 1,000 mL
Case Label**

NDC 17271-701-07

0.9% Sodium Chloride Injection, USP

1,000 mL x 10

Store at 20° to 25°C (68° to
77°F) [see USP Controlled
Room Temperature]. Protect
from freezing.

NDC 17271-701-07

1727170107

0.9% Sodium Chloride Injection, USP
1,000 mL x 10

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from freezing.

LOT

EXP



Becton, Dickinson and Company
1 Becton Drive
Franklin Lakes, NJ 07417 USA

For product inquiry:

1-800-523-0502

Distributed by BD.

Manufactured by Fresenius Kabi.

QTY 10



(01)50317271701075

63944

SODIUM CHLORIDE

sodium chloride injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:17271-701
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of	Strength
-----------------	----------	----------

Ingredient Name		Strength	Strength	
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32Z N48698)		SODIUM CHLORIDE	9 mg in 1 mL	
Inactive Ingredients				
Ingredient Name		Strength		
water (UNII: 059QF0KO0R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17271-701-02	50 mL in 1 BAG; Type 0: Not a Combination Product	09/19/2017	
2	NDC:17271-701-03	100 mL in 1 BAG; Type 0: Not a Combination Product	09/19/2017	
3	NDC:17271-701-05	250 mL in 1 BAG; Type 0: Not a Combination Product	09/19/2017	
4	NDC:17271-701-06	500 mL in 1 BAG; Type 0: Not a Combination Product	09/19/2017	
5	NDC:17271-701-07	1000 mL in 1 BAG; Type 0: Not a Combination Product	09/19/2017	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA207310	09/19/2017	

Labeler - Becton Dickinson and Company (124987988)

Establishment

Name	Address	ID/FEI	Business Operations
Fresenius Kabi Deutschland GmbH		506719546	analysis(17271-701) , manufacture(17271-701) , api manufacture(17271-701)

Establishment

Name	Address	ID/FEI	Business Operations
HP Halden Pharma AS		347747373	analysis(17271-701) , manufacture(17271-701) , pack(17271-701)

Establishment

Name	Address	ID/FEI	Business Operations
Fresenius Medical Care North America		078401205	analysis(17271-701) , label(17271-701) , manufacture(17271-701) , pack(17271-701) , sterilize(17271-701)

Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

Fresenius Kabi USA, LLC		964475045	analysis(17271-701) , manufacture(17271-701)
-------------------------	--	-----------	--

Revised: 5/2024

Becton Dickinson and Company