

**IBUPROFEN- ibuprofen tablet, film coated**  
**Pharbest Pharmaceuticals, Inc.**

-----  
**Drug Facts**

**Active ingredient (in each tablet)**

Ibuprofen 200 mg (NSAID)\*

\*nonsteroidal anti- inflammatory drug

**Purpose**

Pain Reliever/ Fever Reducer

**Uses**

- temporarily relieves minor aches and pain due to:
- headache
- toothache
- backache
- menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to Aspirin.

Symptoms may include

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood-thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/ fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**when using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**stop use and ask a doctor if**

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- Do not take more than directed
- The smallest effective dose should be used

adults and children 12 years of age	Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours unless directed by a doctor.
-------------------------------------	---

and over	exceed 6 tablets in 24 hours unless directed by a doctor
children under 12 years	consult a doctor

### Other information

- **Tamper Evident: do not use if safety seal under cap is broken or missing**
- store at room temperature (20°- 25°C)
- avoid excessive heat above 40°C (104°F)

### Inactive ingredients

carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, stearic acid, sodium starch glycolate, titanium dioxide

### Questions or comments?

(866) 562-2756 Mon-Fri: 8 AM to 4 PM

**PHARBEST**

**NDC 16103-407-11**

**\*Compare to the active ingredient in ADVIL®**

**Pain Reliever/Fever Reducer (NSAID)**

**Ibuprofen**

**Tablets**

**200 mg**

**1000 Tablets**

<b>This Package for Households Without Young Children</b>	<b>Drug Facts</b> <b>Active ingredient (in each tablet)</b> <b>Purpose</b> Ibuprofen 200mg (NSAID).....Pain Reliever/ Fever Reducer <small>*nonsteroidal anti-inflammatory drug</small>	<b>Drug Facts (continued)</b> ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed <b>Stop use and ask a doctor if</b> ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear <b>If pregnant or breast-feeding,</b> ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).
	<b>Uses</b> ■ temporarily relieves minor aches and pains due to: ■ headache ■ toothache ■ backache ■ menstrual cramps ■ the common cold ■ muscular aches ■ minor pain of arthritis ■ temporarily reduces fever	<b>Directions</b> ■ Do not take more than directed ■ The smallest effective dose should be used adults and children 12 years of age and over      Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours unless directed by a doctor. children under 12 years      consult a doctor
	<b>Warnings</b> <b>Allergy alert:</b> Ibuprofen may cause a severe allergic reaction, especially in people allergic to Aspirin. Symptoms may include ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters If an allergic reaction occurs, stop use and seek medical help right away. <b>Stomach bleeding warning:</b> This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood-thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed <b>Do not use</b> ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery	<b>Other information</b> ■ <b>Tamper Evident: do not use if safety seal under cap is broken or missing</b> ■ store at room temperature (20°-25°C) ■ avoid excessive heat above 40°C (104°F) <b>Inactive ingredients</b> carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, stearic acid, sodium starch glycolate, titanium dioxide
	<b>Ask a doctor before use if</b> ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma ■ you are taking a diuretic <b>Ask a doctor or pharmacist before use if you are</b> ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug <b>When using this product</b> ■ take with food or milk if stomach upset occurs	<b>Questions or comments?</b> (866) 562-2756 Mon-Fri: 8 AM to 4 PM

<b>IBUPROFEN</b>			
ibuprofen tablet, film coated			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:16 103-407

<b>Route of Administration</b>	ORAL
--------------------------------	------

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

### Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	brown	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;291
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:16103-407-06	1 in 1 CARTON	10/01/2018	
1		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:16103-407-08	1 in 1 CARTON	10/01/2018	
2		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:16103-407-11	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/01/2018	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

ANDA	ANDA075010	10/01/2018	
------	------------	------------	--

**Labeler** - Pharbest Pharmaceuticals, Inc. (557054835)

**Registrant** - Pharbest Pharmaceuticals, Inc. (557054835)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Pharbest Pharmaceuticals, Inc.		557054835	relabel(16103-407) , repack(16103-407)

Revised: 10/2018

Pharbest Pharmaceuticals, Inc.