


## **NYSTATIN- nystatin cream cream** **Cosette Pharmaceuticals, Inc.**

-----

### **DESCRIPTION**

Nystatin is a polyene antifungal antibiotic obtained from *Streptomyces noursei*.

Structural formula:



**C<sub>47</sub>H<sub>75</sub>NO<sub>17</sub> Molecular Weight: 926.13**

Nystatin Cream, USP is for dermatologic use.

Nystatin cream for topical use, contains 100,000 USP nystatin units per gram in a cream base containing aluminum hydroxide gel, cetareth-15, glyceryl monostearate, polyethylene glycol 400 monostearate, propylene glycol, purified water, simethicone emulsion, sorbitol solution, titanium dioxide, white petrolatum, methylparaben, propylparaben, and sodium hydroxide.

### **CLINICAL PHARMACOLOGY**

#### **Pharmacokinetics**

Nystatin is not absorbed from intact skin or mucous membrane.

#### **Microbiology**

Nystatin is an antibiotic which is both fungistatic and fungicidal *in vitro* against a wide variety of yeasts and yeast-like fungi, including *Candida albicans*, *C. parapsilosis*, *C. tropicalis*, *C. guilliermondi*, *C. pseudotropicalis*, *C. krusei*, *Torulopsis glabrata*, *Trichophyton rubrum*, *T. mentagrophytes*. Nystatin acts by binding to sterols in the cell membrane of susceptible species resulting in a change in membrane permeability and the subsequent leakage of intracellular components. On repeated subculturing with increasing levels of nystatin, *Candida albicans* does not develop resistance to nystatin. Generally, resistance to nystatin does not develop during therapy. However, other species of *Candida* (*C. tropicalis*, *C. guilliermondi*, *C. krusei*, and *C. stellatoides*) become quite resistant on treatment with nystatin and simultaneously become cross resistant to amphotericin as well. This resistance is lost when the antibiotic is removed. Nystatin exhibits no

appreciable activity against bacteria, protozoa, or viruses.

## **INDICATIONS AND USAGE**

Nystatin Cream, USP is indicated in the treatment of cutaneous or mucocutaneous mycotic infections caused by *Candida albicans* and other susceptible *Candida* species.

**This cream is not indicated for systemic, oral, intravaginal or ophthalmic use.**

## **CONTRAINDICATIONS**

Nystatin cream is contraindicated in patients with a history of hypersensitivity to any of its components.

## **PRECAUTIONS**

### **General**

**Nystatin cream should not be used for the treatment of systemic, oral, intravaginal or ophthalmic infections.**

If irritation or sensitization develops, treatment should be discontinued and appropriate measures taken as indicated. It is recommended that KOH smears, cultures, or other diagnostic methods be used to confirm the diagnosis of cutaneous or mucocutaneous candidiasis and to rule out infection caused by other pathogens.

## **INFORMATION FOR THE PATIENT**

Patients using this medication should receive the following information and instructions:

1. The patient should be instructed to use this medication as directed (including the replacement of missed doses). This medication is not for any disorder other than that for which it is prescribed.
2. Even if symptomatic relief occurs within the first few days of treatment, the patient should be advised not to interrupt or discontinue therapy until the prescribed course of treatment is completed.
3. If symptoms of irritation develop, the patient should be advised to notify the physician promptly.

### **Laboratory Tests**

If there is a lack of therapeutic response, KOH smears, cultures or other diagnostic methods should be repeated.

### **Carcinogenesis, Mutagenesis, and Impairment of Fertility**

No long-term animal studies have been performed to evaluate the carcinogenic potential of nystatin. No studies have been performed to determine the mutagenicity of nystatin

or the effects on male or female fertility.

## **Pregnancy**

Teratogenic Effects

### *Category C*

Animal reproduction studies have not been conducted with any nystatin cream. It also is not known whether this cream can cause fetal harm when used by a pregnant woman or can affect reproductive capacity. Nystatin cream should be prescribed for a pregnant woman only if the potential benefit to the mother outweighs the potential risk to the fetus.

## **Nursing Mothers**

It is not known whether nystatin is excreted in human milk. Caution should be exercised when nystatin is prescribed for a nursing woman.

## **Pediatric Use**

Safety and effectiveness have been established in the pediatric population from birth to 16 years.

(See **DOSAGE AND ADMINISTRATION.**)

## **Geriatric Use**

Clinical studies with nystatin cream did not include sufficient numbers of subjects aged 65 years and older to determine whether they respond differently than younger subjects. Other reported clinical experience has not identified differences in responses between elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

## **ADVERSE REACTIONS**

The frequency of adverse events reported in patients using nystatin cream is less than 0.1%. The more common events that were reported include allergic reactions, burning, itching, rash, eczema, and pain on application.

(See **PRECAUTIONS: General.**)

**To report SUSPECTED ADVERSE REACTIONS, contact Cosette Pharmaceuticals, Inc. at 1-800-922-1038 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

## **DOSAGE AND ADMINISTRATION**

## **Adults and Pediatric Patients (Neonates and Older)**

Apply liberally to affected areas twice daily or as indicated until healing is complete.

### **HOW SUPPLIED**

Nystatin Cream, USP is a smooth yellow cream.

Nystatin Cream, USP is supplied in 15 g (NDC 0713-0678-15) and 30 g (NDC 0713-0678-31) tubes providing 100,000 USP Nystatin units per gram.

**Store at 20° to 25°C (68° to 77°F)** [see USP Controlled Room Temperature]. Avoid freezing.

### **Distributed by:**

Cosette Pharmaceuticals, Inc.  
South Plainfield, NJ 07080

Rev. 03/2022  
8-0678CPLNC2 VC7625

### **PRINCIPAL DISPLAY PANEL**

**NDC 0713-0678-15**

**Nystatin Cream, USP**

**15 g**

**Rx only**

**FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.**

**Cosette Pharmaceuticals, Inc.**



# Nystatin Cream, USP

NDC 0713-0678-15

Rx only

**100,000 units per gram**

**15 g**

FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.

**Usual Dosage:** Apply liberally to affected area twice daily.  
See package insert for full prescribing information.

**Distributed by:**

Cosette Pharmaceuticals, Inc.  
South Plainfield, NJ 07080  
10-67815CPLNC2 Rev. 03/2022 VC7626

GTIN 00307130678155



# Nystatin Cream, USP

NDC 0713-0678-15

Rx only

**100,000 units per gram**

**15 g**

FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.

**Each gram contains:** 100,000 USP Nystatin Units in a cream base containing aluminum hydroxide gel, osteareth-15, glyceryl monostearate, polyethylene glycol 400 monostearate, propylene glycol, purified water, simethicone emulsion, sorbitol solution, titanium dioxide, white petrolatum, methylparaben, propylparaben, and sodium hydroxide.

**Store at 20° to 25°C (68° to 77° F) [see USP Controlled Room Temperature]. Avoid freezing.**

**KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.**

See carton or crimp of tube for lot number and expiration date.

FPO 7626

FPO 7626



NDC 0713-0678-15

# Nystatin Cream, USP

Rx only

**100,000 units per gram**

**15 g**

**FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.**

**Usual Dosage:** Apply liberally to affected area twice daily.

See package insert for full prescribing information.

**Each gram contains:** 100,000 USP Nystatin Units in a cream base containing aluminum hydroxide gel, cetareth-15, glyceryl monostearate, polyethylene glycol 400 monostearate, propylene glycol, purified water, simethicone emulsion, sorbitol solution, titanium dioxide, white petrolatum, methylparaben, propylparaben, and sodium hydroxide.

Store at 20° to 25°C (68° to 77°F)

[See USP Controlled Room Temperature]. Avoid freezing.

**KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.**

See carton or crimp of tube for lot number and expiration date.

**Distributed by:** Cosette Pharmaceuticals, Inc.

South Plainfield, NJ 07080

11-67815CPLNC1 Iss. 08/2020 VC126



FP0 126

Read Direction

**NDC 0713-0678-31**


**Nystatin Cream, USP**

**30 g**

**Rx only**

**FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.**

**Cosette Pharmaceuticals, Inc.**



**NYSTATIN**  
nystatin cream cream

**Product Information**

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0713-0678
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
NYSTATIN (UNII: BDF1O1C72E) (NYSTATIN - UNII:BDF1O1C72E)	NYSTATIN	100000 [USP'U] in 1 g

**Inactive Ingredients**

Ingredient Name	Strength
ALGELDRATE (UNII: 03J11K103C)	
CETEARETH-15 (UNII: 867H4YOZ8Z)	
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SORBITOL (UNII: 506T60A25R)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
PETROLATUM (UNII: 4T6H12BN9U)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	

**Product Characteristics**

<b>Color</b>	yellow	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0713-0678-15	1 in 1 CARTON	04/01/2015	
1		15 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:0713-0678-31	1 in 1 CARTON	04/01/2015	
2		30 g in 1 TUBE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date



ANDA	ANDA061966	04/01/2015	
------	------------	------------	--

**Labeler** - Cosette Pharmaceuticals, Inc. (116918230)

**Registrant** - Cosette Pharmaceuticals, Inc. (116918230)

**Establishment**

Name	Address	ID/FEI	Business Operations
Cosette Pharmaceuticals NC Laboratories, LLC		079419931	analysis(0713-0678) , label(0713-0678) , manufacture(0713-0678) , pack(0713-0678)

Revised: 11/2022

Cosette Pharmaceuticals, Inc.