VICKS NYQUIL ALCOHOL FREE COLD AND FLU NIGHTTIME RELIEFacetaminophen, chlorpheniramine maleate, and dextromethorphan hydrobromide liquid

The Procter & Gamble Manufacturing Company

VICKS ® NyQuil Alcohol Free COLD & FLU

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg Chlorpheniramine maleate 4 mg Dextromethorphan HBr 30 mg

Purpose

Pain reliever/fever reducer Antihistamine Cough suppressant

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not

notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & 30 mL every 6 over hrs children 4 to under 12 yrs ask a doctor children under 4 yrs do not use

Other information

- each 30 mL dose cup contains:sodium 25 mg
- store at no greater than 25°C and do not refrigerate

Inactive ingredients

citric acid, FD&C Red No. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions?

1-800-362-1683

TAMPER EVIDENT: Do not use if printed shrinkband is broken or missing.

DIST. BY PROCTER & GAMBLE, CINCINNATI OH 45202

PRINCIPAL DISPLAY PANEL - 354 ml Bottle Label

VICKS®

NyQuil™

Alcohol Free

COLD & FLU

Acetaminophen, Chlorpheniramine maleate, Dextromethorphan HBr Headache, Fever, Sore Throat, Minor Aches & Pains Sneezing, Runny Nose


Cough


12 FL OZ (354 ml)

Nighttime Relief Alcohol Free









VICKS NYQUIL ALCOHOL FREE COLD AND FLU NIGHTTIME RELIEF

acetaminophen, chlorpheniramine maleate, and dextromethorphan hydrobromide liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69423-899
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	650 mg in 30 mL	
CHLORPHENIRAMINE MALEATE (UNII: V1Q0090J9Z) (CHLORPHENIRAMINE - UNII: 3U6I01965U)	CHLORPHENIRAMINE MALEATE	4 mg in 30 mL	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL	

Inactive Ingredients			
Ingredient Name	Strength		
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			
XANTHAN GUM (UNII: TTV12P4NEE)			
GLYCERIN (UNII: PDC6A3C0OX)			

WATER (UNII: 059QF0KO0R) SACCHARIN SODIUM (UNII: SB8ZUX40TY) SODIUM BENZOATE (UNII: OJ245FE5EU) SODIUM CITRATE (UNII: 1Q73Q2JULR)
SODIUM BENZOATE (UNII: OJ245FE5EU)
SODIUM CITRATE (UNII: 107302IULR)
SUCRALOSE (UNII: 96K6UQ3ZD4)
SORBITOL (UNII: 506T60A25R)

Product Characteristics			
Color	red	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging						
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
		NDC:69423- 899-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/20/2021		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/20/2021	

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 6/2024 The Procter & Gamble Manufacturing Company