

**SULFASOL - sulfadimethoxine sodium powder, for solution**  
**Dechra Veterinary Products**

-----  
**SULFASOL SOLUBLE POWDER**

(Brand of Sulfadimethoxine)

**Antibacterial**

**CAUTION:** Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

Approved by FDA under ANADA #200-238

Restricted Drug (California) - Use Only as Directed. Not for Human Use.

**Net Wt.: 3.77 oz (107g)**

**Each packet contains 3.34 oz. (94.6 g) sulfadimethoxine in the form of the soluble sodium salt and disodium edetate.**

**INDICATIONS FOR USE:**

**For Broiler and Replacement Chickens Only** - Use for the treatment of disease outbreaks of coccidiosis, fowl cholera, and infectious coryza.

**For Meat-producing Turkeys Only** - Use for the treatment of disease outbreaks of coccidiosis, and fowl cholera.

**For Dairy Calves, Dairy Heifers, and Beef Cattle** - Use for the treatment of shipping fever complex and bacterial pneumonia associated with *Pasteurella* spp. sensitive to sulfadimethoxine; and calf diphtheria and foot rot associated with *Sphaerophorus necrophorus* sensitive to sulfadimethoxine.

		<b>SULFADIMETHOXINE</b>	
		SOLUBLE POWDER	
<b>Dosage and Administration:</b>			
<b>Species</b>	<b>Concentration</b>	<b>Use of Directions</b>	
Chickens	0.050%	Contents of packet to 50 gal of water	
Turkeys	0.025%	Contents of packet to 100 gal of water	
<b>Automatic Proportioners:</b> To make stock solution, add contents of 5 packets to 2 gal of water for chickens and to 4 gal of water for turkeys. Set proportioner to feed at a rate of 1 fl oz stock solution per gal of water.			
<b>Treatment Period: 6 consecutive days</b>			
<b>Dairy Calves, Dairy Heifers, and Beef Cattle</b>			
<b>Dosage:</b> 25 mg/lb first day followed by 12.5 mg/lb/day for 4 days			
		<b>Sulfadimethoxine in Water</b>	
		<b>Water Consumption</b>	
	<b>Amount of Stock Solution for Cattle*</b>	<b>(Summer)* 1 gal/100 lb body weight**</b>	<b>(Winter)* 1 gal/150 lb body weight**</b>

<b>First Day Add</b>	1 qt	10 gal	7 gal
	2 qt	20 gal	14 gal
	1 gal	40 gal	28 gal
<b>Next 4 Days Add</b>	1 qt	20 gal	14 gal
	2 qt	40 gal	28 gal
	1 gal	80 gal	56 gal

**\*Note:** Make a cattle stock solution by adding 1 packet of Sulfadimethoxine Soluble Powder to 1 gal of water.

\*Twenty fl oz of cattle stock solution will medicate 1-600 lb animal initially or 2-600 lb animals on maintenance dose. Contents of packet will medicate 6-600 lb animals initially or 12-600 lb animals on maintenance dose.

\*\*This dosage recommendation is based on a water consumption of 1 gal per 100 lb of body weight per day, the expected water consumption rate for summer. Water consumption during cold months (winter) may drop markedly (30%-40%). Accordingly, adjustments must be made in the dilution rates to compensate for this and insure proper drug intake.

For treatment of individual cattle, Sulfadimethoxine Soluble Powder stock solution for cattle may be given as a drench.

**Treatment Period: 5 consecutive days**

### **CAUTION:**

*Chickens and Turkeys* - If animals show no improvement within 5 days, discontinue treatment and reevaluate diagnosis. Prepare a fresh stock solution daily. Handle the recommended dilutions (chickens 0.05% and turkeys 0.025%) as regular drinking water. Administer as sole source of drinking water and sulfonamide medication. Chickens and turkeys that have survived fowl cholera outbreaks should not be kept for replacements or breeders.

*Cattle* - During treatment period, make certain that animals maintain adequate water intake. If animals show no improvement within 2 or 3 days, reevaluate diagnosis. Treatment should not be continued beyond 5 days.

### **WARNINGS:**

*Chickens and Turkeys* - Withdraw 5 days before slaughter. Do not administer to chickens over 16 weeks (112 days) of age or to turkeys over 24 weeks (168 days) of age.

*Cattle* - Withdraw 7 days before slaughter. For dairy calves, dairy heifers and beef cattle only.

A withdrawal period has not been established for this product in pre-ruminating calves.

### **Do Not Use in Calves to be Processed For Veal**

**CONTACT INFORMATION:** To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Dechra at (866) 933-2472. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at <http://www.fda.gov/reportanimalae>

SULFASOL is a trademark of Med-Pharmex, Inc.

### **Manufactured for:**

NDC 17033-428-03

# SULFASOL™

## (sulfadimethoxine)

### Antibacterial SOLUBLE POWDER

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Each packet contains 3.34 oz. (94.6 g) sulfadimethoxine in the form of the soluble sodium salt and disodium edetate.

**Restricted Drug (California) - Use Only as Directed. Not for Use in Humans.**

**For Broiler and Replacement Chickens Only** - Use for the treatment of disease outbreaks of coccidiosis, fowl cholera, and infectious coryza.

**For Meat-producing Turkeys Only** - Use for the treatment of disease outbreaks of coccidiosis, and fowl cholera.

**For Dairy Calves, Dairy Heifers, and Beef Cattle** - Use for the treatment of shipping fever complex and bacterial pneumonia associated with *Pasteurella* spp. sensitive to sulfadimethoxine; and calf diphtheria and foot rot associated with *Sphaerophorus necrophorus* sensitive to sulfadimethoxine.

Approved by FDA under ANADA # 200-238

**Net Wt.: 3.77 oz (107g)**



### SULFADIMETHOXINE SOLUBLE POWDER

Species	Concentration	Use of Directions
Chickens	0.05%	Contents of packet to 50 gal of water
Turkeys	0.025%	Contents of packet to 100 gal of water

**Automatic Proportions:** To make stock solution, add contents of 5 packets to 2 gal of water for chickens and to 4 gal of water for turkeys. Set proportioner to feed at a rate of 1 lb of stock solution per gal of water.  
**Treatment Period:** 6 consecutive days

Dairy Calves, Dairy Heifers, and Beef Cattle
Dosage: 25 mg/100 lb first day followed by 12.5 mg/100 lb for 4 days

	Sulfadimethoxine in Water		
	Amount of Stock Solution for Cattle*	Summer** 1 gal/100 lb body weight**	Winter** 1 gal/150 lb body weight**
First Day Add	1 qt	10 gal	7 gal
	2 qt	20 gal	14 gal
Next 4 Days Add	1 qt	40 gal	28 gal
	2 qt	80 gal	56 gal
	1 gal	80 gal	56 gal

\*Note: Make a cattle stock solution by adding 1 packet of Sulfadimethoxine Soluble Powder to 1 gal of water. Twenty 8 oz of cattle stock solution will medicate 1-800 lb animal initially or 2-800 lb animals on maintenance dose. Contents of packet will medicate 6-800 lb animals initially or 12-800 lb animals on maintenance dose.

\*\*This dosage recommendation is based on a water consumption of 1 gal per 100 lb of body weight per day, the expected water consumption rate for summer. Water consumption during cold months (winter) may drop markedly (30%-40%). Accordingly, adjustments must be made in the dilution rates to compensate for this and insure proper drug intake.

For treatment of individual cattle, Sulfadimethoxine Soluble Powder stock solution for cattle may be given as a drench. **Treatment Period:** 5 consecutive days

**Caution: Chickens and Turkeys** - If animals show no improvement within 5 days, discontinue treatment and reevaluate diagnosis. Prepare a fresh stock solution daily. Handle the recommended dilutions (chickens 0.05% and turkeys 0.025%) as regular drinking water. Administer as sole source of drinking water and substitute medication. Chickens and turkeys that have survived foot cholera outbreaks should not be kept for replacements or breeders.  
**Cattle** - During treatment period, make certain that animals maintain adequate water intake. If animals show no improvement within 2 or 3 days, reevaluate diagnosis. **Treatment should not be continued beyond 5 days.**

**Warning:**  
Chickens and Turkeys - Withdraw 5 days before slaughter. Do not administer to chickens over 16 weeks (112 days) of age or to turkeys over 24 weeks (168 days) of age.  
Cattle - Withdraw 7 days before slaughter. For dairy calves, dairy heifers and beef cattle only. A withdrawal period has not been established for this product in pre-ruminating calves.  
**Do Not Use in Calves to be Processed For Veal**

**CONTACT INFORMATION:** To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Dechra at (888) 833-2422.  
For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-800-FDA-VETS or online at <http://www.fda.gov/reportadverse>.  
SULFASOL is a trademark of NeoPharm, Inc.  
MANUFACTURED FOR: Dechra Veterinary Products, Overland Park, KS 66211 USA



Rev. April 2023

## SULFASOL

sulfadimethoxine sodium powder, for solution

### Product Information

<b>Product Type</b>	PRESCRIPTION ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:17033-428
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>SULFADIMETHOXINE SODIUM</b> (UNII: 49DG2B481W) (SULFADIMETHOXINE - UNII:30CPC5LDEX)	SULFADIMETHOXINE SODIUM	94.6 g in 107 g

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17033-428-03	107 g in 1 PACKET		

### Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANADA	ANADA200238	05/17/2023	

**Labeler** - Dechra Veterinary Products (362142734)

Revised: 5/2023

Dechra Veterinary Products