# SOOTHE ULTRA STRENGTH- bismuth subsalicylate liquid Walgreens

-----

# **Drug Facts**

# Active ingredient (in each 15 mL)

Bismuth subsalicylate 525 mg

# **Purpose**

Upset stomach reliever/Antidiarrheal

#### Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
  - heartburn
  - indigestion
  - nausea
  - gas
  - fullness
  - belching

# Warnings

**Reye's syndrome**: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms ahould not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

# Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

#### Do not use

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

# Ask a doctor before use if you have

- fever
- mucus in the stool

## Ask a doctor or pharmacist before use if you are

taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

## When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur.

# Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

# If pregnant or breast-feeding,

ask a health professional before use.

# Keep out of reach of childen.

In case of overdose, get medical help or contact a Posion Control Center (1-800-222-1222) right away.

#### **Directions**

- do not take more than 8 doses (120 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- mL = milliliter
- shake well before using
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- adults and children 12 years and over:
  - 15 mL (1dose) every 1/2 or 30 mL (2 doses) every hour as needed diarrhea/traveler's diarrhea
  - 15 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- children under 12 years of age: ask a doctor

#### Other information

each 15 mL contains: sodium 5 mg
each 15 mL contains: salicylate 206 mg

- low sodium
- keep tightly closed

- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

# **Inactive ingredients**

benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

# **Questions or comments?**

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

# **Principal Display Panel**

Compare to Pepto-Bismol® Ultra active ingredient††

Soothe

BISMUTH SUBSALICYLATE 525 mg /

UPSET STOMACH RELIEVER / ANTIDIARRHEAL

**ULTRA STRENGTH** 

#### SUGAR FREE ALCOHOL FREE

- Provides protective coating action to relieve heartburn, indigestion, nausea, upset stomach, & diarrhea
- 2x strength per ounce\*

FL OZ (mL)

\*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

# TAMPER EVIDENT; DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY: WALGREEN CO., 200 WILMOT RD., DEERFIELD, IL 60015

# Package Label



# Drug Facts (continued) When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur. Stop use and ask a doctor if symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. **Directions** ■ do not take more than 8 doses (120 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea ■ mL = milliliter ■ shake well before using ■ measure only with dosing cup provided. Do not use any other dosing device. ■ keep dosing cup with product ■ adults and children 12 years and over: ■ 15 mL (1 dose) every ½ hour or 30 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea 15 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) children under 12 years of age: ask a doctor Other information ■ each 15 mL contains: sodium 5 mg ■ each 15 mL contains: salicylate 206 mg ■ low sodium ■ keep tightly closed ■ protect from freezing ■ avoid excessive heat (over 104°F or 40°C) Inactive ingredients benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

#### WALGREENS Soothe

# SOOTHE ULTRA STRENGTH

bismuth subsalicylate liquid

# Product Information Product Type HUMAN OTC DRUG Item Code (Source) NDC:0363-7370 Route of Administration ORAL

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD118YE, SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 15 mL		

Inactive Ingredients		
	Ingredient Name	Strength

BENZOIC ACID (UNII: 85KN0B0MIM)	
D&C RED NO. 22 (UNII: 1678RKX8RT)	
<b>D&amp;C RED NO. 28</b> (UNII: 767IP0Y5NH)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0KO0R)	
SUCRALOSE (UNII: 96K6UQ3Z D4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging			
# Item Code	Package Description	Marketing Start Date	Marketing End Date
<b>1</b> NDC:0363-7370-04	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/31/2020	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M008	01/31/2020		

# Labeler - Walgreens (008965063)

Revised: 10/2023 Walgreens