

**BISACODYL- bisacodyl tablet, delayed release**  
**Apnar Pharma LP**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Bisacodyl Delayed Release Tablets USP 5 mg- Pink**

***Active ingredient (in each tablet)***

Bisacodyl (USP) 5 mg

***Purpose***

Stimulant laxative

***Uses***

- for relief of occasional constipation and irregularity
- this product generally produces bowel movement in 6 to 12 hours

***Warnings***

**Do not use**

- if you cannot swallow without chewing

**Ask a doctor before use if you have:**

- stomach pain, nausea, or vomiting
- noticed a sudden change in bowel habits that lasts more than 2 weeks

**When using this product**

- it may cause stomach discomfort, faintness and cramps
- do not chew or crush tablets(s)
- do not use within 1 hour after taking antacid or milk

**Stop use and ask a doctor if**

- you have rectal bleeding or fail to have a bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

**If pregnant or breast-feeding,**

ask a health care professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- take with glass of water
- adult and children 12 years of age and over- 1 to 3 tablets in a single daily dose
- children 6 to under 12 years of age- 1 tablet in a single daily dose
- children under 6 years of age- ask a doctor

**Other Information**

- store at 15°-30°C (59° - 86° F).

**Inactive ingredients**

Lactose monohydrate, Anhydrous lactose, Pregelatinized starch, Magnesium stearate, Hypromellose, Triacetin, Talc, Methacrylic acid-ethyl acrylate copolymer (1:1) Type-A, Titanium dioxide, Triethyl citrate, D&C red no. 27, Colloidal anhydrous silica, Sodium bicarbonate, Sodium lauryl sulfate, FD&C blue no. 1, Carboxymethyl cellulose sodium, Maltodextrin, Dextrose monohydrate, Lecithin Soya.

**Questions or comments?**

- 1-855-642-2594

Distributed By:

**APNAR PHARMA LP**

CHINO, CA, 91710

Rev.: 07/2022

Compare to **Dulcolax Pink**®

active ingredient\*

\*This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., owner of the registered trademark of Dulcolax Pink®

**Caution:** Do not use if imprinted seal under cap is broken or missing.

**NDC 24689-129-02**

Bisacodyl Delayed Release Tablets USP

5 mg- Pink

Stimulant Laxative

20 Tablets

NDC 24689-129-02

# Bisacodyl Delayed Release Tablets USP

5 mg-Pink

STIMULANT LAXATIVE

Compare to Dulcolax Pink®  
active ingredient\*

20 Tablets



Distributed By:  
APNAR PHARMA LP  
CHINO, CA, 91710

Caution: Do not use if imprinted seal under cap is broken or missing.

### Drug Facts

**Active ingredient (in each tablet)**  
Bisacodyl (USP) 5 mg.....Stimulant laxative

**Purpose**  
Stimulant laxative

**Uses** • for relief of occasional constipation and irregularity • this product generally produces bowel movement in 6 to 12 hours

### Warnings

**Do not use** • if you cannot swallow without chewing

**Ask a doctor before use if you have:**

- stomach pain, nausea or vomiting • noticed a sudden change in bowel habits that lasts more than 2 weeks

### When using this product

• it may cause stomach discomfort, faintness and cramps • do not chew or crush tablets(s) • do not use within 1 hour after taking antacid or milk

### Stop use and ask a doctor if

• you have rectal bleeding or fail to have a bowel movement after using this product. These could be signs of a serious condition. • you need to use a laxative for more than 1 week

**if pregnant or breast-feeding**, ask a health care professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

• take with glass of water

• adult and children 12 years of age and over – 1 to 3 tablets in a single daily dose • children 6 to under 12 years of age – 1 tablet in a single daily dose • children under 6 years of age – ask a doctor

### Other information

• Store at 15° - 30°C (59° - 86° F).

**Inactive ingredients** Lactose monohydrate, Anhydrous lactose, Pregelatinized starch, Magnesium stearate, Hypromellose, Triacetin, Talc, Methacrylic acid-ethyl acrylate copolymer (1:1) Type-A, Titanium dioxide, Triethyl citrate, D&C red no. 27, Colloidal anhydrous silica, Sodium bicarbonate, Sodium lauryl sulfate, FD&C blue no. 1, Carboxymethyl cellulose sodium, Maltodextrin, Dextrose monohydrate, Lecithin Soya. 300072

**Questions or comments?** • 1-855-642-2594 Rev.: 07/2022

\* This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., owner of the registered trademark of Dulcolax Pink®



Unvarnished Area/Coding Area  
(7 x 25 mm)

## NDC 24689-129-03

Bisacodyl Delayed Release Tablets USP

5 mg- Pink

Stimulant Laxative

100 Tablets

NDC 24689-129-03

# Bisacodyl Delayed Release Tablets USP

5 mg-Pink

STIMULANT LAXATIVE

Compare to Dulcolax Pink®  
active ingredient\*

100 Tablets



Distributed By:  
APNAR PHARMA LP  
CHINO, CA, 91710

Caution: Do not use if imprinted seal under cap is broken or missing.

### Drug Facts

**Active ingredient (in each tablet)**  
Bisacodyl (USP) 5 mg.....Stimulant laxative

**Purpose**  
Stimulant laxative

**Uses** • for relief of occasional constipation and irregularity • this product generally produces bowel movement in 6 to 12 hours

### Warnings

**Do not use** • if you cannot swallow without chewing

**Ask a doctor before use if you have:**

- stomach pain, nausea or vomiting • noticed a sudden change in bowel habits that lasts more than 2 weeks

### When using this product

• it may cause stomach discomfort, faintness and cramps • do not chew or crush tablets(s) • do not use within 1 hour after taking antacid or milk

### Stop use and ask a doctor if

• you have rectal bleeding or fail to have a bowel movement after using this product. These could be signs of a serious condition. • you need to use a laxative for more than 1 week

**if pregnant or breast-feeding**, ask a health care professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

• take with glass of water

• adult and children 12 years of age and over – 1 to 3 tablets in a single daily dose • children 6 to under 12 years of age – 1 tablet in a single daily dose • children under 6 years of age – ask a doctor

### Other information

• Store at 15° - 30°C (59° - 86° F).

**Inactive ingredients** Lactose monohydrate, Anhydrous lactose, Pregelatinized starch, Magnesium stearate, Hypromellose, Triacetin, Talc, Methacrylic acid-ethyl acrylate copolymer (1:1) Type-A, Titanium dioxide, Triethyl citrate, D&C red no. 27, Colloidal anhydrous silica, Sodium bicarbonate, Sodium lauryl sulfate, FD&C blue no. 1, Carboxymethyl cellulose sodium, Maltodextrin, Dextrose monohydrate, Lecithin Soya. 300073

**Questions or comments?** • 1-855-642-2594 Rev.: 10/2022

\* This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., owner of the registered trademark of Dulcolax Pink®



Unvarnished Area/Coding Area  
(7 x 25 mm)

## NDC 24689-129-04

Bisacodyl Delayed Release Tablets USP

5 mg- Pink

Stimulant Laxative

250 Tablets

NDC 24689-129-04

# Bisacodyl Delayed Release Tablets USP

**5 mg-Pink**

STIMULANT LAXATIVE

Compare to Dulcolax Pink®  
active ingredient\*

**250 Tablets**



Distributed By:  
APNAR PHARMA LP  
CHINO, CA, 91710

**Caution:** Do not use if imprinted seal under cap is broken or missing.

## Drug Facts

**Active ingredient (in each tablet)**  
Bisacodyl (USP) 5 mg.....Stimulant laxative

**Uses** • for relief of occasional constipation and irregularity • this product generally produces bowel movement in 6 to 12 hours

## Warnings

**Do not use** • if you cannot swallow without chewing

**Ask a doctor before use if you have:**

- stomach pain, nausea or vomiting
- noticed a sudden change in bowel habits that lasts more than 2 weeks

**When using this product**

- it may cause stomach discomfort, faintness and cramps
- do not chew or crush tablets(s)
- do not use within 1 hour after taking antacid or milk

**Stop use and ask a doctor if**

- you have rectal bleeding or fail to have a bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

**If pregnant or breast-feeding,** ask a health care professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** • take with glass of water

- adult and children 12 years of age and over – 1 to 3 tablets in a single daily dose
- children 6 to under 12 years of age – 1 tablet in a single daily dose
- children under 6 years of age – ask a doctor

**Other information** • Store at 15° - 30°C (59° - 86° F).

**Inactive ingredients** Lactose monohydrate, Anhydrous lactose, Pregelatinized starch, Magnesium stearate, Hypromellose, Triacetin, Talc, Methacrylic acid-ethyl acrylate copolymer (1:1) Type-A, Titanium dioxide, Triethyl citrate, D&C red no. 27, Colloidal anhydrous silica, Sodium bicarbonate, Sodium lauryl sulfate, FD&C blue no. 1, Carboxymethyl cellulose sodium, Maltodextrin, Dextrose monohydrate, Lecithin Soya. 300075

**Questions or comments?** • 1-855-642-2594 Rev.: 10/2022

\* This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., owner of the registered trademark of Dulcolax Pink®



Unvarnished Area/Coding Area  
(7 x 25 mm)

## NDC 24689-129-05

Bisacodyl Delayed Release Tablets USP

5 mg- Pink

Stimulant Laxative

500 Tablets

NDC 24689-129-05

# Bisacodyl Delayed Release Tablets USP

**5 mg-Pink**

STIMULANT LAXATIVE

Compare to Dulcolax Pink®  
active ingredient\*

**500 Tablets**



Distributed By:  
APNAR PHARMA LP  
CHINO, CA, 91710

**Caution:** Do not use if imprinted seal under cap is broken or missing.

## Drug Facts

**Active ingredient (in each tablet)**  
Bisacodyl (USP) 5 mg.....Stimulant laxative

**Uses** • for relief of occasional constipation and irregularity • this product generally produces bowel movement in 6 to 12 hours

## Warnings

**Do not use** • if you cannot swallow without chewing

**Ask a doctor before use if you have:**

- stomach pain, nausea or vomiting
- noticed a sudden change in bowel habits that lasts more than 2 weeks

**When using this product**

- it may cause stomach discomfort, faintness and cramps
- do not chew or crush tablets(s)
- do not use within 1 hour after taking antacid or milk

**Stop use and ask a doctor if**

- you have rectal bleeding or fail to have a bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

**If pregnant or breast-feeding,** ask a health care professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** • take with glass of water

- adult and children 12 years of age and over – 1 to 3 tablets in a single daily dose
- children 6 to under 12 years of age – 1 tablet in a single daily dose
- children under 6 years of age – ask a doctor

**Other information** • Store at 15° - 30°C (59° - 86° F).

**Inactive ingredients** Lactose monohydrate, Anhydrous lactose, Pregelatinized starch, Magnesium stearate, Hypromellose, Triacetin, Talc, Methacrylic acid-ethyl acrylate copolymer (1:1) Type-A, Titanium dioxide, Triethyl citrate, D&C red no. 27, Colloidal anhydrous silica, Sodium bicarbonate, Sodium lauryl sulfate, FD&C blue no. 1, Carboxymethyl cellulose sodium, Maltodextrin, Dextrose monohydrate, Lecithin Soya. 300075

**Questions or comments?** • 1-855-642-2594 Rev.: 07/2022

\* This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., owner of the registered trademark of Dulcolax Pink®



Unvarnished Area/Coding Area  
(8.5 x 25 mm)

## BISACODYL

bisacodyl tablet, delayed release

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:24689-129
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

<b>BISACODYL</b> (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	5 mg
<b>Inactive Ingredients</b>		
Ingredient Name		Strength
<b>DEXTROSE MONOHYDRATE</b> (UNII: LX22YL083G)		
<b>TRIA CETIN</b> (UNII: XHX3C3X673)		
<b>D&amp;C RED NO. 27</b> (UNII: 2LRS185U6K)		
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)		
<b>METHACRYLIC ACID-ETHYL ACRYLATE COPOLYMER (1:1) TYPE A</b> (UNII: NX76LV5T8J)		
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)		
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)		
<b>TALC</b> (UNII: 7SEV7J4R1U)		
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)		
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)		
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)		
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)		
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)		
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)		
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)		
<b>CARBOXYMETHYLCELLULOSE SODIUM</b> (UNII: K679OBS311)		
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)		
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)		
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)		

Product Characteristics			
<b>Color</b>	pink	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	6mm
<b>Flavor</b>		<b>Imprint Code</b>	F
<b>Contains</b>			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:24689-129-02	20 in 1 BOTTLE; Type 0: Not a Combination Product	11/21/2022	
2	NDC:24689-129-03	100 in 1 BOTTLE; Type 0: Not a Combination Product	11/21/2022	
3	NDC:24689-129-04	250 in 1 BOTTLE; Type 0: Not a Combination Product	11/21/2022	
4	NDC:24689-129-05	500 in 1 BOTTLE; Type 0: Not a Combination Product	11/21/2022	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	11/21/2022	

**Labeler** - Apnar Pharma LP (079568229)

**Establishment**

Name	Address	ID/FEI	Business Operations
Apnar Pharma Private Limited		876730408	manufacture(24689-129) , pack(24689-129) , analysis(24689-129)

Revised: 11/2022

Apnar Pharma LP