

**UP AND UP FAMOTIDINE- famotidine tablet, film coated**  
**Target Corporation**

-----  
**Target Corporation Famotidine Tablets, 20 mg Drug Facts**

**Active ingredient (in each tablet)**

Famotidine 20 mg

**Purpose**

Acid reducer

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings**

**Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60minutes before** eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

**Other information**

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture

**Inactive ingredients**

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, titanium dioxide

**Questions?**

**Call 1-888-547-7400**

**Principal Display Panel**

see new warnings

Compare to active ingredient in Maximum Strength Pepcid® AC

maximum strength

famotidine tablets, 20 mg

acid reducer

just one tablet prevents and relieves heartburn due to acid indigestion

ACTUAL SIZE

50 + 50

100 TABLETS

2 x 50 TABLETS, 100 TOTAL

ACTUAL SIZE




see new warnings

NDC 11673-061-78

Compare to active ingredient in Maximum Strength Pepcid® AC\*

# maximum strength famotidine tablets, 20 mg acid reducer

just one tablet prevents and relieves heartburn due to acid indigestion



2 x 50 TABLETS, 100 TOTAL

MANUFACTURED BY



A8002 UM C1



246 06 0702 R000  
C-001727-01-002  
Distributed by Target Corporation  
Minneapolis, MN 55403  
TM & ©2020 Target Brands, Inc.  
**GLUTEN FREE**



### Drug Facts

**Active ingredient**      **Purpose**  
(in each tablet)  
Famotidine 20 mg.....Acid reducer

**Uses**  

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

**Warnings**  
**Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

**Do not use**  

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**Ask a doctor before use if you have**  

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or light-headedness
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

### Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if**  

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)**

**Directions**  

- Adults and children 12 years and over:
  - to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
  - to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn
  - do not use more than 2 tablets in 24 hours
  - children under 12 years: ask a doctor

**Other information**  

- read the insert to learn more about this drug before use
- keep the carton. It contains important information.
- store at 20°-25° C (68°-77° F)
- protect from moisture

**Inactive ingredients** canuba wax, croscarmellose sodium, hydroxypropyl methylcellulose, hydroxypropyl cellulose, polyethylene glycol, polyvinyl alcohol, talc, titanium dioxide

maximum strength famotidine tablets, 20 mg acid reducer

DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

Drug Facts (continued)      Tips for Managing Heartburn      Do not lie flat or bend over

Questions? Call 1-888-547-7400

Just one tablet prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Maximum Strength PeppiD® AC.

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

## UP AND UP FAMOTIDINE

famotidine tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11673-061
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>FAMOTIDINE</b> (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	8mm
<b>Flavor</b>		<b>Imprint Code</b>	L194
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC 11673-061			

1	NDC:11673-061-71	1 in 1 CARTON	05/14/2012	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:11673-061-02	25 in 1 CARTON	05/03/2012	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:11673-061-72	1 in 1 CARTON	10/30/2013	10/10/2016
3		60 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:11673-061-78	2 in 1 CARTON	04/17/2020	
4		50 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:11673-061-82	1 in 1 CARTON	11/24/2021	
5		200 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077351	05/03/2012	

**Labeler** - Target Corporation (006961700)

Revised: 11/2021

Target Corporation