

UP AND UP FAMOTIDINE- famotidine tablet
Target Corporation

Target Corporation Famotidine Tablets, 20 mg Drug Facts

Active ingredient (in each tablet)

Famotidine 20 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- if you have kidney disease, except under the advice and supervision of a doctor
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years and over:
- to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture and light

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, lactose (monohydrate), magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, and titanium dioxide

Questions?

Call 1-888-547-7400

Principal Display Panel

Compare to active ingredient in Maximum Strength Pepcid® AC

maximum strength

famotidine tablets, 20 mg

acid reducer

just one tablet prevents and relieves heartburn due to acid indigestion

ACTUAL SIZE

50 TABLETS

50 TABLETS

NDC 11673-061-71

Compare to active ingredient
in **Maximum Strength Pepcid® AC***

maximum strength famotidine tablets, 20 mg acid reducer

just one tablet prevents and relieves
heartburn due to acid indigestion



50 TABLETS



maximum
strength
**famotidine
tablets,
20 mg**
acid reducer



LOT NO.



7

245 05 0164 R01 10285492
Distributed by Target Corporation
Minneapolis, MN 55403
©2015 Target Brands, Inc.
Shop target.com

GLUTEN FREE

EXP.

:34473 UU C3

Drug Facts (continued)

Directions

- adults and children 12 years and over: to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other Information

◀ read the directions and warnings before use

Drug Facts (continued)

■ keep the carton. It contains important information.

■ store at 20°-25°C (68°-77°F)


■ protect from moisture and light

Inactive ingredients carnauba wax, colloidal silicon dioxide, croscarmellose sodium, lactose (mono hydrate), magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, and titanium dioxide

Questions? Call 1-888-547-7400

*This product is not manufactured or distributed by MERCK & Co., Inc., owner of the registered trademark Pepcid® AC.





Tips for managing heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Drug Facts

Active ingredient (in each tablet)	Purpose
Famotidine 20 mg.....	Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- if you have kidney disease, except under the advice and supervision of a doctor
- with other acid reducers

Drug Facts (continued)

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

UP AND UP FAMOTIDINE

famotidine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-061
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	

MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	WHITE	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	L194
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-061-71	1 in 1 CARTON	05/14/2012	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:11673-061-02	25 in 1 CARTON	05/03/2012	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:11673-061-72	1 in 1 CARTON	10/30/2013	
3		60 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077351	05/03/2012	

Labeler - Target Corporation (006961700)