

TOPIRAMATE- topiramate tablet
Aidarex Pharmaceuticals LLC


Topiramate Tablets

Rx only

DESCRIPTION

Topiramate is a sulfamate-substituted monosaccharide. Topiramate Tablets are available as 25 mg, 50 mg, 100 mg, and 200 mg round tablets for oral administration.

Topiramate USP is a white crystalline powder with a bitter taste. Topiramate USP is most soluble in alkaline solutions containing sodium hydroxide or sodium phosphate and having a pH of 9 to 10. It is freely soluble in acetone, chloroform, dimethylsulfoxide, and ethanol. The solubility in water is 9.8 mg/mL. Its saturated solution has a pH of 6.3. Topiramate USP has the molecular formula $C_{12}H_{21}NO_8S$ and a molecular weight of 339.37. Topiramate USP is designated chemically as 2,3:4,5-Di-O-isopropylidene- β -D-fructopyranose sulfamate and has the following structural formula:



Topiramate tablets contain the following inactive ingredients: lactose monohydrate, microcrystalline cellulose, pre-gelatinized starch, lactose monohydrate, sodium starch glycolate, magnesium stearate, opadry white (titanium dioxide, hypromellose 3cp, hypromellose 6cp, PEG 400, polysorbate 80) for 25 mg tablets, opadry yellow (titanium dioxide, hypromellose 3cp, hypromellose 6cp, PEG 400, polysorbate 80, iron oxide yellow) for 50 mg tablets, opadry yellow (hypromellose 3cp, hypromellose 6cp titanium dioxide, PEG 400, iron oxide yellow, polysorbate 80, iron oxide red) for 100 mg tablets and), opadry pink (titanium dioxide, hypromellose 6cp, PEG 400, iron oxide red) for 200 mg tablets.

CLINICAL PHARMACOLOGY

Mechanism of Action:

The precise mechanisms by which topiramate exerts its anticonvulsant effects are unknown; however, preclinical studies have revealed four properties that may contribute to topiramate's efficacy for epilepsy. Electrophysiological and biochemical evidence suggests that topiramate, at pharmacologically relevant concentrations, blocks voltage-dependent sodium channels, augments the activity of the neurotransmitter gamma-aminobutyrate at some subtypes of the GABA-A receptor, antagonizes the AMPA/kainate subtype of the glutamate receptor, and inhibits the carbonic anhydrase enzyme, particularly isozymes II and IV.

Pharmacodynamics:

Topiramate has anticonvulsant activity in rat and mouse maximal electroshock seizure (MES) tests. Topiramate is only weakly effective in blocking clonic seizures induced by the GABAA receptor antagonist, pentylenetetrazole. Topiramate is also effective in rodent models of epilepsy, which include tonic and absence-like seizures in the spontaneous epileptic rat (SER) and tonic and clonic seizures induced in rats by kindling of the amygdala or by global ischemia.

Pharmacokinetics:

The sprinkle formulation is bioequivalent to the immediate release tablet formulation and, therefore, may be substituted as a therapeutic equivalent.

Absorption of topiramate is rapid, with peak plasma concentrations occurring at approximately 2 hours following a 400 mg oral dose. The relative bioavailability of topiramate from the tablet formulation is about 80% compared to a solution. The bioavailability of topiramate is not affected by food.

The pharmacokinetics of topiramate are linear with dose proportional increases in plasma concentration over the dose range studied (200 to 800 mg/day). The mean plasma elimination half-life is 21 hours after single or multiple doses. Steady state is thus reached in about 4 days in patients with normal renal function. Topiramate is 15 to 41% bound to human plasma proteins over the blood concentration range of 0.5 to 250 mcg/mL. The fraction bound decreased as blood concentration increased.

Carbamazepine and phenytoin do not alter the binding of topiramate. Sodium valproate, at 500 mcg/mL (a concentration 5 to 10 times higher than considered therapeutic for valproate) decreased the protein binding of topiramate from 23% to 13%. Topiramate does not influence the binding of sodium valproate.

Metabolism and Excretion:

Topiramate is not extensively metabolized and is primarily eliminated unchanged in the urine (approximately 70% of an administered dose). Six metabolites have been identified in humans, none of which constitutes more than 5% of an administered dose. The metabolites are formed via hydroxylation, hydrolysis, and glucuronidation. There is evidence of renal tubular reabsorption of topiramate. In rats, given probenecid to inhibit tubular reabsorption, along with topiramate, a significant increase in renal clearance of topiramate was observed. This interaction has not been evaluated in humans. Overall, oral plasma clearance (CL/F) is approximately 20 to 30 mL/min in humans following oral administration.

Pharmacokinetic Interactions (see also Drug Interactions):

Antiepileptic Drugs

Potential interactions between topiramate and standard AEDs were assessed in controlled clinical pharmacokinetic studies in patients with epilepsy. The effect of these interactions on mean plasma AUCs are summarized under **PRECAUTIONS**

(Table 3).

Special Populations:

Renal Impairment:

The clearance of topiramate was reduced by 42% in moderately renally impaired (creatinine clearance 30 to 69 mL/min/1.73m²) and by 54% in severely renally impaired subjects (creatinine clearance <30 mL/min/1.73m²) compared to normal renal function subjects (creatinine clearance >70 mL/min/1.73m²). Since topiramate is presumed to undergo significant tubular reabsorption, it is uncertain whether this experience can be generalized to all situations of renal impairment. It is conceivable that some forms of renal disease could differentially affect glomerular filtration rate and tubular reabsorption resulting in a clearance of topiramate not predicted by creatinine clearance. In general, however, use of one-half the usual starting and maintenance dose is recommended in patients with moderate or severe renal impairment (see **PRECAUTIONS Adjustment of Dose in Renal Failure** and **DOSAGE AND ADMINISTRATION**).

Hemodialysis:

Topiramate is cleared by hemodialysis. Using a high efficiency, counterflow, single pass-dialysate hemodialysis procedure, topiramate dialysis clearance was 120 mL/min with blood flow through the dialyzer at 400 mL/min. This high clearance (compared to 20 to 30 mL/min total oral clearance in healthy adults) will remove a clinically significant amount of topiramate from the patient over the hemodialysis treatment period. Therefore, a supplemental dose may be required (see **DOSAGE AND ADMINISTRATION**).

Hepatic Impairment:

In hepatically impaired subjects, the clearance of topiramate may be decreased; the mechanism underlying the decrease is not well understood.

Age, Gender, and Race:

The pharmacokinetics of topiramate in elderly subjects (65 to 85 years of age, N=16) were evaluated in a controlled clinical study. The elderly subject population had reduced renal function [creatinine clearance (-20%)] compared to young adults. Following a single oral 100 mg dose, maximum plasma concentration for elderly and young adults was achieved at approximately 1 to 2 hours. Reflecting the primary renal elimination of topiramate, topiramate plasma and renal clearance were reduced 21% and 19%, respectively, in elderly subjects, compared to young adults. Similarly, topiramate half-life was longer (13%) in the elderly. Reduced topiramate clearance resulted in slightly higher maximum plasma concentration (23%) and AUC (25%) in elderly subjects than observed in young adults. Topiramate clearance is decreased in the elderly only to the extent that renal function is reduced. As recommended for all patients, dosage adjustment may be indicated in the elderly patient when impaired renal function (creatinine clearance rate ≤ 70 mL/min/1.73 m²) is evident. It may be useful to monitor renal function in the elderly patient (see **Special Populations:Renal Impairment, PRECAUTIONSAdjustment of Dose in Renal Failure** and **DOSAGE AND ADMINISTRATION**).

Clearance of topiramate in adults was not affected by gender or race.

Pediatric Pharmacokinetics:

Pharmacokinetics of topiramate were evaluated in patients ages 4 to 17 years receiving one or two other antiepileptic drugs. Pharmacokinetic profiles were obtained after one week at doses of 1, 3, and 9 mg/kg/day. Clearance was independent of dose.

Pediatric patients have a 50% higher clearance and consequently shorter elimination half-life than adults. Consequently, the plasma concentration for the same mg/kg dose may be lower in pediatric patients compared to adults. As in adults, hepatic enzyme inducing antiepileptic drugs decrease the steady state plasma concentrations of topiramate.

CLINICAL STUDIES

The studies described in the following sections were conducted using topiramate tablets.

Epilepsy


Monotherapy Controlled Trial

The effectiveness of topiramate as initial monotherapy in adults and children 10 years of age and older with partial onset or primary generalized seizures was established in a multicenter, randomized, double-blind, parallel-group trial.

The trial was conducted in 487 patients diagnosed with epilepsy (6 to 83 years of age) who had 1 or 2 well-documented seizures during the 3-month retrospective baseline phase who then entered the study and received topiramate 25 mg/day for 7 days in an open-label fashion. Forty-nine percent of subjects had no prior AED treatment and 17% had a diagnosis of epilepsy for greater than 24 months. Any AED therapy used for temporary or emergency purposes was discontinued prior to randomization. In the

double-blind phase, 470 patients were randomized to titrate up to 50 mg/day or 400 mg/day. If the target dose could not be achieved, patients were maintained on the maximum tolerated dose. Fifty eight percent of patients achieved the maximal dose of 400 mg/day for ≥ 2 weeks, and patients who did not tolerate 150 mg/day were discontinued. The primary efficacy assessment was a between group comparison of time to first seizure during the double-blind phase. Comparison of the Kaplan-Meier survival curves of time to first seizure favored the topiramate 400 mg/day group over the topiramate 50 mg/day group ($p=0.0002$, log rank test; Figure 1). The treatment effects with respect to time to first seizure were consistent across various patient subgroups defined by age, sex, geographic region, baseline body weight, baseline seizure type, time since diagnosis, and baseline AED use.

Figure 1: Kaplan-Meier Estimates of Cumulative Rates for Time to First Seizure



Adjunctive Therapy Controlled Trials in Adult Patients With Partial Onset Seizures

The effectiveness of topiramate as an adjunctive treatment for adults with partial onset seizures was established in six multicenter, randomized, double-blind, placebo controlled trials, two comparing several dosages of topiramate and placebo and four comparing a single dosage with placebo, in patients with a history of partial onset seizures, with or without secondarily generalized seizures.

Patients in these studies were permitted a maximum of two antiepileptic drugs (AEDs) in addition to Topiramate Tablets or placebo. In each study, patients were stabilized on optimum dosages of their concomitant AEDs during baseline phase lasting between 4 and 12 weeks. Patients who experienced a prespecified minimum number of partial onset seizures, with or without secondary generalization, during the baseline phase (12 seizures for 12-week baseline, 8 for 8-week baseline, or 3 for 4-week baseline) were randomly assigned to placebo or a specified dose of Topiramate Tablets in addition to their other AEDs.

Following randomization, patients began the double-blind phase of treatment. In five of the six studies, patients received active drug beginning at 100 mg per day; the dose was then increased by 100 mg or 200 mg/day increments weekly or every other week until the assigned dose was reached, unless intolerance prevented increases. In the sixth study (119), the 25 or 50 mg/day initial doses of topiramate were followed by respective weekly increments of 25 or 50 mg/day until the target dose of 200 mg/day was reached. After titration, patients entered a 4, 8, or 12-week stabilization period.

The numbers of patients randomized to each dose, and the actual mean and median doses in the stabilization period are shown in Table 1.

Adjunctive Therapy Controlled Trial in Pediatric Patients Ages 2 to 16 Years With Partial Onset Seizures

The effectiveness of topiramate as an adjunctive treatment for pediatric patients ages 2 to 16 years with

partial onset seizures was established in a multicenter, randomized, double-blind, placebo-controlled trial, comparing topiramate and placebo in patients with a history of partial onset seizures, with or without secondarily generalized seizures.

Patients in this study were permitted a maximum of two antiepileptic drugs (AEDs) in addition to Topiramate Tablets or placebo. In this study, patients were stabilized on optimum dosages of their concomitant AEDs during an 8 week baseline phase. Patients who experienced at least six partial onset seizures, with or without secondarily generalized seizures, during the baseline phase were randomly assigned to placebo or Topiramate Tablets in addition to their other AEDs.

Following randomization, patients began the double-blind phase of treatment. Patients received active drug beginning at 25 or 50 mg per day; the dose was then increased by 25 mg to 150 mg/day increments every other week until the assigned dosage of 125, 175, 225, or 400 mg/day based on patients' weight to approximate a dosage of 6 mg/kg per day was reached, unless intolerance prevented increases. After titration, patients entered an 8-week stabilization period.

Adjunctive Therapy Controlled Trial in Patients With Primary Generalized Tonic-Clonic Seizures

The effectiveness of topiramate as an adjunctive treatment for primary generalized tonic-clonic seizures in patients 2 years old and older was established in a multicenter randomized, double-blind, placebo-controlled trial, comparing a single dosage of topiramate and placebo.

Patients in this study were permitted a maximum of two antiepileptic drugs (AEDs) in addition to Topiramate or placebo. Patients were stabilized on optimum dosages of their concomitant AEDs during an 8-week baseline phase. Patients who experienced at least three primary generalized tonic-clonic seizures during the baseline phase were randomly assigned to placebo or Topiramate in addition to their other AEDs.

Following randomization, patients began the double-blind phase of treatment. Patients received active drug beginning at 50 mg per day for four weeks; the dose was then increased by 50 mg to 150 mg/day increments every other week until the assigned dose of 175, 225, or 400 mg/day based on patients' body weight to approximate a dosage of 6 mg/kg per day was reached, unless intolerance prevented increases. After titration, patients entered a 12-week stabilization period.

Adjunctive Therapy Controlled Trial in Patients With Lennox-Gastaut Syndrome

The effectiveness of topiramate as an adjunctive treatment for seizures associated with Lennox-Gastaut syndrome was established in a multicenter, randomized, double-blind, placebo-controlled trial comparing a single dosage of topiramate with placebo in patients 2 years of age and older.

Patients in this study were permitted a maximum of two antiepileptic drugs (AEDs) in addition to Topiramate or placebo. Patients who were experiencing at least 60 seizures per month before study entry were stabilized on optimum dosages of their concomitant AEDs during a 4-week baseline phase. Following baseline, patients were randomly assigned to placebo or Topiramate in addition to their other AEDs. Active drug was titrated beginning at 1 mg/kg per day for a week; the dose was then increased to 3 mg/kg per day for one week then to 6 mg/kg per day. After titration, patients entered an 8-week stabilization period. The primary measures of effectiveness were the percent reduction in drop attacks and a parental global rating of seizure severity.

Table 1: Topiramate Dose Summary During the Stabilization Periods of Each of Six Double-Blind, Placebo-Controlled, Add-On Trials in Adults with Partial Onset Seizures^b

Protocol	Stabilization Dose	Placebo ^a	Target Topiramate Dosage (mg/day)				
			200	400	600	800	1,000
YD	N	42	42	40	41	-	-
	Mean Dose	5.9	200	390	556	-	-
	Median Dose	6.0	200	400	600	-	-
YE	N	44	-	-	40	45	40

	Mean Dose	9.7	-	-	544	739	796
	Median Dose	10.0	-	-	600	800	1,000
Y1	N	23	-	19	-	-	-
	Mean Dose	3.8	-	395	-	-	-
	Median Dose	4.0	-	400	-	-	-
Y2	N	30	-	-	28	-	-
	Mean Dose	5.7	-	-	522	-	-
	Median Dose	6.0	-	-	600	-	-
Y3	N	28	-	-	-	25	-
	Mean Dose	7.9	-	-	-	568	-
	Median Dose	8.0	-	-	-	600	-
119	N	90	157	-	-	-	-
	Mean Dose	8	200	-	-	-	-
	Median Dose	8	200	-	-	-	-

^a Placebo dosages are given as the number of tablets. Placebo target dosages were as follows: Protocol Y1, 4 tablets/day; Protocols YD and Y2, 6 tablets/day; Protocol Y3 and 119, 8 tablets/day; Protocol YE, 10 tablets/day.

^b Dose-response studies were not conducted for other indications or pediatric partial onset seizures.

In all add-on trials, the reduction in seizure rate from baseline during the entire double-blind phase was measured. The median percent reductions in seizure rates and the responder rates (fraction of patients with at least a 50% reduction) by treatment group for each study are shown below in Table 2. As described above, a global improvement in seizure severity was also assessed in the Lennox-Gastaut trial.

Table 2: Efficacy Results in Double-Blind, Placebo-Controlled, Add-On Trials

		Target Topiramate Dosage (mg/day)						
Protocol	Efficacy Results	Placebo	200	400	600	800	1,000	≈6mg/kg/day*
Partial Onset Seizures Studies in Adults								
YD	N	45	45	45	46	-	-	-
	Median % Reduction	11.6	27.2 ^a	47.5 ^b	44.7 ^c	-	-	-
	% Responders	18	24	44 ^d	46 ^d	-	-	-
YE	N	47	-	-	48	48	47	-
	Median % Reduction	1.7	-	-	40.8 ^c	41.0 ^c	36.0 ^c	-
	% Responders	9	-	-	40 ^c	41 ^c	36 ^d	-
Y1	N	24	-	23	-	-	-	-
	Median % Reduction	1.1	-	40.7 ^e	-	-	-	-
	% Responders	8	-	35 ^d	-	-	-	-
Y2	N	30	-	-	30	-	-	-
	Median % Reduction	-12.2	-	-	46.4 ^f	-	-	-
	% Responders	10	-	-	47 ^c	-	-	-
Y3	N	28	-	-	-	28	-	-
	Median % Reduction	-20.6	-	-	-	24.3 ^c	-	-
	% Responders	0	-	-	-	43 ^c	-	-
119	N	91	168	-	-	-	-	-
	Median % Reduction	20.0	44.2 ^c	-	-	-	-	-
	% Responders	24	45 ^c	-	-	-	-	-
Studies in Pediatric Patients								
YP	N	45	-	-	-	-	-	41
	Median % Reduction	10.5	-	-	-	-	-	33.1 ^d

% Responders	20	-	-	-	-	-	39
Primary Generalized Tonic-Clonic							
YTC N	40	-	-	-	-	-	39
Median % Reduction	9.0	-	-	-	-	-	56.7 ^d
% Responders	20	-	-	-	-	-	56 ^c

Comparisons with placebo: ^a p=0.080; ^b p≤0.010; ^c p≤0.001; ^d p≤0.050; ^e p=0.065; ^f p≤0.005; ^g p=0.071;

^h Median % reduction and % responders are reported for PGTC Seizures;

ⁱ Median % reduction and % responders for drop attacks, i.e., tonic or atonic seizures;

^j Percent of subjects who were minimally, much, or very much improved from baseline

* For Protocols YP and YTC, protocol-specified target dosages (<9.3 mg/kg/day) were assigned based on subject's weight to approximate a dosage of 6 mg/kg per day; these dosages corresponded to mg/day dosages of 125, 175, 225, and 400 mg/day.

Subset analyses of the antiepileptic efficacy of Topiramate Tablets in these studies showed no differences as a function of gender, race, age, baseline seizure rate, or concomitant AED.

INDICATIONS AND USAGE

Monotherapy Epilepsy

Topiramate Tablets are indicated as initial monotherapy in patients 10 years of age and older with partial onset or primary generalized tonic-clonic seizures.

Effectiveness was demonstrated in a controlled trial in patients with epilepsy who had no more than 2 seizures in the 3 months prior to enrollment. Safety and effectiveness in patients who were converted to monotherapy from a previous regimen of other anticonvulsant drugs have not been established in controlled trials.

Adjunctive Therapy Epilepsy

Topiramate Tablets are indicated as adjunctive therapy for adults and pediatric patients ages 2 to 16 years with partial onset seizures, or primary generalized tonic-clonic seizures, and in patients 2 years of age and older with seizures associated with Lennox-Gastaut syndrome.

CONTRAINDICATIONS

Topiramate is contraindicated in patients with a history of hypersensitivity to any component of this product.

WARNINGS

Metabolic Acidosis

Hyperchloremic, non-anion gap, metabolic acidosis (i.e., decreased serum bicarbonate below the normal reference range in the absence of chronic respiratory alkalosis) is associated with topiramate treatment. This metabolic acidosis is caused by renal bicarbonate loss due to the inhibitory effect of topiramate on carbonic anhydrase. Such electrolyte imbalance has been observed with the use of topiramate in placebo-controlled clinical trials and in the post-marketing period. Generally, topiramate-induced metabolic acidosis occurs early in treatment although cases can occur at any time during treatment. Bicarbonate decrements are usually mild/moderate (average decrease of 4 mEq/L at daily doses of 400 mg in adults and at approximately 6 mg/kg/day in pediatric patients); rarely, patients can experience severe decrements to values below 10 mEq/L. Conditions or therapies that predispose to acidosis (such as renal disease, severe respiratory disorders, status epilepticus, diarrhea, surgery,

ketogenic diet, or drugs) may be additive to the bicarbonate lowering effects of topiramate.

In adults, the incidence of persistent treatment-emergent decreases in serum bicarbonate (levels of <20 mEq/L at two consecutive visits or at the final visit) in controlled clinical trials for adjunctive treatment of epilepsy was 32% for 400 mg/day, and 1% for placebo. Metabolic acidosis has been observed at doses as low as 50 mg/day. The incidence of persistent treatment-emergent decreases in serum bicarbonate in adults in the epilepsy controlled clinical trial for monotherapy was 15% for 50 mg/day and 25% for 400 mg/day. The incidence of a markedly abnormally low serum bicarbonate (i.e., absolute value <17 mEq/L and >5 mEq/L decrease from pretreatment) in the adjunctive therapy trials was 3% for 400 mg/day, and 0% for placebo and in the monotherapy trial was 1% for 50 mg/day and 7% for 400 mg/day. Serum bicarbonate levels have not been systematically evaluated at daily doses greater than 400 mg/day.

In pediatric patients (<16 years of age), the incidence of persistent treatment emergent decreases in serum bicarbonate in placebo-controlled trials for adjunctive treatment of Lennox-Gastaut syndrome or refractory partial onset seizures was 67% for topiramate (at approximately 6 mg/kg/day), and 10% for placebo. The incidence of a markedly abnormally low serum bicarbonate (i.e., absolute value <17 mEq/L and >5 mEq/L decrease from pretreatment) in these trials was 11% for topiramate and 0% for placebo. Cases of moderately severe metabolic acidosis have been reported in patients as young as 5 months old, especially at daily doses above 5 mg/kg/day.

In pediatric patients (10 years up to 16 years of age), the incidence of persistent treatment-emergent decreases in serum bicarbonate in the epilepsy controlled clinical trial for monotherapy was 7% for 50 mg/day and 20% for 400 mg/day. The incidence of a markedly abnormally low serum bicarbonate (i.e., absolute value <17 mEq/L and >5 mEq/L decrease from pretreatment) in this trial was 4% for 50 mg/day and 4% for 400 mg/day.

The incidence of a markedly abnormally low serum bicarbonate (i.e., absolute value <17 mEq/L and >5 mEq/L decrease from pretreatment) in these trials was 11% for 200 mg/day, 9% for 100 mg/day, 2% for 50 mg/day, and <1% for placebo.

Some manifestations of acute or chronic metabolic acidosis may include hyperventilation, nonspecific symptoms such as fatigue and anorexia, or more severe sequelae including cardiac arrhythmias or stupor. Chronic, untreated metabolic acidosis may increase the risk for nephrolithiasis or nephrocalcinosis, and may also result in osteomalacia (referred to as rickets in pediatric patients) and/or osteoporosis with an increased risk for fractures. Chronic metabolic acidosis in pediatric patients may also reduce growth rates. A reduction in growth rate may eventually decrease the maximal height achieved. The effect of topiramate on growth and bone-related sequelae has not been systematically investigated.

Measurement of baseline and periodic serum bicarbonate during topiramate treatment is recommended. If metabolic acidosis develops and persists, consideration should be given to reducing the dose or discontinuing topiramate (using dose tapering). If the decision is made to continue patients on topiramate in the face of persistent acidosis, alkali treatment should be considered.

Acute Myopia and Secondary Angle Closure Glaucoma

A syndrome consisting of acute myopia associated with secondary angle closure glaucoma has been reported in patients receiving Topiramate. Symptoms include acute onset of decreased visual acuity and/or ocular pain. Ophthalmologic findings can include myopia, anterior chamber shallowing, ocular hyperemia (redness) and increased intraocular pressure. Mydriasis may or may not be present. This syndrome may be associated with supraciliary effusion resulting in anterior displacement of the lens and iris, with secondary angle closure glaucoma. Symptoms typically occur within 1 month of initiating Topiramate therapy. In contrast to primary narrow angle glaucoma, which is rare under 40 years of age, secondary angle closure glaucoma associated with topiramate has been reported in pediatric patients as well as adults. The primary treatment to reverse symptoms is discontinuation of Topiramate as rapidly as possible, according to the judgment of the treating physician. Other measures, in conjunction with

discontinuation of Topiramate, may be helpful.

Elevated intraocular pressure of any etiology, if left untreated, can lead to serious sequelae including permanent vision loss.

Oligohidrosis and Hyperthermia

Oligohidrosis (decreased sweating), infrequently resulting in hospitalization, has been reported in association with Topiramate use. Decreased sweating and an elevation in body temperature above normal characterized these cases. Some of the cases were reported after exposure to elevated environmental temperatures.

The majority of the reports have been in children. Patients, especially pediatric patients, treated with Topiramate should be monitored closely for evidence of decreased sweating and increased body temperature, especially in hot weather.

Caution should be used when Topiramate is prescribed with other drugs that predispose patients to heat-related disorders; these drugs include, but are not limited to, other carbonic anhydrase inhibitors and drugs with anticholinergic activity.

Withdrawal of AEDs

Antiepileptic drugs, including Topiramate should be withdrawn gradually to minimize the potential of increased seizure frequency.

Cognitive/Neuropsychiatric Adverse Events

Adults

Adverse events most often associated with the use of Topiramate were related to the central nervous system and were observed in the epilepsy populations. In adults, the most frequent of these can be classified into three general categories: 1) Cognitive-related dysfunction (e.g. confusion, psychomotor slowing, difficulty with concentration/attention, difficulty with memory, speech or language problems, particularly word-finding difficulties); 2) Psychiatric/behavioral disturbances (e.g. depression or mood problems); and 3) Somnolence or fatigue.

Cognitive-Related Dysfunction

The majority of cognitive-related adverse events were mild to moderate in severity, and they frequently occurred in isolation. Rapid titration rate and higher initial dose were associated with higher incidences of these events. Many of these events contributed to withdrawal from treatment [see **ADVERSE REACTIONS, Table 4 and Table 6**].

In the original add-on epilepsy controlled trials (using rapid titration such as 100 to 200 mg/day weekly increments), the proportion of patients who experienced one or more cognitive-related adverse events was 42% for 200 mg/day, 41% for 400 mg/day, 52% for 600 mg/day, 56% for 800 and 1000 mg/day, and 14% for placebo. These dose-related adverse reactions began with a similar frequency in the titration or in the maintenance phase, although in some patients the events began during titration and persisted into the maintenance phase. Some patients who experienced one or more cognitive-related adverse events in the titration phase had a dose-related recurrence of these events in the maintenance phase.

In the monotherapy epilepsy controlled trial, the proportion of patients who experienced one or more cognitive-related adverse events was 19% for Topiramate 50 mg/day and 26% for 400 mg/day.

Psychiatric/Behavioral Disturbances

Psychiatric/behavioral disturbances (depression or mood problems) were dose-related for the epilepsy populations.

Somnolence/Fatigue

Somnolence and fatigue were the adverse events most frequently reported during clinical trials of

Topiramate for adjunctive epilepsy. For the adjunctive epilepsy population, the incidence of somnolence did not differ substantially between 200 mg/day and 1000 mg/day, but the incidence of fatigue was dose-related and increased at dosages above 400 mg/day. For the monotherapy epilepsy population in the 50 mg/day and 400 mg/day groups, the incidence of somnolence was dose-related (9% for the 50 mg/day group and 15% for the 400 mg/day group) and the incidence of fatigue was comparable in both treatment groups (14% each).

Additional nonspecific CNS events commonly observed with topiramate in the add-on epilepsy population include dizziness or ataxia.

Pediatric Patients

In double-blind adjunctive therapy and monotherapy epilepsy clinical studies, the incidences of cognitive/neuropsychiatric adverse events in pediatric patients were generally lower than observed in adults. These events included psychomotor slowing, difficulty with concentration/attention, speech disorders/related speech problems and language problems. The most frequently reported neuropsychiatric events in pediatric patients during adjunctive therapy double-blind studies were somnolence and fatigue. The most frequently reported neuropsychiatric events in pediatric patients in the 50 mg/day and 400 mg/day groups during the monotherapy double-blind study were headache, dizziness anorexia, and somnolence.

No patients discontinued treatment due to any adverse events in the adjunctive epilepsy double-blind trials. In the monotherapy epilepsy double-blind trial, 1 pediatric patient (2%) in the 50 mg/day group and 7 pediatric patients (12%) in the 400 mg/day group discontinued treatment due to any adverse events. The most common adverse event associated with discontinuation of therapy was difficulty with concentration/attention; all occurred in the 400 mg/day group.

Sudden Unexplained Death in Epilepsy (SUDEP)

During the course of premarketing development of Topiramate Tablets, 10 sudden and unexplained deaths were recorded among a cohort of treated patients (2,796 subject years of exposure). This represents an incidence of 0.0035 deaths per patient year. Although this rate exceeds that expected in a healthy population matched for age and sex, it is within the range of estimates for the incidence of sudden unexplained deaths in patients with epilepsy not receiving Topiramate (ranging from 0.0005 for the general population of patients with epilepsy, to 0.003 for a clinical trial population similar to that in the Topiramate program, to 0.005 for patients with refractory epilepsy).

PRECAUTIONS

Hyperammonemia and Encephalopathy Associated with Concomitant Valproic Acid Use

Concomitant administration of topiramate and valproic acid has been associated with hyperammonemia with or without encephalopathy in patients who have tolerated either drug alone. Clinical symptoms of hyperammonemic encephalopathy often include acute alterations in level of consciousness and/or cognitive function with lethargy or vomiting. In most cases, symptoms and signs abated with discontinuation of either drug. This adverse event is not due to a pharmacokinetic interaction.

It is not known if topiramate monotherapy is associated with hyperammonemia.

Patients with inborn errors of metabolism or reduced hepatic mitochondrial activity may be at an increased risk for hyperammonemia with or without encephalopathy. Although not studied, an interaction of topiramate and valproic acid may exacerbate existing defects or unmask deficiencies in susceptible persons.

In patients who develop unexplained lethargy, vomiting, or changes in mental status, hyperammonemic encephalopathy should be considered and an ammonia level should be measured.

Kidney Stones

A total of 32/2,086 (1.5%) of adults exposed to topiramate during its adjunctive epilepsy therapy development reported the occurrence of kidney stones, an incidence about 2 to 4 times greater than expected in a similar, untreated population. In the double-blind monotherapy epilepsy study, a total of 4/319 (1.3%) of adults exposed to topiramate reported the occurrence of kidney stones. As in the general population, the incidence of stone formation among topiramate treated patients was higher in men. Kidney stones have also been reported in pediatric patients.

An explanation for the association of Topiramate and kidney stones may lie in the fact that topiramate is a carbonic anhydrase inhibitor. Carbonic anhydrase inhibitors, e.g., acetazolamide or dichlorphenamide, promote stone formation by reducing urinary citrate excretion and by increasing urinary pH. The concomitant use of Topiramate with other carbonic anhydrase inhibitors or potentially in patients on a ketogenic diet may create a physiological environment that increases the risk of kidney stone formation, and should therefore be avoided.

Increased fluid intake increases the urinary output, lowering the concentration of substances involved in stone formation. Hydration is recommended to reduce new stone formation.

Paresthesia

Paresthesia (usually tingling of the extremities), an effect associated with the use of other carbonic anhydrase inhibitors, appears to be a common effect of Topiramate. Paresthesia was more frequently reported in the monotherapy epilepsy trials versus the adjunctive therapy epilepsy trials. In the majority of instances, paresthesia did not lead to treatment discontinuation.

Adjustment of Dose in Renal Failure

The major route of elimination of unchanged topiramate and its metabolites is via the kidney. Dosage adjustment may be required in patients with reduced renal function

(see **DOSAGE AND ADMINISTRATION**).

Decreased Hepatic Function

In hepatically impaired patients, topiramate should be administered with caution as the clearance of topiramate may be decreased.

Information for Patients

Patients taking Topiramate should be told to seek immediate medical attention if they experience blurred vision or periorbital pain.

Patients, especially pediatric patients, treated with Topiramate should be monitored closely for evidence of decreased sweating and increased body temperature, especially in hot weather.

Patients, particularly those with predisposing factors, should be instructed to maintain an adequate fluid intake in order to minimize the risk of renal stone formation [see **PRECAUTIONS:Kidney Stones**, for support regarding hydration as a preventative measure].

Patients should be warned about the potential for somnolence, dizziness, confusion, and difficulty concentrating and advised not to drive or operate machinery until they have gained sufficient experience on topiramate to gauge whether it adversely affects their mental and/or motor performance.

Additional food intake may be considered if the patient is losing weight while on this medication.

Laboratory Tests:

Measurement of baseline and periodic serum bicarbonate during topiramate treatment is recommended (see **WARNINGS**).

Drug Interactions:

In vitro studies indicate that topiramate does not inhibit enzyme activity for CYP1A2, CYP2A6,

CYP2B6, CYP2C9, CYP2C19, CYP2D6, CYP2E1 and CYP3A4/5 isozymes.

Antiepileptic Drugs

Potential interactions between topiramate and standard AEDs were assessed in controlled clinical pharmacokinetic studies in patients with epilepsy. The effects of these interactions on mean plasma AUCs are summarized in Table 3.

In Table 3, the second column (AED concentration) describes what happens to the concentration of the AED listed in the first column when topiramate is added.

The third column (topiramate concentration) describes how the coadministration of a drug listed in the first column modifies the concentration of topiramate in experimental settings when Topiramate was given alone.

Table 3: Summary of AED Interactions with Topiramate

AED Co-administered	AED Concentration	Topiramate Concentration
Phenytoin	NC or 25% increase ^a	48% decrease
Carbamazepine (CBZ)	NC	40% decrease
CBZ epoxide ^b	NC	NE
Valproic acid	11% decrease	4% decrease
Phenobarbital	NC	NE
Primidone	NC	NE
Lamotrigine	NC at TPM doses up to 400 mg/day	15% increase

^a = Plasma concentration increased 25% in some patients, generally those on a b.i.d. dosing regimen of phenytoin.

^b = Is not administered but is an active metabolite of carbamazepine.

NC = Less than 10% change in plasma concentration.

AED = Antiepileptic drug.

NE = Not Evaluated.

TPM - Topiramate

In addition to the pharmacokinetic interaction described in the above table, concomitant administration of valproic acid and topiramate has been associated with hyperammonemia with and without encephalopathy (see **PRECAUTIONS**, Hyperammonemia and Encephalopathy Associated with Concomitant Valproic Acid Use).

Other Drug Interactions

Digoxin: In a single-dose study, serum digoxin AUC was decreased by 12% with concomitant Topiramate administration. The clinical relevance of this observation has not been established.

CNS Depressants: Concomitant administration of Topiramate and alcohol or other CNS depressant drugs has not been evaluated in clinical studies. Because of the potential of topiramate to cause CNS depression, as well as other cognitive and/or neuropsychiatric adverse events, topiramate should be used with extreme caution if used in combination with alcohol and other CNS depressants.

Oral Contraceptives: In a pharmacokinetic interaction study in healthy volunteers with a concomitantly administered combination oral contraceptive product containing 1 mg norethindrone (NET) plus 35 mcg ethinyl estradiol (EE), Topiramate given in the absence of other medications at doses of 50 to 200 mg/day was not associated with statistically significant changes in mean exposure (AUC) to either component of the oral contraceptive. In another study, exposure to EE was statistically significantly decreased at doses of 200, 400, and 800 mg/day (18%, 21%, and 30%, respectively) when given as

adjunctive therapy in patients taking valproic acid. In both studies, Topiramate (50 mg/day to 800 mg/day) did not significantly affect exposure to NET. Although there was a dose dependent decrease in EE exposure for doses between 200 to 800 mg/day, there was no significant dose dependent change in EE exposure for doses of 50 to 200 mg/day. The clinical significance of the changes observed is not known. The possibility of decreased contraceptive efficacy and increased breakthrough bleeding should be considered in patients taking combination oral contraceptive products with Topiramate. Patients taking estrogen containing contraceptives should be asked to report any change in their bleeding patterns. Contraceptive efficacy can be decreased even in the absence of breakthrough bleeding.

Hydrochlorothiazide (HCTZ): A drug-drug interaction study conducted in healthy volunteers evaluated the steady-state pharmacokinetics of HCTZ (25 mg q24h) and topiramate (96 mg q12h) when administered alone and concomitantly. The results of this study indicate that topiramate C_{max} increased by 27% and AUC increased by 29% when HCTZ was added to topiramate. The clinical significance of this change is unknown. The addition of HCTZ to topiramate therapy may require an adjustment of the topiramate dose. The steady-state pharmacokinetics of HCTZ were not significantly influenced by the concomitant administration of topiramate. Clinical laboratory results indicated decreases in serum potassium after topiramate or HCTZ administration, which were greater when HCTZ and topiramate were administered in combination.

Pioglitazone: A drug-drug interaction study conducted in healthy volunteers evaluated the steady-state pharmacokinetics of topiramate and pioglitazone when administered alone and concomitantly. A 15% decrease in the $AUC_{T,SS}$ of pioglitazone

with no alteration in $C_{max,SS}$ was observed. This finding was not statistically significant. In addition, a 13% and 16% decrease in $C_{max,SS}$ and $AUC_{T,SS}$ respectively, of the active hydroxy-metabolite was noted as well as a 60% decrease in $C_{max,SS}$ and $AUC_{T,SS}$ of the active keto-metabolite. The clinical significance of these findings is not known. When Topiramate is added to pioglitazone therapy or pioglitazone is added to Topiramate therapy, careful attention should be given to the routine monitoring of patients for adequate control of their diabetic disease state.

Lithium: Multiple dosing of topiramate 100 mg every 12 hrs decreased the AUC and C_{max} of Lithium (300 mg every 8 hrs) by 20% (N=12, 6 M; 6 F).

Haloperidol: The pharmacokinetics of a single dose of haloperidol (5 mg) were not affected following multiple dosing of topiramate (100 mg every 12 hr) in 13 healthy adults (6 M, 7 F).

Amitriptyline: There was a 12% increase in AUC and C_{max} for amitriptyline (25 mg per day) in 18 normal subjects (9 male; 9 female) receiving 200 mg/day of topiramate. Some subjects may experience a large increase in amitriptyline concentration in the presence of topiramate and any adjustments in amitriptyline dose should be made according to the patient's clinical response and not on the basis of plasma levels.

Sumatriptan: Multiple dosing of topiramate (100 mg every 12 hrs) in 24 healthy volunteers (14 M, 10 F) did not affect the pharmacokinetics of single dose sumatriptan either orally (100 mg) or subcutaneously (6 mg).

Risperidone: There was a 25% decrease in exposure to risperidone (2 mg single dose) in 12 healthy volunteers (6 M, 6 F) receiving 200 mg/day of topiramate. Therefore, patients receiving risperidone in combination with topiramate should be closely monitored for clinical response.

Propranolol: Multiple dosing of topiramate (200 mg/day) in 34 healthy volunteers (17 M, 17 F) did not affect the pharmacokinetics of propranolol following daily 160 mg doses. Propranolol doses of 160 mg/day in 39 volunteers (27M, 12F) had no effect on the exposure to topiramate at a dose of 200 mg/day of topiramate.

Dihydroergotamine: Multiple dosing of topiramate (200 mg/day) in 24 healthy volunteers (12 M, 12 F) did not affect the pharmacokinetics of a 1 mg subcutaneous dose of dihydroergotamine. Similarly, a 1

mg subcutaneous dose of

dihydroergotamine did not affect the pharmacokinetics of a 200 mg/day dose of topiramate in the same study.

Others: Concomitant use of Topiramate, a carbonic anhydrase inhibitor, with other carbonic anhydrase inhibitors, e.g., acetazolamide or dichlorphenamide, may create a physiological environment that increases the risk of renal stone formation, and should therefore be avoided.

Drug/Laboratory Test Interactions There are no known interactions of topiramate with commonly used laboratory tests.

Carcinogenesis, Mutagenesis, Impairment of Fertility:

An increase in urinary bladder tumors was observed in mice given topiramate (20, 75, and 300 mg/kg) in the diet for 21 months. The elevated bladder tumor incidence, which was statistically significant in males and females receiving 300 mg/kg, was primarily due to the increased occurrence of a smooth muscle tumor considered histomorphologically unique to mice. Plasma exposures in mice receiving 300 mg/kg were approximately 0.5 to 1 times steady-state exposures measured in patients receiving topiramate monotherapy at the recommended human dose (RHD) of 400 mg, and 1.5 to 2 times steady-state topiramate exposures in patients receiving 400 mg of topiramate plus phenytoin. The relevance of this finding to human carcinogenic risk is uncertain. No evidence of carcinogenicity was seen in rats following oral administration of topiramate for 2 years at doses up to 120 mg/kg (approximately 3 times the RHD on a mg/m² basis).

Topiramate did not demonstrate genotoxic potential when tested in a battery of *in vitro* and *in vivo* assays. Topiramate was not mutagenic in the Ames test or the *in vitro* mouse lymphoma assay; it did not increase unscheduled DNA synthesis in rat hepatocytes *in vitro*; and it did not increase chromosomal aberrations in human lymphocytes *in vitro* or in rat bone marrow *in vivo*.

No adverse effects on male or female fertility were observed in rats at doses up to 100 mg/kg (2.5 times the RHD on a mg/m² basis).

Pregnancy: Pregnancy Category C.

Topiramate has demonstrated selective developmental toxicity, including teratogenicity, in experimental animal studies. When oral doses of 20, 100, or 500 mg/kg were administered to pregnant mice during the period of organogenesis, the incidence of fetal malformations (primarily craniofacial defects) was increased at all doses. The low dose is approximately 0.2 times the recommended human dose (RHD=400 mg/day) on a mg/m² basis. Fetal body weights and skeletal ossification were reduced at 500 mg/kg in conjunction with decreased maternal body weight gain.

In rat studies (oral doses of 20, 100, and 500 mg/kg or 0.2, 2.5, 30 and 400 mg/kg), the frequency of limb malformations (ectrodactyly, micromelia, and amelia) was increased among the offspring of dams treated with 400 mg/kg (10 times the RHD on a mg/m² basis) or greater during the organogenesis period of pregnancy. Embryotoxicity (reduced fetal body weights, increased incidence of structural variations) was observed at doses as low as 20 mg/kg (0.5 times the RHD on a mg/m² basis). Clinical signs of maternal toxicity were seen at 400 mg/kg and above, and maternal body weight gain was reduced during treatment with 100 mg/kg or greater.

In rabbit studies (20, 60, and 180 mg/kg or 10, 35, and 120 mg/kg orally during organogenesis), embryo/fetal mortality was increased at 35 mg/kg (2 times the RHD on a mg/m² basis) or greater, and teratogenic effects (primarily rib and vertebral malformations) were observed at 120 mg/kg (6 times the RHD on a mg/m² basis).

Evidence of maternal toxicity (decreased body weight gain, clinical signs, and/or mortality) was seen at 35 mg/kg and above.

When female rats were treated during the latter part of gestation and throughout lactation (0.2, 4, 20, and 100 mg/kg or 2, 20, and 200 mg/kg), offspring exhibited decreased viability and delayed physical

development at 200 mg/kg (5 times the RHD on a mg/m² basis) and reductions in pre- and/or postweaning body weight gain at 2 mg/kg (0.05 times the RHD on a mg/m² basis) and above. Maternal toxicity (decreased body weight gain, clinical signs) was evident at 100 mg/kg or greater.

In a rat embryo/fetal development study with a postnatal component (0.2, 2.5, 30 or 400 mg/kg during organogenesis; noted above), pups exhibited delayed physical development at 400 mg/kg (10 times the RHD on a mg/m² basis) and persistent reductions in body weight gain at 30 mg/kg (1 times the RHD on a mg/m² basis) and higher.

There are no studies using Topiramate in pregnant women. Topiramate should be used during pregnancy only if the potential benefit outweighs the potential risk to the fetus.

In post-marketing experience, cases of hypospadias have been reported in male infants exposed in utero to topiramate, with or without other anticonvulsants; however, a causal relationship with topiramate has not been established.

Labor and Delivery:

In studies of rats where dams were allowed to deliver pups naturally, no drug-related effects on gestation length or parturition were observed at dosage levels up to 200 mg/kg/day.

The effect of Topiramate on labor and delivery in humans is unknown.

Nursing Mothers:

Topiramate is excreted in the milk of lactating rats. The excretion of topiramate in human milk has not been evaluated in controlled studies. Limited observations in patients suggest an extensive secretion of topiramate into breast milk. Since many drugs are excreted in human milk, and because the potential for serious adverse reactions in nursing infants to Topiramate is unknown, the potential benefit to the mother should be weighed against the potential risk to the infant when considering recommendations regarding nursing.

Pediatric Use:

Safety and effectiveness in patients below the age of 2 years have not been established for the adjunctive therapy treatment of partial onset seizures, primary generalized tonic-clonic seizures, or seizures associated with Lennox-Gastaut syndrome. Safety and effectiveness in patients below the age of 10 years have not been established for the monotherapy treatment of epilepsy. Topiramate is associated with metabolic acidosis. Chronic untreated metabolic acidosis in pediatric patients may cause osteomalacia/rickets and may reduce growth rates. A reduction in growth rate may eventually decrease the maximal height achieved. The effect of topiramate on growth and bone-related sequelae has not been systematically investigated (see **WARNINGS**).

Geriatric Use:

In clinical trials, 3% of patients were over 60. No age related difference in effectiveness or adverse effects were evident. However, clinical studies of topiramate did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently than younger subjects. Dosage adjustment may be necessary for elderly with impaired renal function (creatinine clearance rate ≤ 70 mL/min/1.73 m²) due to reduced clearance of topiramate (see **CLINICAL PHARMACOLOGY** and **DOSAGE AND ADMINISTRATION**).

Race and Gender Effects:

Evaluation of effectiveness and safety in clinical trials has shown no race or gender related effects.

ADVERSE REACTIONS

The data described in the following section were obtained using Topiramate Tablets.

Monotherapy Epilepsy

The adverse events in the controlled trial that occurred most commonly in adults in the 400 mg/day group and at a rate higher than the 50 mg/day group were: paresthesia, weight decrease, somnolence, anorexia, dizziness, and difficulty with

memory NOS [see Table 4].

The adverse events in the controlled trial that occurred most commonly in children (10 years up to 16 years of age) in the 400 mg/day group and at a rate higher than the 50 mg/day group were: weight decrease, upper respiratory tract infection, paresthesia, anorexia, diarrhea, and mood problems [see Table 5].

Approximately 21% of the 159 adult patients in the 400 mg/day group who received topiramate as monotherapy in the controlled clinical trial discontinued therapy due to adverse events. Adverse events associated with discontinuing therapy ($\geq 2\%$) included depression, insomnia, difficulty with memory (NOS), somnolence, paresthesia, psychomotor slowing, dizziness, and nausea.

Approximately 12% of the 57 pediatric patients in the 400 mg/day group who received topiramate as monotherapy in the controlled clinical trial discontinued therapy due to adverse events. Adverse events associated with discontinuing therapy ($\geq 5\%$) included difficulty with concentration/attention.

The prescriber should be aware that these data cannot be used to predict the frequency of adverse events in the course of usual medical practice where patient characteristics and other factors may differ from those prevailing during the clinical study. Similarly, the cited frequencies cannot be directly compared with data obtained from other clinical investigations involving different treatments, uses, or investigators. Inspection of these frequencies, however, does provide the prescribing physician with a basis to estimate the relative contribution of drug and non-drug factors to the adverse event incidences in the population studied.

Table 4: Incidence of Treatment-Emergent Adverse Events in the Monotherapy Epilepsy Trial in Adults^a Where Rate Was at Least 2% in the 400 mg/day Topiramate Group and Greater Than the Rate in the 50 mg/day Topiramate Group

Body System/ Adverse Event	Topiramate Dosage (mg/day)	
	50 (N=160)	400 (N=159)
Body as a Whole-		
General Disorders		
Asthenia	4	6
Leg Pain	2	3
Chest Pain	1	2
Central & Peripheral Nervous system Disorders		
Paresthesia		
Dizziness	21	40
Hypoaesthesia	13	14
Ataxia	4	5
Hypertonia	3	4
	0	3
Gastro-Intestinal System Disorders		
Diarrhea	5	6
Constipation	1	4
Gastritis	0	3
Dry Mouth	1	3
Gastroesophageal Reflux	1	2
Liver and Biliary System Disorders		
Gamma-GT Increased	1	3

Metabolic and nutritional Disorders		
Weight Decrease	6	16
Psychiatric Disorders		
Somnolence	9	15
Anorexia	4	14
Difficulty with Memory NOS	5	10
Insomnia	8	9
Depression	7	9
Difficulty with Concentration/Attention	7	8
Anxiety		
Psychomotor Slowing	4	6
Mood Problems	3	5
Confusion	2	5
Cognitive Problem NOS	3	4
Libido Decreased	1	4
	0	3
Reproductive Disorders, female		
Vaginal Hemorrhage	0	3
Red Blood Cell disorders		
Anemia	1	2

^a Values represent the percentage of patients reporting a given adverse event. Patients may have reported more than one adverse event during the study and can be included in more than one adverse event category

Table 5: Incidence of Treatment-Emergent Adverse Events in the Monotherapy Epilepsy Trail in Children Ages 10 up to 16 Yearsa where Rate was at least 5% in the 400 mg/day Topiramate Group and Greater Than the Rate in the 50 mg/day Topiramate Group

Body System/ Adverse Event	Topiramate Dosage (mg/day) ^b	
	50 (N=57)	400 (N=57)
Body as a Whole-General Disorders		
Fever	0	9
Central & Peripheral Nervous system Disorders		
Paresthesia	2	16
Gastro-Intestinal System Disorders		
Diarrhea	5	11
Metabolic and nutritional Disorders		
Weight Decrease	7	21
Psychiatric Disorders		
Anorexia	11	14
Mood Problems	2	11
Difficulty with Concentration/Attention	4	9
Cognitive Problem NOS		
Nervousness	0	7
	4	5
Resistance Mechanism Disorders		
Infection Viral	4	9
Infection	2	7
Respiratory System Disorders	16	

Upper Respiratory Tract Infection	2	18
Rhinitis	2	7
Bronchitis	2	7
Sinusitis		5
Skin and Appendages Disorders		
Alopecia	2	5

^a Values represent the percentage of patients reporting a given adverse event. Patients may have reported more than one adverse event during the study and can be included in more than one adverse event category

Adjunctive Therapy Epilepsy

The most commonly observed adverse events associated with the use of topiramate at dosages of 200 to 400 mg/day in controlled trials in adults with partial onset seizures, primary generalized tonic-clonic seizures, or Lennox-Gastaut syndrome, that were seen at greater frequency in topiramate-treated patients and did not appear to be doserelated were: somnolence, dizziness, ataxia, speech disorders and related speech problems, psychomotor slowing, abnormal vision, difficulty with memory, paresthesia and diplopia [see Table 6]. The most common dose-related adverse events at dosages of 200 to 1,000 mg/day were: fatigue, nervousness, difficulty with concentration or attention, confusion, depression, anorexia, language problems, anxiety, mood problems, and weight decrease [see Table 8].

Adverse events associated with the use of topiramate at dosages of 5 to 9 mg/kg/day in controlled trials in pediatric patients with partial onset seizures, primary generalized tonic-clonic seizures, or Lennox-Gastaut syndrome, that were seen at greater frequency in topiramate-treated patients were: fatigue, somnolence, anorexia, nervousness, difficulty with concentration/attention, difficulty with memory, aggressive reaction, and weight decrease [see Table 9].

In controlled clinical trials in adults, 11% of patients receiving topiramate 200 to 400 mg/day as adjunctive therapy discontinued due to adverse events. This rate appeared to increase at dosages above 400 mg/day. Adverse events associated with discontinuing therapy included somnolence, dizziness, anxiety, difficulty with concentration or attention, fatigue, and paresthesia and increased at dosages above 400 mg/day. None of the pediatric patients who received topiramate adjunctive therapy at 5 to 9 mg/kg/day in controlled clinical trials discontinued due to adverse events.

Approximately 28% of the 1,757 adults with epilepsy who received topiramate at dosages of 200 to 1,600 mg/day in clinical studies discontinued treatment because of adverse events; an individual patient could have reported more than one adverse event. These adverse events were: psychomotor slowing (4.0%), difficulty with memory (3.2%), fatigue (3.2%), confusion (3.1%), somnolence (3.2%), difficulty with concentration/attention (2.9%), anorexia (2.7%), depression (2.6%), dizziness (2.5%), weight decrease (2.5%), nervousness (2.3%), ataxia (2.1%), and paresthesia (2.0%). Approximately 11% of the 310 pediatric patients who received topiramate at dosages up to 30 mg/kg/day discontinued due to adverse events. Adverse events associated with discontinuing therapy included aggravated convulsions (2.3%), difficulty with concentration/attention (1.6%), language problems (1.3%), personality disorder (1.3%), and somnolence (1.3%).

Incidence in Epilepsy Controlled Clinical Trials Adjunctive Therapy– Partial Onset Seizures, Primary Generalized Tonic-Clonic Seizures, and Lennox-Gastaut Syndrome

Table 6 lists treatment-emergent adverse events that occurred in at least 1% of adults treated with 200 to 400 mg/day topiramate in controlled trials that were numerically more common at this dose than in the patients treated with placebo. In general, most patients who experienced adverse events during the first eight weeks of these trials no longer experienced them by their last visit. Table 9 lists treatment-emergent adverse events that occurred in at least 1% of pediatric patients treated with 5 to 9 mg/kg topiramate in controlled trials that were numerically more common than in patients treated with placebo.

The prescriber should be aware that these data were obtained when Topiramate was added to

concurrent antiepileptic drug therapy and cannot be used to predict the frequency of adverse events in the course of usual medical practice where patient characteristics and other factors may differ from those prevailing during clinical studies. Similarly, the cited frequencies cannot be directly compared with data obtained from other clinical investigations involving different treatments, uses, or investigators. Inspection of these frequencies, however, does provide the prescribing physician with a basis to estimate the relative contribution of drug and non-drug factors to the adverse event incidences in the population studied.

Other Adverse Events Observed During Double-Blind Adjunctive Therapy Epilepsy Trials

Other events that occurred in more than 1% of adults treated with 200 to 400 mg of topiramate in placebo-controlled epilepsy trials but with equal or greater frequency in the placebo group were: headache, injury, anxiety, rash, pain, convulsions aggravated, coughing, fever, diarrhea, vomiting, muscle weakness, insomnia, personality disorder, dysmenorrhea, upper respiratory tract infection, and eye pain.

Table 6: Incidence of Treatment-Emergent Adverse Events in Placebo-Controlled, Add-On Epilepsy Trials in Adults^{a,b} Where Rate Was >1% in Any Topiramate Group and Greater Than the Rate in Placebo-Treated Patients

Body System/ Adverse Event ^c	Topiramate Dosage (mg/day)		
	Placebo (N=291)	200-400 (N=183)	600-1,000 (N=414)
Body as a Whole-			
General Disorders			
Fatigue	13	15	30
Asthenia	1	6	3
Back Pain	4	5	3
Chest Pain	3	4	2
Influenza-Like Symptoms	2	3	4
Leg Pain	2	2	4
Hot Flushes	1	2	1
Allergy	1	2	3
Edema	1	2	1
Body Odor	0	1	0
Rigors	0	1	<1
Central & Peripheral Nervous system Disorders			
Dizziness	15	25	32
Ataxia	7	16	14
Speech Disorders/ Related Speech Problems	2	13	11
Paresthesia	4	11	19
Nystagmus	7	10	11
Tremor	6	9	9
Language problems	1	6	10
Coordination Abnormal	2	4	4
Hypoaesthesia	1	2	1
Gait Abnormal	1	3	2
Muscle Contractions Involuntary			
Stupor	1	2	2
Vertigo	0	2	1
	1	1	2
Gastro-Intestinal System Disorders			
Nausea			

Dyspepsia	8	10	12
Abdominal Pain	6	7	6
Constipation	4	6	7
Gastroenteritis	2	4	3
Dry Mouth	1	2	1
Gingivitis	1	2	4
GI Disorder	<1	1	1
	<1	1	0
Hearing and Vestibular Disorders			
Hearing Decreased			
	1	2	1
Metabolic and nutritional Disorders			
Weight Decrease			
	3	9	13
Muscle-Skeletal System Disorders			
Myalgia			
Skeletal Pain	1	2	2
	0	1	0
Platelet, Bleeding, & Clotting Disorders			
Epistaxis			
	1	2	1
Psychiatric Disorders			
Somnolence	12	29	28
Nervousness	6	16	19
Psychomotor Slowing	2	13	21
Difficulty with Memory	3	12	14
Anorexia	4	10	12
Confusion	5	11	14
Depression	5	5	13
Difficulty with Concentration/ Attention			
Mood Problems	2	6	14
Agitation	2	4	9
Aggressive Reaction	2	3	3
Emotional Liability	2	3	3
Cognitive Problems	1	3	3
Libido Decreased	1	3	3
Apathy	1	2	<1
Depersonalization	1	1	3
	1	1	2
Reproductive Disorders, female			
Breast Pain			
Amenorrhea	2	4	0
Menorrhagia	1	2	2
Menstrual Disorder	0	2	1
	1	2	1
Reproductive Disorders, male			
Prostatic Disorder	<1	2	0
Resistance Mechanism Disorders			
Infection			
Infection Viral	1	2	1
Moniliasis	1	2	<1

	<1	1	0
Respiratory System Disorders			
Pharyngitis	2	6	3
Rhinitis	6	7	6
Sinusitis	4	5	6
Dyspnea	1	1	2
Skin and Appendages Disorders			
Skin Disorder			
Sweating Increased	<1	2	1
Rash Erythematous	<1	1	<1
	<1	1	<1
Special Senses Other, Disorders			
Taste Perversion			
	0	2	4
Urinary System Disorders			
Hematuria	1	2	<1
Urinary Tract Infection	1	2	3
Micturition Frequency	1	1	2
Urinary Incontinence	<1	2	1
Urine Abnormal	0	1	<1
Vision Disorders			
Vision Abnormal	2	13	10
Diplopia	5	10	10
■ White Cell and RES Disorders			
Leukopenia	1	2	1

^a Patients in these add-on trials were receiving 1 to 2 concomitant antiepileptic drugs in addition to Topiramate or placebo.

^b Values represent the percentage of patients reporting a given adverse event. Patients may have reported more than one adverse event during the study and can be included in more than one adverse event category.

^c Adverse events reported by at least 1% of patients in the Topiramate 200-400 mg/day group and more common than in the placebo group are listed in this table.

Incidence in Study 119 – Add-On Therapy– Adults with Partial Onset Seizures

Study 119 was a randomized, double-blind, placebo-controlled, parallel group study with 3 treatment arms: 1) placebo; 2) topiramate 200 mg/day with a 25 mg/day starting dose, increased by 25 mg/day each week for 8 weeks until the 200 mg/day maintenance dose was reached; and 3) topiramate 200 mg/day with a 50 mg/day starting dose, increased by 50 mg/day each week for 4 weeks until the 200 mg/day maintenance dose was reached. All patients were maintained on concomitant carbamazepine with or without another concomitant antiepileptic drug.

The incidence of adverse events (Table 7) did not differ significantly between the 2 topiramate regimens. Because the frequencies of adverse events reported in this study were markedly lower than those reported in the previous epilepsy studies, they cannot be directly compared with data obtained in other studies.

Table 7: Incidence of Treatment-Emergent Adverse Events in Study 119^{a,b} Where Rate Was $\geq 2\%$ in the Topiramate Group and Greater Than the Rate in Placebo-Treated Patients

Topiramate Dosage (mg/day)

Body System/ Adverse Event ^c	Placebo (N=92)	200 (N=171)
Body as a Whole-General Disorders		
Fatigue	4	9
Chest Pain	1	2
Cardiovascular Disorders, General		
Hypertension	0	2
Central & Peripheral Nervous system Disorders		
Paresthesia	2	9
Dizziness	4	7
Tremor	2	3
Hypoesthesia	0	2
Leg Cramps	0	2
Language Problems	0	2
Gastro-Intestinal System Disorders		
Abdominal Pain	3	5
Constipation	0	4
Diarrhea	1	2
Dyspepsia	0	2
Dry Mouth	0	2
Hearing and Vestibular Disorders		
Tinnitus	0	2
Metabolic and Nutritional Disorders		
Weight Decrease	4	8
Psychiatric Disorders		
Somnolence	9	15
Anorexia	7	9
Nervousness	2	9
Difficulty with Concentration/Attention	0	5
Insomnia		
Difficulty with Memory	3	4
Aggressive Reaction	1	2
	0	2
Respiratory System Disorders		
Rhinitis	0	4
Urinary System Disorders		
Cystitis	0	2
Vision Disorders		
Diplopia	0	2
Vision Abnormal	0	2

^a Patients in these add-on trials were receiving 1 to 2 concomitant antiepileptic drugs in addition to Topiramate or placebo.

^b Values represent the percentage of patients reporting a given adverse event. Patients may have reported more than one adverse event during the study and can be included in more than one adverse event category.

^c Adverse events reported by at least 2% of patients in the Topiramate 200 mg/day group and more common than in the placebo group are listed in this table.

Table 8: Incidence (%) of Dose-Related Adverse Events From Placebo-Controlled, Add-On Trials in Adults with Partial Onset Seizures^a

Adverse Event	Topiramate Dosage (mg/day)			
	Placebo (N=216)	200 (N=45)	400 (N=68)	600-1,000 (N=414)
Fatigue	13	11	12	30
Nervousness	7	13	18	19
Difficulty with Concentration/ Attention	1	7	9	14
Confusion	4	9	10	14
Depression	6	9	7	13
Anorexia	4	4	6	12
Language problems	<1	2	9	10
Anxiety	6	2	3	10
Mood problems	2	0	6	9
Weight decrease	3	4	9	13

^a Dose-response studies were not conducted for other adult indications or for pediatric indications.

Table 9: Incidence (%) of Treatment-Emergent Adverse Events in Placebo-Controlled, Add-On Epilepsy Trials in Pediatric Patients Ages 2-16 Years^{a,b} (Events that Occurred in at Least 1% of Topiramate-Treated Patients and Occurred More Frequently in Topiramate-Treated Than Placebo-Treated Patients)

■ Body System/ Adverse Event	Placebo (N=101)	Topiramate (N=98)
Body as a Whole-General Disorders		
Fatigue	5	16
Injury	13	14
Allergic Reaction	1	2
Back Pain	0	1
Pallor	0	1
Cardiovascular Disorders, General		
Hypertension	0	1
Central & Peripheral Nervous system Disorders		
Gait Abnormal	5	8
Ataxia	2	6
Hyperkinesia	4	5
Dizziness	2	4
Speech Disorders/Related Speech Problems	2	4
Hyporeflexia		
Convulsions Grand Mal	0	2
Fecal Incontinence	0	1
Paresthesia	0	1
	0	1
Gastro-Intestinal System Disorders		
Nausea	5	6
Saliva Increased	4	6
Constipation	4	5
Gastroenteritis	2	3
Dysphagia	0	1
Flatulence	0	1

Gastroesophageal Reflux	0	1
Glossitis	0	1
Gum Hyperplasia	0	1
Heart Rate and Rhythm Disorders		
Bradycardia	0	1
Metabolic and Nutritional Disorders		
Weight Decrease	1	9
Thirst	1	2
Hypoglycemia	0	1
Weight Increase	0	1
Platelet, Bleeding, & Clotting Disorders		
Purpura		
Epistaxis	4	8
Hematoma	1	4
Prothrombin Increased	0	1
Thrombocytopenia	0	1
	0	1
Psychiatric Disorders		
Somnolence	16	26
Anorexia	15	24
Nervousness	7	14
Personality Disorder (Behavior Problems)	9	11
Difficulty with Concentration/Attention		
Aggressive Reaction	2	10
Insomnia		
Difficulty with Memory NOS	4	9
Confusion	7	8
Psychomotor Slowing	0	5
Appetite Increased	3	4
Neurosis	2	3
	0	1
	0	1
Reproductive Disorders, female		
Leukorrhoea	0	2
Resistance Mechanism Disorders		
Infection Viral	3	7
Respiratory System Disorders		
Pneumonia	1	5
Respiratory Disorder	0	1

Placebo(N=101) Topiramate(N=98)

▪ **Body System/
Adverse Event**

Skin and Appendages Disorders		
Skin Disorder	2	3
Alopecia	1	2
Dermatitis	0	2
Hypertrichosis	1	2
Rash Erythematous	0	2
Eczema	0	1
Seborrhoea	0	1

Skin Discoloration	0	1
Urinary System Disorders		
Urinary Incontinence	2	4
Nocturia	0	1
Vision Disorders		
Eye Abnormality	1	2
Vision Abnormal	1	2
Diplopia	0	1
Lacrimation Abnormal	0	1
Myopia	0	1
White Cell and RES Disorders	0	2
Leukopenia		

^a Patients in these add-on trials were receiving 1 to 2 concomitant antiepileptic drugs in addition to Topiramate or placebo.

^b Values represent the percentage of patients reporting a given adverse event. Patients may have reported more than one adverse event during the study and can be included in more than one adverse event category.

Other Adverse Events Observed During All Epilepsy Clinical Trials

Topiramate has been administered to 2,246 adults and 427 pediatric patients with epilepsy during all clinical studies, only some of which were placebo controlled. During these studies, all adverse events were recorded by the clinical investigators using terminology of their own choosing. To provide a meaningful estimate of the proportion of individuals having adverse events, similar types of events were grouped into a smaller number of standardized categories using modified WHOART dictionary terminology. The frequencies presented represent the proportion of patients who experienced an event of the type cited on at least one occasion while receiving topiramate. Reported events are included except those already listed in the previous tables or text, those too general to be informative, and those not reasonably associated with the use of the drug.

Events are classified within body system categories and enumerated in order of decreasing frequency using the following definitions: *frequent* occurring in at least 1/100 patients; *infrequent* occurring in 1/100 to 1/1000 patients; *rare* occurring in fewer than 1/1000 patients.

Autonomic Nervous System Disorders:*Infrequent:* vasodilation.

Body as a Whole:*Frequent:* syncope. *Infrequent:* abdomen enlarged. *Rare:* alcohol intolerance.

Cardiovascular Disorders, General:*Infrequent:* hypotension, postural hypotension, angina pectoris.

Central & Peripheral Nervous System Disorders:*Infrequent:* neuropathy, apraxia, hyperaesthesia, dyskinesia, dysphonia, scotoma, ptosis, dystonia, visual field defect, encephalopathy, EEG abnormal. *Rare:* upper motor neuron lesion, cerebellar syndrome, tongue paralysis.

Gastrointestinal System Disorders:*Infrequent:* hemorrhoids, stomatitis, melena, gastritis, esophagitis. *Rare:* tongue edema.

Heart Rate and Rhythm Disorders:*Infrequent:* AV block.

Liver and Biliary System Disorders: *Infrequent:* SGPT increased, SGOT increased.

Metabolic and Nutritional Disorders:*Infrequent:* dehydration, hypokalemia, alkaline phosphatase increased, hypocalcemia, hyperlipemia, hyperglycemia, xerophthalmia, diabetes mellitus. *Rare:* hyperchloremia, hypernatremia, hyponatremia, hypocholesterolemia, hypophosphatemia, creatinine increased.

Musculoskeletal System Disorders: *Frequent:* Arthralgia. *Infrequent:* arthrosis.

Neoplasms: *Infrequent:* thrombocythemia. *Rare:* polycythemia.

Platelet, Bleeding, and Clotting Disorders: *Infrequent:* gingival bleeding, pulmonary embolism.

Psychiatric Disorders: *Frequent:* impotence, hallucination, psychosis, suicide attempt. *Infrequent:* euphoria, paranoid reaction, delusion, paranoia, delirium, abnormal dreaming. *Rare:* libido increased, manic reaction.

Red Blood Cell Disorders: *Frequent:* anemia. *Rare:* marrow depression, pancytopenia.

Reproductive Disorders, Male: *Infrequent:* ejaculation disorder, breast discharge.

Skin and Appendages Disorders: *Infrequent:* urticaria, photosensitivity reaction, abnormal hair texture. *Rare:* chloasma.

Special Senses Other, Disorders: *Infrequent:* taste loss, parosmia.

Urinary System Disorders: *Infrequent:* urinary retention, face edema, renal pain, albuminuria, polyuria, oliguria.

Vascular (Extracardiac) Disorders: *Infrequent:* flushing, deep vein thrombosis, phlebitis. *Rare:* vasospasm.

Vision Disorders: *Frequent:* conjunctivitis. *Infrequent:* abnormal accommodation, photophobia, strabismus.

Rare: mydriasis, iritis.

White Cell and Reticuloendothelial System Disorders: *Infrequent:* lymphadenopathy, eosinophilia, lymphopenia, granulocytopenia. *Rare:* lymphocytosis.

Postmarketing and Other Experience

In addition to the adverse experiences reported during clinical testing of Topiramate, the following adverse experiences have been reported worldwide in patients receiving topiramate post-approval. These adverse experiences have not been listed above and data are insufficient to support an estimate of their incidence or to establish causation. The listing is alphabetized: bullous skin reactions (including erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis), hepatic failure (including fatalities), hepatitis, pancreatitis, pemphigus, and renal tubular acidosis.

DRUG ABUSE AND DEPENDENCE

The abuse and dependence potential of Topiramate has not been evaluated in human studies.

OVERDOSAGE

Overdoses of Topiramate have been reported. Signs and symptoms included convulsions, drowsiness, speech disturbance, blurred vision, diplopia, mentation impaired, lethargy, abnormal coordination, stupor, hypotension, abdominal pain, agitation, dizziness and depression. The clinical consequences were not severe in most cases, but deaths have been reported after poly-drug overdoses involving Topiramate.

Topiramate overdose has resulted in severe metabolic acidosis (see **WARNINGS**).

A patient who ingested a dose between 96 and 110 g topiramate was admitted to hospital with coma lasting 20-24 hours followed by full recovery after 3 to 4 days.

In acute Topiramate overdose, if the ingestion is recent, the stomach should be emptied immediately by lavage or by induction of emesis. Activated charcoal has been shown to adsorb topiramate *in vitro*.

Treatment should be appropriately supportive. Hemodialysis is an effective means of removing topiramate from the body.

DOSAGE AND ADMINISTRATION

Epilepsy

In the controlled add-on trials, no correlation has been demonstrated between trough plasma concentrations of topiramate and clinical efficacy. No evidence of tolerance has been demonstrated in humans. Doses above 400 mg/day (600, 800, or 1000 mg/day) have not been shown to improve responses in dose-response studies in adults with partial onset seizures.

It is not necessary to monitor topiramate plasma concentrations to optimize Topiramate therapy. On occasion, the addition of Topiramate to phenytoin may require an adjustment of the dose of phenytoin to achieve optimal clinical outcome. Addition or withdrawal of phenytoin and/or carbamazepine during adjunctive therapy with Topiramate may require adjustment of the dose of Topiramate. Because of the bitter taste, tablets should not be broken.

Topiramate can be taken without regard to meals.

Monotherapy Use

The recommended dose for topiramate monotherapy in adults and children 10 years of age and older is 400 mg/day in two divided doses. Approximately 58% of patients randomized to 400 mg/day achieved this maximal dose in the monotherapy controlled trial; the mean dose achieved in the trial was 275 mg/day. The dose should be achieved by titrating according to the following schedule:

	Morning Dose	Evening Dose
Week 1	25 mg	25 mg
Week 2	50 mg	50 mg
Week 3	75 mg	75 mg
Week 4	100 mg	100 mg
Week 5	150 mg	150 mg
Week 6	200 mg	200 mg

Adjunctive Therapy Use

Adults (17 Years of Age and Over) - Partial Seizures, Primary Generalized Tonic-Clonic Seizures, or Lennox-Gastaut Syndrome

The recommended total daily dose of Topiramate as adjunctive therapy in adults with partial seizures is 200-400 mg/day in two divided doses, and 400 mg/day in two divided doses as adjunctive treatment in adults with primary generalized tonic-clonic seizures. It is recommended that therapy be initiated at 25-50 mg/day followed by titration to an effective dose in increments of 25-50 mg/week. Titrating in increments of 25 mg/week may delay the time to reach an effective dose. Daily doses above 1,600 mg have not been studied.

In the study of primary generalized tonic-clonic seizures the initial titration rate was slower than in previous studies; the assigned dose was reached at the end of 8 weeks (see **CLINICAL STUDIES, Adjunctive Therapy Controlled Trials in Patients With Primary Generalized Tonic-Clonic Seizures**).

Pediatric Patients (Ages 2 - 16 Years) - Partial Seizures, Primary Generalized Tonic-Clonic Seizures, or Lennox-Gastaut Syndrome

The recommended total daily dose of Topiramate as adjunctive therapy for patients with partial seizures, primary generalized tonic-clonic seizures, or seizures associated with Lennox-Gastaut

Syndrome is approximately 5 to 9 mg/kg/day in two divided doses. Titration should begin at 25 mg (or less, based on a range of 1 to 3 mg/kg/day) nightly for the first week. The dosage should then be increased at 1- or 2-week intervals by increments of 1 to 3 mg/kg/day (administered in two divided doses), to achieve optimal clinical response. Dose titration should be guided by clinical outcome.

In the study of primary generalized tonic-clonic seizures the initial titration rate was slower than in previous studies; the assigned dose of 6 mg/kg/day was reached at the end of 8 weeks (see **CLINICAL STUDIES, Adjunctive Therapy Controlled Trial in Patients With Primary Generalized Tonic-Clonic Seizures**).

Patients with Renal Impairment:

In renally impaired subjects (creatinine clearance less than 70 mL/min/1.73m²), one half of the usual adult dose is recommended. Such patients will require a longer time to reach steady-state at each dose.

Geriatric Patients (Ages 65 Years and Over):

Dosage adjustment may be indicated in the elderly patient when impaired renal function (creatinine clearance rate ≤ 70 mL/min/1.73 m²) is evident (see **DOSAGE AND ADMINISTRATION:Patients with Renal Impairment and CLINICAL PHARMACOLOGY:Special Populations:Age, Gender, and Race**).

Patients Undergoing Hemodialysis:

Topiramate is cleared by hemodialysis at a rate that is 4 to 6 times greater than a normal individual. Accordingly, a prolonged period of dialysis may cause topiramate concentration to fall below that required to maintain an anti-seizure effect. To avoid rapid drops in topiramate plasma concentration during hemodialysis, a supplemental dose of topiramate may be required. The actual adjustment should take into account 1) the duration of dialysis period, 2) the clearance rate of the dialysis system being used, and 3) the effective renal clearance of topiramate in the patient being dialyzed.

Patients with Hepatic Disease:

In hepatically impaired patients topiramate plasma concentrations may be increased. The mechanism is not well understood.

HOW SUPPLIED

Topiramate Tablets are available containing 50 mg or 100 of topiramate USP.

The 50 mg tablets are yellow, round, biconvex tablets de-bossed with **IG** on one side and **279** on other. They are available as follows:

NDC 33261-0400-07 bottles of 7 tablets

NDC 33261-0400-14 bottles of 14 tablets

NDC 33261-0400-28 bottles of 28 tablets

NDC 33261-0400-30 bottles of 30 tablets

NDC 33261-0400-60 bottles of 60 tablets

NDC 33261-0400-90 bottles of 90 tablets

NDC 33261-0400-02 bottles of 120 tablets

The 100 mg tablets are light yellow, round, biconvex tablets de-bossed with **IG** on one side and **280** on other. They are available as follows:

NDC 33261-480-60 bottles of 60 tablets

Store at 20° to 25°C (68° to 77°F); [See USP Controlled Room Temperature].


Protect from moisture.

PATIENT INFORMATION

Topiramate Tablets

What do TOPIRAMATE Tablets look like?

TOPIRAMATE Tablets



25mg	50mg	100mg	200mg
White	Yellow	Light	Pink

Yellow

Note: The pictures above show the shapes and lettering of TOPIRAMATE tablets. The wording describes the strength and colors of the medication. Before taking your medicine, it is important to compare the tablets you receive from your healthcare professional or pharmacist with these pictures to make sure you have received the correct medicine.

Please read this patient information carefully before you take TOPIRAMATE and each time you obtain a refill, in case any information has changed. This summary does not contain all the information about TOPIRAMATE and is not meant to take the place of talking with your healthcare professional. If you have any questions about TOPIRAMATE, discuss them with your healthcare professional or pharmacist.

What is TOPIRAMATE?

TOPIRAMATE is a prescription medicine used:

- alone to treat seizures in patients 10 years and older
- with other medicines to treat seizures in adults and children over age 2

Who Should Not Take TOPIRAMATE?

Do not take TOPIRAMATE if you are allergic to anything in it. See the end of this leaflet for a complete list of ingredients in TOPIRAMATE.

What Should I Tell My Healthcare Professional Before Taking TOPIRAMATE?

Tell your healthcare professional about all of your medical conditions, including if you:

- have kidney problems, especially kidney stones, or are getting kidney dialysis
- have a history of metabolic acidosis (blood and body fluid abnormality)
- have liver problems

- have osteoporosis (weak or brittle bones) and/or soft bones (osteomalacia) or decreased bone density (osteopenia)
- have lung or breathing problems
- have eye problems, especially glaucoma
- have diarrhea
- have a growth problem
- are on a diet high in fat called a ketogenic diet
- are having surgery
- are pregnant or planning to become pregnant. It is not known if TOPIRAMATE can harm your unborn baby.
- are breastfeeding. TOPIRAMATE may pass into your milk. Talk to your healthcare professional about the best way to feed your baby while taking TOPIRAMATE.
- suffer from depression, mood problems or suicidal thoughts or behavior

Tell your healthcare professional about all the medicines you take including prescription and nonprescription medicines, vitamins and herbal supplements. TOPIRAMATE and certain other medicines can affect each other. Sometimes the dose of some of your other medicines or TOPIRAMATE will have to be adjusted. Especially, tell your healthcare professional if you are taking:

- other medicines that impair or decrease your thinking, concentration, or muscle coordination (e.g. central nervous system depressant medicines).
- birth control pills. TOPIRAMATE may make your birth control pills less effective. Tell your healthcare professional if your menstrual bleeding changes while you are taking birth control pills and TOPIRAMATE.

Keep a list of all the medicines you take. Show this list to your healthcare professionals and pharmacists before you start a new medicine.

How Should I Take TOPIRAMATE?

- Take TOPIRAMATE exactly as prescribed. Your healthcare professional will usually start you on a low dose of TOPIRAMATE and slowly increase your dose until the best dose is found for you.
- TOPIRAMATE Tablets should be swallowed whole. Avoid, chewing the tablets as they may leave a bitter taste.
- Never store any medicine and food mixture for use at a later time.
- TOPIRAMATE can be taken before, during, or after a meal. Drink plenty of fluids during the day to prevent kidney stones while taking TOPIRAMATE.
- If you take too much TOPIRAMATE, call your healthcare professional or poison control center right away or go to an emergency room.
- If you miss a single dose of TOPIRAMATE, take it as soon as you can. However, if you are within 6 hours of taking your next scheduled dose, wait until then to take your usual dose of TOPIRAMATE, and skip the missed dose. Do not double your dose. If you have missed more than one dose, you should call your healthcare professional for advice.
- Do not stop taking TOPIRAMATE unless a healthcare professional tells you to stop taking TOPIRAMATE. Your healthcare professional will tell you how to slowly stop taking TOPIRAMATE.

What Should I Avoid While Taking TOPIRAMATE?

- If you are taking Topiramate or other antiepileptic drugs for epilepsy or seizures, you may need to avoid activities where loss of consciousness (passing out) could result in serious danger to yourself or those around you (including swimming, driving a car, climbing in high places, etc.). Talk to your doctor before engaging in such activities.
- Unless prescribed by your healthcare professional, you should avoid other medicines that also impair or decrease your thinking, concentration, or muscle coordination (e.g. central nervous system depressant medicines).

- You should avoid drinking alcohol while taking TOPIRAMATE. Alcohol with TOPIRAMATE can make side effects such as sleepiness and dizziness worse.
- Do not drive a car or operate heavy machinery until you know how TOPIRAMATE affects you. TOPIRAMATE can impair your thinking, motor skills, and/or vision.

What are the Possible Side Effects of TOPIRAMATE?

TOPIRAMATE may cause the following side effects which can be serious:

- **metabolic acidosis.** Metabolic acidosis is a condition that happens when there is too much acid in your blood. Metabolic acidosis can cause symptoms such as tiredness, loss of appetite, irregular heartbeat, and impaired consciousness. **Call your healthcare professional right away if you get these symptoms with TOPIRAMATE.** Your healthcare professional should do a blood test (measurement of serum bicarbonate) to monitor your bicarbonate level while you are taking TOPIRAMATE.
- eye problems. Serious eye problems include:
- a sudden decrease in vision (acute myopia) with or without eye pain and
- a blockage of fluid in the eye causing increased pressure in the eye (secondary angle closure glaucoma).

Call your healthcare professional right away if you have a loss in vision or get eye pain. These problems can lead to blindness if not treated right away. Your healthcare professional will probably stop TOPIRAMATE and may recommend other therapy.

- **decreased sweating (oligohidrosis) and increased body temperature (fever).** Patients, especially children, should be watched closely for signs of decreased sweating and fever (increased body temperature), especially in hot temperatures. Some patients may need hospital treatment for this condition.
- **effects on thinking and alertness.** TOPIRAMATE may affect thinking skills and cause confusion, problems with concentration, attention, memory, and/or speech. TOPIRAMATE may cause depression or mood problems, tiredness, and sleepiness. Call your healthcare professional right away if you experience any of these side effects.
- **dizziness or loss of muscle coordination** in patients who take TOPIRAMATE alone or with other seizure medicines.
- **high blood ammonia levels and effects on mental activities.** High ammonia in the blood can affect your mental activities and decrease alertness, can make you feel tired or fatigued, or can cause vomiting. This has happened when TOPIRAMATE has been used with a medicine called valproic acid.
- **kidney stones.** Drink plenty of fluids when taking TOPIRAMATE to decrease your chances of getting kidney stones.
- **tingling of the arms and legs (paresthesia)** is a common side effect of TOPIRAMATE.

Other side effects with TOPIRAMATE include loss of appetite, nausea, a change in the way foods taste, diarrhea, weight loss, nervousness, aggression, upper respiratory tract infection. Call your healthcare professional if you have any symptoms that concern you or that do not go away. These are not all the side effects with TOPIRAMATE. For more information, ask your healthcare professional or pharmacist.

What Should I Do If I Get Pregnant While Taking TOPIRAMATE?

It is not clear if there is a risk to the fetus/baby if you are exposed to TOPIRAMATE and you are pregnant. Various abnormalities have been described in the offspring of animals exposed to TOPIRAMATE during pregnancy. If you use TOPIRAMATE while you are pregnant, ask your healthcare professional about reporting your experience to the North American Drug Pregnancy Registry at Massachusetts General Hospital (Boston, MA). This registry collects information about the babies born to women who are taking drugs to treat various conditions. Information about the North

American Drug Pregnancy Registry can be found at <http://www.massgeneral.org/aed/>. You can also join the registry by calling 1-877-376-3872.

How Should I Store TOPIRAMATE?

- Store at 20° to 25°C (68° to 77°F); [See USP Controlled Room Temperature]. Protect from moisture.
- **Keep TOPIRAMATE and all medicines out of the reach of children.**

General Information About TOPIRAMATE.

Medicines are sometimes prescribed for purposes other than those listed in Patient Information leaflets. Do not use TOPIRAMATE for a condition for which it was not prescribed. Do not give TOPIRAMATE to other people, even if they have the same symptoms that you have. It may harm them.

This leaflet summarizes the most important information about TOPIRAMATE. If you would like more information, talk to your healthcare professional. You can ask your health care professional or pharmacist for information about TOPIRAMATE that is written for health professionals.

What Are the Ingredients of TOPIRAMATE?

Active Ingredient: topiramate

Inactive Ingredients:

- **Tablets** - lactose monohydrate, microcrystalline cellulose, pre-gelatinized starch, lactose monohydrate, sodium starch glycolate, magnesium stearate, opadry white (titanium dioxide, hypromellose 3cp, hypromellose 6cp, PEG 400, polysorbate 80) for 25 mg tablets, opadry yellow (titanium dioxide, hypromellose 3cp, hypromellose 6cp, PEG 400, polysorbate 80, iron oxide yellow) for 50 mg tablets, opadry yellow , hypromellose 6cp titanium dioxide, PEG 400, iron oxide yellow, polysorbate 80, iron oxide red) for 100 mg tablets and), opadry pink (titanium dioxide, hypromellose 6cp, PEG 400, iron oxide red) for 200 mg tablets.

Manufactured by:
InvaGen Pharmaceuticals, Inc.
Hauppauge, NY 11788

Manufactured for
Camber Pharmaceuticals, Inc
Piscataway, NJ 08854

Repackaged By:
Aidarex Pharmaceuticals, LLC.
Corona, CA 92880

Rev: 03/09

Barcode
281-03-2009

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT. KEEP OUT OF REACH OF CHILDREN. STORE AT CONTROLLED ROOM TEMP 15-30C (59-86F)

Packaged and Distributed by: **AIDAREX PHARMACEUTICALS LLC.**

TOPIRAMATE

50mg
60 TABS

EACH TABLET CONTAINS THE FOLLOWING ACTIVE INGREDIENTS:
TOPIRAMATE.....50 MG

YELLOW ROUND TABLET W/ 279 ON ONE SIDE & IG ON REVERSE

RX QLS0000

GENERIC FOR : TOPAMAX
NDC: 33261-0400-60

TAKE _____ EVERY _____ HOURS _____ TIMES A DAY
TOME _____ CADA _____ HORAS _____ VECES AL DIA

MFG: INVAGEN PHARMACEUTICALS, INC., HAUPPAUGE, NY 11788

TOPIRAMATE 50mg
NDC: 33261-0400-60
RX QLS0000 60

TOPIRAMATE 50mg
NDC: 33261-0400-60
RX QLS0000 60

TOPIRAMATE 50mg
NDC: 33261-0400-60
RX QLS0000 60

PATIENT LOG CHART BILL

PEEL HERE

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT. KEEP OUT OF REACH OF CHILDREN. STORE AT CONTROLLED ROOM TEMP 15-30C (59-86F)

Packaged and Distributed by: **AIDAREX PHARMACEUTICALS LLC.**

TOPIRAMATE

100 MG
60 TABS

EACH TABLET CONTAINS THE FOLLOWING ACTIVE INGREDIENTS:
TOPIRAMATE.....100 MG
FILM-COATED

YELLOW ROUND TABLET W/ I G ON ONE SIDE & 280 ON REVERSE

RX QLS0000

GENERIC FOR : TOPAMAX
NDC: 33261-0480-60

TAKE _____ EVERY _____ HOURS _____ TIMES A DAY
TOME _____ CADA _____ HORAS _____ VECES AL DIA

MFG: INVAGEN PHARMACEUTICALS, INC., HAUPPAUGE, NY 11788

TOPIRAMATE 100 MG
NDC: 33261-0480-60
RX QLS0000 60

TOPIRAMATE 100 MG
NDC: 33261-0480-60
RX QLS0000 60

TOPIRAMATE 100 MG
NDC: 33261-0480-60
RX QLS0000 60

PATIENT LOG CHART BILL

PEEL HERE

TOPIRAMATE

topiramate tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:33261-400(NDC:31722-279)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TOPIRAMATE (UNII: 0H73WJJ39 1) (TOPIRAMATE - UNII:0H73WJJ39 1)	TOPIRAMATE	50 mg

Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYPROMELLOSE 2910 (3 MPA.S) (UNII: 0VUT3PMY82)	
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ8WG20P6)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	

Product Characteristics

Color	YELLOW	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	IG;279
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:33261-400-02	120 in 1 BOTTLE, PLASTIC		
2	NDC:33261-400-07	7 in 1 BOTTLE, PLASTIC		
3	NDC:33261-400-14	14 in 1 BOTTLE, PLASTIC		
4	NDC:33261-400-28	28 in 1 BOTTLE, PLASTIC		
5	NDC:33261-400-30	30 in 1 BOTTLE, PLASTIC		
6	NDC:33261-400-60	60 in 1 BOTTLE, PLASTIC		
7	NDC:33261-400-90	90 in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079162	01/01/2011	

TOPIRAMATE

topiramate tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:33261-480(NDC:31722-280)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TOPIRAMATE (UNII: 0H73WJJ391) (TOPIRAMATE - UNII:0H73WJJ391)	TOPIRAMATE	100 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYPROMELLOSE 2910 (3 MPAS) (UNII: 0VUT3PMY82)	
HYPROMELLOSE 2910 (6 MPAS) (UNII: 0WZ8WG20P6)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

Product Characteristics

Color	YELLOW (Light yellow)	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	IG;280
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:33261-480-60	60 in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079162	01/01/2011	

Labeler - Aidarex Pharmaceuticals LLC (801503249)

Revised: 1/2014

Aidarex Pharmaceuticals LLC