WEGOVY- semaglutide injection, solution
WEGOVY- semaglutide tablet
Novo Nordisk Pharmaceutical Industries, LP

HIGHLIGHTS OF PRESCRIBING INFORMATION
These highlights do not include all the information needed to use WEGOVY® safely and effectively. See full prescribing information for WEGOVY.
WEGOVY (semaglutide) injection, for subcutaneous use
WEGOVY (semaglutide) tablets, for oral use
Initial U.S. Approval: 2017

WARNING: RISK OF THYROID C-CELL TUMORS

See full prescribing information for complete boxed warning.

- **In rodents, semaglutide causes thyroid C-cell tumors at clinically relevant exposures. It is unknown whether WEGOVY causes thyroid C-cell tumors, including medullary thyroid carcinoma (MTC), in humans as the human relevance of semaglutide-induced rodent thyroid C-cell tumors has not been determined. (5.1, 13.1)**
- **WEGOVY is contraindicated in patients with a personal or family history of MTC or in patients with Multiple Endocrine Neoplasia syndrome type 2 (MEN 2). Counsel patients regarding the potential risk of MTC and symptoms of thyroid tumors. (4, 5.1)**

RECENT MAJOR CHANGES

Indications and Usage (1) 08/2025, 12/2025
Dosage and Administration (2)..... 08/2025, 12/2025, 03/2026
Warnings and Precautions,

Acute Kidney Injury due to Volume Depletion (5.5).....08/2025
Severe Gastrointestinal Adverse Reactions (5.6).....10/2025
Suicidal Behavior and Ideation (5.10).....(Removed) 02/2026
Never Share WEGOVY FlexTouch Between Patients (5.11).....06/2026

INDICATIONS AND USAGE

WEGOVY is a glucagon-like peptide-1 (GLP-1) receptor agonist.
WEGOVY injection is indicated in combination with a reduced calorie diet and increased physical activity:

- to reduce the risk of major adverse cardiovascular (CV) events (CV death, non-fatal myocardial infarction, or non-fatal stroke) in adults with established CV disease and either obesity or overweight. (1)
- to reduce excess body weight and maintain weight reduction long term in:
  - adults and pediatric patients aged 12 years and older with obesity. (1)
  - adults with overweight in the presence of at least one weight-related comorbid condition. (1)
- for the treatment of noncirrhotic metabolic dysfunction-associated steatohepatitis (MASH), formerly known as nonalcoholic steatohepatitis (NASH), with moderate to advanced liver fibrosis (consistent with stages F2 to F3 fibrosis) in adults. This indication is approved under accelerated approval based on improvement of MASH and fibrosis. Continued approval for this indication may be contingent upon the verification and description of clinical benefit in a confirmatory trial. (1)

WEGOVY tablets are indicated in combination with a reduced calorie diet and increased physical activity:

- to reduce the risk of major adverse CV events (CV death, non-fatal myocardial infarction, or non-fatal stroke) in adults with established CV disease and either obesity or overweight. (1)
- to reduce excess body weight and maintain weight reduction long term in adults with obesity, or in adults with overweight in the presence of at least one weight-related comorbid condition. (1)

Limitations of Use:

- Concomitant use of WEGOVY (semaglutide) tablets or WEGOVY (semaglutide) injection with other semaglutide-containing products or with any other GLP-1 receptor agonist is not recommended. (1)

DOSAGE AND ADMINISTRATION

In patients with diabetes, monitor blood glucose prior to starting and during WEGOVY treatment. (2.1)

WEGOVY Injection

- Administer WEGOVY injection once weekly as an adjunct to diet and increased physical activity, on the same day each week, at any time of day, with or without meals. (2.1)
- Inject subcutaneously in the abdomen, thigh, or upper arm. (2.1)
- Initiate at 0.25 mg once weekly for 4 weeks. Then follow the dosage escalation schedule in Table 1, titrating every 4 weeks to achieve the maintenance dosage. (2.2)
- The usual recommended maintenance dosage of WEGOVY injection is 2.4 mg once weekly. Refer to the full PI for maintenance dosages based on the indication. (2.2)

WEGOVY Tablets

- Take WEGOVY tablets orally once-daily on an empty stomach in the morning with water (up to 4 ounces). Do not take with other liquids besides water. (2.1)
- Swallow tablets whole. Do not split, crush, chew or dissolve. (2.1)
- After taking WEGOVY tablet wait at least 30 minutes before eating food, drinking beverages or taking other oral medications. (2.1)
- Initiate WEGOVY tablet with a dosage of 1.5 mg once daily for 30 days. Then follow the dosage escalation schedule, titrating every 30 days to achieve the maintenance dosage. (2.2)
- The recommended maintenance dosage of WEGOVY tablets is 25 mg orally once daily for cardiovascular risk reduction and weight reduction in adults. (2.3)

DOSAGE FORMS AND STRENGTHS

- Injection (3):
  - 0.25 mg/0.5 mL, 0.5 mg/0.5 mL, 1 mg/0.5 mL, 1.7 mg/0.75 mL, 2.4 mg/0.75 mL, or 7.2 mg/0.75 mL (WEGOVY HD) in a single-dose pen
  - 0.25 mg/0.5 mL, 0.5 mg/0.5 mL, 1 mg/0.5 mL, 1.7 mg/0.75 mL, or 2.4 mg/0.75 mL in a single-dose syringe
  - 9.6 mg/3 mL (3.2 mg/mL) solution in a single-patient-use pen (WEGOVY FlexTouch) that delivers four 2.4 mg/0.75 mL doses
- Tablets: 1.5 mg, 4 mg, 9 mg and 25 mg (3)

CONTRAINDICATIONS

- Personal or family history of MTC or in patients with MEN 2. (4)
- Known hypersensitivity to semaglutide or any of the excipients in WEGOVY tablets or WEGOVY injection. (4)

WARNINGS AND PRECAUTIONS

- *Acute Pancreatitis*: Has been observed in patients treated with GLP-1 receptor agonists, including WEGOVY. Discontinue if pancreatitis is suspected. (5.2)
- *Acute Gallbladder Disease*: Has occurred in clinical trials. If cholelithiasis is suspected, gallbladder studies and clinical follow-up are indicated. (5.3)
- *Hypoglycemia*: Concomitant use with insulin or an insulin secretagogue may increase the risk of hypoglycemia, including severe hypoglycemia. Reducing the dose of insulin or insulin secretagogue may be necessary. Inform all patients of the risk of hypoglycemia and educate them on the signs and symptoms of hypoglycemia. (5.4)
- *Acute Kidney Injury Due to Volume Depletion*: Monitor renal function in patients reporting adverse reactions that could lead to volume depletion. (5.5)
- *Severe Gastrointestinal Adverse Reactions*: Use has been associated with gastrointestinal adverse reactions, sometimes severe. WEGOVY is not recommended in patients with severe gastroparesis. (5.6)
- *Hypersensitivity Reactions*: Anaphylactic reactions and angioedema have been reported postmarketing. Discontinue WEGOVY if suspected and promptly seek medical advice. (5.7)
- *Diabetic Retinopathy Complications in Patients with Type 2 Diabetes*: Has been reported in trials with semaglutide. Patients with a history of diabetic retinopathy should be monitored. (5.8)
- *Heart Rate Increase*: Monitor heart rate at regular intervals. (5.9)
- *Pulmonary Aspiration During General Anesthesia or Deep Sedation*: Has been reported in patients receiving GLP-1 receptor agonists undergoing elective surgeries or procedures. Instruct patients to inform healthcare providers of any planned surgeries or procedures. (5.10)
- *Never Share WEGOVY FlexTouch Between Patients*, even if the needle is changed. (5.11)

ADVERSE REACTIONS

Most common adverse reactions (incidence ≥5%) in adults or pediatric patients aged 12 years and older are: nausea, diarrhea, vomiting, constipation, abdominal pain, dysesthesia, headache, fatigue, dyspepsia, dizziness, abdominal distension, eructation, hypoglycemia in patients with type 2 diabetes, flatulence, gastroenteritis, gastroesophageal reflux disease, and hair loss. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Novo Nordisk Inc., at 1-833-934-6891 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

WEGOVY delays gastric emptying. May impact absorption of concomitantly administered oral medications. Consider increased clinical or laboratory monitoring when used concomitantly with other oral medications that have a narrow therapeutic index or that require clinical monitoring. (7.2)

USE IN SPECIFIC POPULATIONS

- *Pregnancy*: May cause fetal harm. For patients receiving WEGOVY for CV risk reduction or weight reduction, discontinue WEGOVY when pregnancy is recognized. For patients with MASH, use during pregnancy only if the potential benefit justifies the potential risk to the fetus. (8.1)
- *Lactation*: Breastfeeding not recommended during treatment with WEGOVY tablets. (8.2)
- *Females and Males of Reproductive Potential*: For patients receiving WEGOVY for CV risk reduction or weight reduction, or for MASH where the potential risk outweighs the potential benefit, discontinue WEGOVY at least 2 months before a planned pregnancy because of the long half-life of semaglutide. (8.3)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 6/2026

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: RISK OF THYROID C-CELL TUMORS
1 INDICATIONS AND USAGE
2 DOSAGE AND ADMINISTRATION
2.1 Important Monitoring and Administration Instructions
2.2 Recommended Dosage for WEGOVY Injection
2.3 Recommended Dosage of WEGOVY Tablets
2.4 Recommendations Regarding Missed Dose(s)
2.5 Switching Between WEGOVY Injection and WEGOVY Tablets
3 DOSAGE FORMS AND STRENGTHS
4 CONTRAINDICATIONS
5 WARNINGS AND PRECAUTIONS
5.1 Risk of Thyroid C-Cell Tumors
5.2 Acute Pancreatitis
5.3 Acute Gallbladder Disease
5.4 Hypoglycemia
5.5 Acute Kidney Injury Due to Volume Depletion
5.6 Severe Gastrointestinal Adverse Reactions
5.7 Hypersensitivity Reactions
5.8 Diabetic Retinopathy Complications in Patients with Type 2 Diabetes
5.9 Heart Rate Increase
5.10 Pulmonary Aspiration During General Anesthesia or Deep Sedation
5.11 Never Share WEGOVY FlexTouch Between Patients
6 ADVERSE REACTIONS
6.1 Clinical Trials Experience
6.2 Postmarketing Experience
7 DRUG INTERACTIONS

7.1 Concomitant Use with Insulin or an Insulin Secretagogue (e.g., Sulfonylurea)
7.2 Oral Medications
8 USE IN SPECIFIC POPULATIONS
8.1 Pregnancy
8.2 Lactation
8.3 Females and Males of Reproductive Potential
8.4 Pediatric Use
8.5 Geriatric Use
8.6 Type 2 Diabetes Mellitus
10 OVERDOSAGE
11 DESCRIPTION
12 CLINICAL PHARMACOLOGY
12.1 Mechanism of Action
12.2 Pharmacodynamics
12.3 Pharmacokinetics
12.6 Immunogenicity
13 NONCLINICAL TOXICOLOGY
13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
13.2 Animal Toxicology and/or Pharmacology
14 CLINICAL STUDIES
14.1 Cardiovascular Outcomes Trial in Adult Patients with Cardiovascular Disease and Either Obesity or Overweight
14.2 Weight Reduction and Long-term Maintenance Studies in Adults with Obesity or Overweight
14.3 Weight Reduction and Long-term Maintenance Study of WEGOVY Injection in Pediatric Patients Aged 12 Years and Older with Obesity
14.4 Noncirrhotic Metabolic Dysfunction-associated Steatohepatitis with Moderate to Advanced Liver Fibrosis in Adults Treated with WEGOVY Injection
16 HOW SUPPLIED/STORAGE AND HANDLING
17 PATIENT COUNSELING INFORMATION
* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: RISK OF THYROID C-CELL TUMORS

- **In rodents, semaglutide causes dose-dependent and treatment-duration-dependent thyroid C-cell tumors at clinically relevant exposures. It is unknown whether WEGOVY causes thyroid C-cell tumors, including medullary thyroid carcinoma (MTC), in humans as human relevance of semaglutide-induced rodent thyroid C-cell tumors has not been determined *[see Warnings and Precautions (5.1), Nonclinical Toxicology (13.1)]*.**
- **WEGOVY is contraindicated in patients with a personal or family history of MTC or in patients with Multiple Endocrine Neoplasia syndrome type 2 (MEN 2) *[see Contraindications (4)]*. Counsel patients regarding the potential risk for MTC with the use of WEGOVY and inform them of symptoms of thyroid tumors (e.g., a mass in the neck, dysphagia, dyspnea, persistent hoarseness). Routine monitoring of serum calcitonin or using thyroid ultrasound is of uncertain value for early detection of MTC in patients treated with WEGOVY *[see Contraindications (4), Warnings and Precautions (5.1)]*.**

1 INDICATIONS AND USAGE

WEGOVY injection is indicated in combination with a reduced calorie diet and increased physical activity:

- to reduce the risk of major adverse cardiovascular (CV) events (CV death, non-fatal myocardial infarction, or non-fatal stroke) in adults with established CV disease and either obesity or overweight.
- to reduce excess body weight and maintain weight reduction long term in:
    - adults and pediatric patients aged 12 years and older with obesity.
    - adults with overweight in the presence of at least one weight-related comorbid condition.
- for the treatment of noncirrhotic metabolic dysfunction-associated steatohepatitis (MASH), formerly known as nonalcoholic steatohepatitis (NASH), with moderate to advanced liver fibrosis (consistent with stages F2 to F3 fibrosis) in adults. This indication is approved under accelerated approval based on improvement of MASH and fibrosis *[see Clinical Studies (14.4)]*. Continued approval for this indication may be contingent upon the verification and description of clinical benefit in a confirmatory trial.

WEGOVY tablets are indicated in combination with a reduced calorie diet and increased physical activity:

- to reduce the risk of major adverse CV events (CV death, non-fatal myocardial infarction, or non-fatal stroke) in adults with established CV disease and either obesity or overweight.
- to reduce excess body weight and maintain weight reduction long term in adults with obesity, or in adults with overweight in the presence of at least one weight-

related comorbid condition.

Limitations of Use

Concomitant use of WEGOVY (semaglutide) tablets or WEGOVY (semaglutide) injection with other semaglutide-containing products or with any other GLP-1 receptor agonist is not recommended.

2 DOSAGE AND ADMINISTRATION

2.1 Important Monitoring and Administration Instructions

In patients with diabetes mellitus, monitor blood glucose prior to starting WEGOVY and during WEGOVY treatment *[see Warnings and Precautions (5.4)]*.

Administer WEGOVY in combination with a reduced-calorie diet and increased physical activity.

WEGOVY Injection

- Inform patients and their caregiver(s) which WEGOVY presentation (e.g., single-dose pen or syringe, or WEGOVY FlexTouch single-patient-use pen) they will receive and ensure they receive training appropriate for that specific presentation. If the prescribed WEGOVY presentation changes, ensure patients and caregivers receive appropriate training and instruct them to consult the Instructions for Use for the newly prescribed presentation.
- Prior to initiation, train patients and their caregiver(s) on proper injection technique for the prescribed WEGOVY presentation *[see Instructions for Use]*. After training, a patient may self-inject WEGOVY if the healthcare provider determines that it can be properly administered, except for the following:
    - WEGOVY FlexTouch is not recommended for self-administration by those who are visually impaired.
- Visually inspect the WEGOVY injection prior to each administration. Only use if the solution is clear, colorless, and contains no particles.
- Administer WEGOVY injection once weekly, on the same day each week, at any time of day, with or without meals.
- Inject WEGOVY subcutaneously in the abdomen, thigh, or upper arm. The time of day and the injection site can be changed without the need for a dosage modification.
- Rotate injection sites with each dose.

WEGOVY Tablets

- Take one WEGOVY tablet orally once daily on an empty stomach in the morning with water (up to 4 ounces). Do not take WEGOVY tablets with other liquids besides water *[see Clinical Pharmacology (12.3)]*.
- Swallow tablets whole. Do not split, crush, chew or dissolve in any solution.
- Do not take more than one tablet per day.
- After taking a WEGOVY tablet, wait at least 30 minutes before eating food, drinking beverages or taking other oral medications *[see Clinical Pharmacology (12.3)]*.

2.2 Recommended Dosage for WEGOVY Injection

Recommended Starting Dosage and Dosage Escalation of WEGOVY Injection for All Approved Indications

- The recommended starting dosage of WEGOVY injection is 0.25 mg administered subcutaneously once weekly.
- Follow the dosage escalation in **Table 1** to reduce the risk of gastrointestinal adverse reactions *[see Warnings and Precautions (5.6), Adverse Reactions (6.1)]*.
- If patients do not tolerate a dose during dosage escalation, consider delaying dosage escalation for 4 weeks.

Table 1. Recommended Starting Dosage and Dosage Escalation of WEGOVY Injection for All Approved Indications in Adults and Pediatric Patients Aged 12 Years and Older

	Weeks	**Once-weekly Subcutaneous Injection Dosage**
Starting Dosage	1 through 4	0.25 mg
Dosage Escalation	5 through 8	0.5 mg
	9 through 12	1 mg
	13 through 16	1.7 mg
Maintenance Dosage	17 and onward	See the indication below for the recommended maintenance dosage(s)

Recommended Maintenance Dosage of WEGOVY Injection for All Approved Indications

Cardiovascular Risk Reduction in Adults

- The maintenance dosage of WEGOVY injection for cardiovascular risk reduction in adults is either 2.4 mg (recommended) or 1.7 mg once weekly.
- Consider treatment response and tolerability when selecting the maintenance dosage *[see Adverse Reactions (6.1), Clinical Studies (14.1)]*.

Weight Reduction in Adults

- The maintenance dosage of WEGOVY for weight reduction in adults is either 1.7 mg or 2.4 mg (recommended) injected subcutaneously once weekly.
- For patients who tolerate the 2.4 mg dosage for at least 4 weeks and additional weight reduction is clinically indicated, the dosage may be increased to a maximum dosage of 7.2 mg subcutaneously once weekly.
- Consider treatment response and tolerability when selecting the maintenance dosage *[see Adverse Reactions (6.1),Clinical Studies (14.2)].*

Weight Reduction in Pediatric Patients Aged 12 Years and Older

- The maintenance dosage of WEGOVY injection for weight reduction in pediatric patients aged 12 years and older is either 2.4 mg (recommended), or 1.7 mg once weekly.
- Consider treatment response and tolerability when selecting the maintenance dosage *[see Adverse Reactions (6.1),Clinical Studies (14.3)]*.

Noncirrhotic MASH with Moderate to Advanced Liver Fibrosis in Adults

- The recommended maintenance dosage of WEGOVY injection for the treatment of noncirrhotic MASH with moderate to advanced liver fibrosis in adults is 2.4 mg injected subcutaneously once weekly. If patients do not tolerate the maintenance dosage of 2.4 mg once weekly, the dosage can be decreased to 1.7 mg once weekly. Consider reescalation to 2.4 mg once weekly *[see Adverse Reactions (6.1), Clinical Studies (14.4)]*.

2.3 Recommended Dosage of WEGOVY Tablets

Recommended Dosage of WEGOVY Tablets for Cardiovascular Risk Reduction or Weight Reduction in Adults

- The recommended starting dosage of WEGOVY tablets is 1.5 mg taken orally once daily. Follow the dosage escalation in **Table 2** to reduce the risk of gastrointestinal adverse reactions *[see Warnings and Precautions (5.6), Adverse Reactions (6.1)]*.
- If patients do not tolerate a dose during dosage escalation, consider delaying dosage escalation.
- The recommended maintenance dosage of WEGOVY tablets is 25 mg orally once daily.
- If patients do not tolerate the 25 mg once daily maintenance dosage, consider switching to WEGOVY injection 1.7 mg once weekly *[see Dosage and Administration (2.5)]*.

Table 2. Recommended Dosage of WEGOVY Tablets for Cardiovascular Risk Reduction or Weight Reduction in Adults

	Days	Once Daily Tablet Dosage
Starting Dosage	1 through 30	1.5 mg
Dosage Escalation	31 through 60	4 mg
	61 through 90	9 mg
Maintenance Dosage	91 and onward	25 mg

2.4 Recommendations Regarding Missed Dose(s)

WEGOVY Injection

- If one dose of WEGOVY injection is missed and the next scheduled dose is:
  - more than 2 days away, administer WEGOVY injection as soon as possible.
  - less than 2 days away, do not administer the WEGOVY injection dose. Resume dosing on the regularly scheduled day of the week.
- If 2 or more consecutive doses of WEGOVY injection are missed, reinitiate dosage escalation at a lower dosage to reduce the risk of gastrointestinal adverse reactions

[see Dosage and Administration (2.2), Warnings and Precautions (5.6), Adverse Reactions (6.1)].

WEGOVY Tablets

If a dose of WEGOVY tablets is missed, skip the missed dose and take the next dose the following day.

2.5 Switching Between WEGOVY Injection and WEGOVY Tablets

Switching from WEGOVY Injection to WEGOVY Tablets

- Patients taking WEGOVY 2.4 mg injection for cardiovascular risk reduction or weight reduction in adults may switch to WEGOVY 25 mg tablets.
- One week after discontinuing WEGOVY 2.4 mg injection, initiate 25 mg of WEGOVY tablets orally once daily.

Switching from WEGOVY Tablets to WEGOVY Injection

- Patients may switch from WEGOVY 25 mg tablets to WEGOVY injection.
- The day after discontinuing WEGOVY tablets 25 mg once daily, initiate WEGOVY 2.4 mg subcutaneous injection once weekly. For patients who do not tolerate WEGOVY 25 mg tablets, consider switching to WEGOVY 1.7 mg injection.
- If additional weight reduction is needed in patients with type 2 diabetes mellitus treated with WEGOVY 25 mg tablets, consider switching to WEGOVY 1.7 mg injection and follow the recommended dosage escalation for WEGOVY injection *[see Dosage and Administration (2.2), Use in Specific Populations (8.6), Clinical Pharmacology (12.3)].*

3 DOSAGE FORMS AND STRENGTHS

Injection: clear, colorless solution available as follows:

Single-Dose Pens	
Dose per Injection	**Total Strength per Total Volume**
0.25 mg	0.25 mg/0.5 mL
0.5 mg	0.5 mg/0.5 mL
1 mg	1 mg/0.5 mL
1.7 mg	1.7 mg/0.75 mL
2.4 mg	2.4 mg/0.75 mL
7.2 mg (WEGOVY HD)	7.2 mg/0.75 mL

Single-Patient-Use Pen (4 doses per Pen) (WEGOVY FlexTouch)	
Dose per Injection	**Total Strength per Total Volume**
2.4 mg	9.6 mg/3 mL (3.2 mg/mL)

Single-Dose Syringes	
Dose per Injection	**Total Strength per Total Volume**
0.25 mg	0.25 mg/0.5 mL
0.5 mg	0.5 mg/0.5 mL
1 mg	1 mg/0.5 mL
1.7 mg	1.7 mg/0.75 mL

2.4 mg	2.4 mg/0.75 mL

Tablets: available as:

Tablet Strength	Description
1.5 mg	White to light yellow, round shaped debossed with “1.5” on one side and “novo” on the other side
4 mg	White to light yellow, round shaped debossed with “4” on one side and “novo” on the other side
9 mg	White to light yellow, round shaped debossed with “9” on one side and “novo” on the other side
25 mg	White to light yellow, oval shaped debossed with “25” on one side and “novo” on the other side

4 CONTRAINDICATIONS

WEGOVY is contraindicated in the following conditions:

- a personal or family history of MTC or in patients with MEN 2 *[see Warnings and Precautions (5.1)]*.
- a prior serious hypersensitivity reaction to semaglutide or to any of the excipients in WEGOVY injection or WEGOVY tablet. Serious hypersensitivity reactions, including anaphylaxis and angioedema, have been reported with WEGOVY *[see Warnings and Precautions (5.7)]*.

5 WARNINGS AND PRECAUTIONS

5.1 Risk of Thyroid C-Cell Tumors

In mice and rats, semaglutide caused a dose-dependent and treatment-duration-dependent increase in the incidence of thyroid C-cell tumors (adenomas and carcinomas) after lifetime exposure at clinically relevant plasma exposures *[see Nonclinical Toxicology (13.1)]*. It is unknown whether WEGOVY causes thyroid C-cell tumors, including MTC, in humans, as human relevance of semaglutide-induced rodent thyroid C-cell tumors has not been determined.

Cases of MTC in patients treated with liraglutide, another GLP-1 receptor agonist, have been reported in the postmarketing period; the data in these reports are insufficient to establish or exclude a causal relationship between MTC and GLP-1 receptor agonist use in humans.

WEGOVY is contraindicated in patients with a personal or family history of MTC or in patients with MEN 2. Counsel patients regarding the potential risk for MTC with the use of WEGOVY and inform them of symptoms of thyroid tumors (e.g., a mass in the neck, dysphagia, dyspnea, persistent hoarseness).

Routine monitoring of serum calcitonin or using thyroid ultrasound is of uncertain value for early detection of MTC in patients treated with WEGOVY. Such monitoring may

increase the risk of unnecessary procedures, due to the low-test specificity for serum calcitonin and a high background incidence of thyroid disease. Significantly elevated serum calcitonin value may indicate MTC and patients with MTC usually have calcitonin values greater than 50 ng/L. If serum calcitonin is measured and found to be elevated, the patient should be further evaluated. Patients with thyroid nodules noted on physical examination or neck imaging should also be further evaluated.

5.2 Acute Pancreatitis

Acute pancreatitis, including fatal and non-fatal hemorrhagic or necrotizing pancreatitis, has been observed in patients treated with GLP-1 receptor agonists, including WEGOVY *[see Adverse Reactions (6)]*. After initiation of WEGOVY, observe patients carefully for signs and symptoms of acute pancreatitis, which may include persistent or severe abdominal pain (sometimes radiating to the back), and which may or may not be accompanied by nausea or vomiting. If pancreatitis is suspected, discontinue WEGOVY and initiate appropriate management.

5.3 Acute Gallbladder Disease

Treatment with WEGOVY is associated with an increased occurrence of cholelithiasis and cholecystitis. The incidence of cholelithiasis and cholecystitis was higher in WEGOVY injection-treated pediatric patients aged 12 years and older than in WEGOVY injection-treated adults. In randomized clinical trials in adults for weight reduction, cholelithiasis was reported by 1.6% of WEGOVY injection-treated patients and 0.7% of placebo injection-treated patients, and by 2.5% of WEGOVY tablet-treated patients and 1% of placebo tablet-treated patients. Cholecystitis was reported by 0.6% of WEGOVY injection-treated adult patients and 0.2% of placebo- treated patients. In a clinical trial in pediatric patients aged 12 years and older for weight reduction, cholelithiasis was reported by 3.8% of WEGOVY injection-treated patients and 0% placebo-treated patients. Cholecystitis was reported by 0.8% of WEGOVY injection-treated pediatric patients and 0% placebo-treated patients *[see Adverse Reactions (6.1)]*.

Substantial or rapid weight loss can increase the risk of cholelithiasis; however, the incidence of acute gallbladder disease was greater in WEGOVY-treated patients than in placebo-treated patients, even after accounting for the degree of weight loss. If cholelithiasis is suspected, gallbladder studies and appropriate clinical follow-up are indicated.

5.4 Hypoglycemia

WEGOVY lowers blood glucose and can cause hypoglycemia.

In a trial of WEGOVY injection in adult patients with type 2 diabetes and body mass index (BMI) greater than or equal to 27 kg/m^2 for weight reduction (Study 3), hypoglycemia (defined as a plasma glucose less than 54 mg/dL) was reported in more patients treated with WEGOVY versus placebo *[see Adverse Reactions (6.1)]*. One episode of severe hypoglycemia (requiring the assistance of another person) was reported in one WEGOVY treated patient versus no placebo-treated patients *[see Clinical Studies (14.2)]*.

In glycemic control clinical trials, the risk of hypoglycemia was increased when semaglutide injection or tablet was used concomitantly with insulin or an insulin secretagogue (e.g., sulfonylurea). Patients with diabetes mellitus taking WEGOVY in combination with insulin or an insulin secretagogue may have an increased risk of hypoglycemia, including severe hypoglycemia. The use of WEGOVY in patients with type

1 diabetes mellitus or in combination with insulin has not been evaluated.

Inform patients of the risk of hypoglycemia and educate them on the signs and symptoms of hypoglycemia. In patients with diabetes, monitor blood glucose prior to starting WEGOVY and during WEGOVY treatment. When initiating WEGOVY, consider reducing the dose of concomitantly administered insulin or insulin secretagogue to reduce the risk of hypoglycemia *[see Drug Interactions (7.1)]*.

5.5 Acute Kidney Injury Due to Volume Depletion

There have been postmarketing reports of acute kidney injury, in some cases requiring hemodialysis, in patients treated with semaglutide *[see Adverse Reactions (6)]*. The majority of the reported events occurred in patients who experienced gastrointestinal adverse reactions leading to dehydration such as nausea, vomiting, or diarrhea *[see Adverse Reactions (6)]*.

Monitor renal function in patients reporting adverse reactions to WEGOVY that could lead to volume depletion, especially during dosage initiation and escalation of WEGOVY.

5.6 Severe Gastrointestinal Adverse Reactions

Use of WEGOVY has been associated with gastrointestinal adverse reactions, sometimes severe *[see Adverse Reactions (6)]*. In clinical trials for adults for weight reduction, severe gastrointestinal adverse reactions were reported more frequently among patients receiving WEGOVY than placebo. Severe gastrointestinal adverse reactions were reported in 4.1% and 0.9% of WEGOVY injection and placebo-treated patients, respectively, and in 2% of WEGOVY tablet-treated and 0% of placebo-treated patients, respectively. Severe gastrointestinal adverse reactions have also been reported postmarketing with GLP-1 receptor agonists.

WEGOVY is not recommended in patients with severe gastroparesis.

5.7 Hypersensitivity Reactions

Serious hypersensitivity reactions (e.g., anaphylaxis, angioedema) have been reported with WEGOVY. If hypersensitivity reactions occur, discontinue use of WEGOVY, treat promptly per standard of care, and monitor until signs and symptoms resolve. WEGOVY is contraindicated in patients with a prior serious hypersensitivity reaction to semaglutide or to any of the excipients in WEGOVY *[see Adverse Reactions (6.2)]*.

Anaphylaxis and angioedema have been reported with other GLP-1 receptor agonists. Use caution in a patient with a history of anaphylaxis or angioedema with another GLP-1 receptor agonist because it is unknown whether such patients will be predisposed to these reactions with WEGOVY.

5.8 Diabetic Retinopathy Complications in Patients with Type 2 Diabetes

In a 2-year trial with semaglutide 0.5 mg and 1 mg once-weekly injection in adult patients with type 2 diabetes and high CV risk, diabetic retinopathy complications (which was a 4-component adjudicated endpoint) occurred in patients treated with semaglutide injection (3%) compared to placebo (1.8%). The absolute risk increase for diabetic retinopathy complications was larger among patients with a history of diabetic retinopathy at baseline (semaglutide injection 8.2%, placebo 5.2%) than among patients without a known history of diabetic retinopathy (semaglutide injection 0.7%, placebo 0.4%).

In a trial of adult patients with type 2 diabetes and BMI greater than or equal to 27 kg/m^2

for weight reduction (Study 3), diabetic retinopathy was reported by 4% of WEGOVY injection-treated patients and 2.7% placebo-treated patients *[see Clinical Studies (14.2)]*.

In a glycemic control trial evaluating a dose comparable to the 9 mg dose and the 25 mg semaglutide tablet dose in patients with type 2 diabetes, a similar proportion of patients in each dose group reported diabetic retinopathy related adverse reactions during the trial; 1.3% and 1.9% of patients in the 9 mg and 25 mg semaglutide group, respectively, reported moderate-severe non-proliferative diabetic retinopathy events, and 0% and 0.4% reported proliferative retinopathy events, respectively.

Rapid improvement in glucose control has been associated with a temporary worsening of diabetic retinopathy. The effect of long-term glycemic control with semaglutide on diabetic retinopathy complications has not been studied. Patients with a history of diabetic retinopathy should be monitored for progression of diabetic retinopathy.

5.9 Heart Rate Increase

Treatment with WEGOVY was associated with increases in resting heart rate. Mean increases in resting heart rate of 1 to 4 beats per minute (bpm) were observed in WEGOVY injection-treated adult patients compared to placebo in clinical trials for weight reduction. More adult patients treated with WEGOVY injection compared with placebo had maximum changes from baseline at any visit of 10 to 19 bpm (41% versus 34%, respectively) and 20 bpm or more (26% versus 16%, respectively). In a clinical trial in pediatric patients aged 12 years and older with normal baseline heart rate, more patients treated with WEGOVY injection compared to placebo had maximum changes in heart rate of 20 bpm or more (54% versus 39%) *[see Adverse Reactions (6.1)]*. Findings were similar in a trial with the WEGOVY tablets.

Monitor heart rate at regular intervals consistent with usual clinical practice. Instruct patients to inform their healthcare providers of palpitations or feelings of a racing heartbeat while at rest during WEGOVY treatment. If patients experience a sustained increase in resting heart rate, discontinue WEGOVY.

5.10 Pulmonary Aspiration During General Anesthesia or Deep Sedation

WEGOVY delays gastric emptying *[see Clinical Pharmacology (12.2)]*. There have been rare postmarketing reports of pulmonary aspiration in patients receiving GLP-1 receptor agonists undergoing elective surgeries or procedures requiring general anesthesia or deep sedation who had residual gastric contents despite reported adherence to preoperative fasting recommendations.

Available data are insufficient to inform recommendations to mitigate the risk of pulmonary aspiration during general anesthesia or deep sedation in patients taking WEGOVY, including whether modifying preoperative fasting recommendations or temporarily discontinuing WEGOVY could reduce the incidence of retained gastric contents. Instruct patients to inform healthcare providers prior to any planned surgeries or procedures if they are taking WEGOVY.

5.11 Never Share WEGOVY FlexTouch Between Patients

WEGOVY FlexTouch must never be shared between patients, even if the needle is changed. Pen-sharing poses a risk for transmission of blood-borne pathogens.

6 ADVERSE REACTIONS

The following serious adverse reactions are described below or elsewhere in the prescribing information:

- Risk of Thyroid C-cell Tumors *[see Warnings and Precautions (5.1)]*
- Acute Pancreatitis *[see Warnings and Precautions (5.2)]*
- Acute Gallbladder Disease *[see Warnings and Precautions (5.3)]*
- Hypoglycemia *[see Warnings and Precautions (5.4)]*
- Acute Kidney Injury Due to Volume Depletion *[see Warnings and Precautions (5.5)]*
- Severe Gastrointestinal Adverse Reactions *[see Warnings and Precautions (5.6)]*
- Hypersensitivity Reactions *[see Warnings and Precautions (5.7)]*
- Diabetic Retinopathy Complications in Patients with Type 2 Diabetes *[see Warnings and Precautions (5.8)]*
- Heart Rate Increase *[see Warnings and Precautions (5.9)]*
- Pulmonary Aspiration During General Anesthesia or Deep Sedation *[see Warnings and Precautions (5.10)]*

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical studies of another drug and may not reflect the rates observed in practice.

Adverse Reactions in Clinical Trials in Adults with Obesity or Overweight for Weight Reduction

WEGOVY 2.4 mg Subcutaneous Injection Weekly Dosage

WEGOVY was evaluated for safety in 3 randomized, double-blind, placebo-controlled trials that included 2,116 adult patients with obesity or overweight treated with 2.4 mg WEGOVY injection for up to 68 weeks and a 7-week off-drug follow-up period *[see Clinical Studies (14.2)]*. Baseline characteristics included a mean age of 48 years, 71% female, 72% White, 14% Asian, 9% Black or African American, and 5% reported as other or unknown; and 85% were not Hispanic or Latino ethnicity, 13% were Hispanic or Latino ethnicity, and 2% reported as unknown. The baseline characteristics were 42% with hypertension, 19% with type 2 diabetes, 43% with dyslipidemia, 28% with a BMI greater than 40 kg/m^2 and 4% with CV disease.

In these clinical trials, 6.8% of patients treated with 2.4 mg WEGOVY injection and 3.2% of patients treated with placebo permanently discontinued treatment as a result of adverse reactions. The most common adverse reactions leading to discontinuation were nausea (1.8% versus 0.2%), vomiting (1.2% versus 0%), and diarrhea (0.7% versus 0.1%) for WEGOVY and placebo, respectively.

Table 3 shows adverse reactions reported in greater than or equal to 2% of WEGOVY 2.4 mg injection-treated adult patients and more frequently than in the placebo group from these trials.

Table 3. Adverse Reactions (≥2% and Greater Than Placebo) in WEGOVY 2.4 mg Injection-treated Adults with Obesity or Overweight for Weight Reduction

	Placebo N=1,261 %	**WEGOVY Injection (2.4 mg Once Weekly) N=2,116**

		%
Nausea	16	44
Diarrhea	16	30
Vomiting	6	24
Constipation	11	24
Abdominal Pain[a]	10	20
Headache	10	14
Fatigue[b]	5	11
Dyspepsia	3	9
Dizziness	4	8
Abdominal Distension	5	7
Eructation	<1	7
Hypoglycemia in T2DM[c]	2	6
Flatulence	4	6
Gastroenteritis	4	6
Gastroesophageal Reflux Disease	3	5
Gastritis[d]	1	4
Gastroenteritis Viral	3	4
Hair Loss	1	3
Dysesthesia[e]	1	2

[a] Includes abdominal pain, abdominal pain upper, abdominal pain lower, gastrointestinal pain, abdominal tenderness, abdominal discomfort and epigastric discomfort.

[b] Includes fatigue and asthenia.

[c] Defined as blood glucose <54 mg/dL with or without symptoms of hypoglycemia or severe hypoglycemia (requiring the assistance of another person) in patients with type 2 diabetes not on concomitant insulin (Study 3, WEGOVY N=403, Placebo N=402). See text below for further information regarding hypoglycemia in patients with and without type 2 diabetes. T2DM = type 2 diabetes mellitus.
[d] Includes chronic gastritis, gastritis, gastritis erosive, and reflux gastritis.
[e] Includes paresthesia, hyperesthesia, burning sensation, allodynia, dysesthesia, skin burning sensation, pain of skin, and sensitive skin.

CV Outcomes Trial

In a CV outcomes trial, 8,803 patients were exposed to WEGOVY injection for a median of 37.3 months and 8,801 patients were exposed to placebo for a median of 38.6 months *[see Clinical Studies (14.1)]*. Safety data collection was limited to serious adverse events (including death), adverse events leading to discontinuation, and adverse events of special interest. Sixteen percent (16%) of WEGOVY injection-treated patients and 8% of placebo-treated patients, respectively, discontinued study drug due to an adverse event. Additional information from this trial is included in subsequent sections below when relevant.

WEGOVY 7.2 mg Subcutaneous Injection Weekly Dosage

The safety of WEGOVY 7.2 mg injection was evaluated in two 72-week double-blinded randomized placebo-controlled 3-armed trials (Study 8 and Study 9), including 1,311 patients with obesity (BMI ≥30 kg/m^2) exposed in the treatment arm with WEGOVY 7.2

mg injection, 304 patients exposed in the treatment arm with WEGOVY 2.4 mg injection, and 303 patients exposed to placebo *[see Clinical Studies (14.2)]*. After dosage escalation, 1,149 patients received at least one dose of WEGOVY 7.2 mg injection, and 578 patients received 7.2 mg for at least 1 year. Baseline characteristics in the WEGOVY 7.2 mg injection arm included a mean age of 49 years, 70% female, 85% White, 5% Asian, 8% Black or African American, and 2% reported as other or unknown; and 5% were Hispanic or Latino ethnicity. At baseline, mean BMI was 39.5 kg/m^2, 52% with hypertension, 38% with dyslipidemia, 40% with a BMI greater than 40 kg/m^2, and 14% with CV disease.

In these clinical trials, 5% of patients treated with 7.2 mg WEGOVY injection, 5% of patients treated with 2.4 mg of WEGOVY injection and 2% of patients treated with placebo permanently discontinued treatment as a result of adverse reactions. The most common adverse reactions leading to discontinuation were gastrointestinal disorders: 3%, 3%, and 0.3%, for WEGOVY 7.2 mg injection, WEGOVY 2.4 mg injection, and placebo, respectively.

Table 4 shows adverse reactions reported in greater than or equal to 2% of WEGOVY 7.2 mg injection and more frequently than 2.4 mg injection-treated and placebo-treated patients from these studies.

Table 4. Adverse Reactions (2% and Greater Than WEGOVY 2.4 mg and Placebo) in WEGOVY 7.2 mg Injection-treated Adults with Obesity for Weight Reduction

	Placebo N=303 %	**WEGOVY Injection (2.4 mg Once Weekly) N=304 %**	**WEGOVY Injection (7.2 mg Once Weekly) N=1,311 %**
Nausea	13	35	39
Vomiting	6	16	22
Dysesthesia[a]	0	6	22
Constipation	8	19	20
Abdominal pain[b]	7	9	12
Fatigue[c]	5	9	11
Headache	7	8	9
Dizziness[d]	1	5	6
Hair loss	1	3	6
Flatulence	2	2	4

[a] Includes allodynia, burning sensation, dysesthesia, hyperesthesia, hyperpathia, pain of skin, paresthesia, sensitive skin, skin burning sensation, skin discomfort, and skin sensitization.

[b] Includes abdominal pain, abdominal pain upper, abdominal pain lower, gastrointestinal pain, abdominal tenderness, abdominal discomfort and epigastric discomfort.

[c] Includes fatigue and asthenia.

[d] Includes dizziness and dizziness postural.

WEGOVY Tablet 25 mg Oral Daily Dosage

WEGOVY tablet was evaluated for safety in a randomized, double-blind, placebo-controlled trial that included 204 adult patients with obesity or overweight and at least one weight-related comorbidity, treated with WEGOVY 25 mg tablets for up to 64 weeks and a 7-week off drug follow-up period (Study 7) *[see Clinical Studies (14.2)]*. Patients with type 2 diabetes mellitus were excluded. Baseline characteristics included a mean age of 48 years, 76% were female, 93% were White, 0.5% were Asian, 6% were Black or African American, and 8% were Hispanic or Latino ethnicity. Mean baseline body weight was 106.4 kg and mean BMI was 37.5 kg/m^2. The baseline characteristics were 44% with hypertension, 31% with dyslipidemia, 27% with a BMI greater than 40 kg/m^2 and 1.5% with coronary artery disease.

In the clinical trial, 6.9% of patients treated with WEGOVY 25 mg tablets and 5.9% patients treated with placebo permanently discontinued treatment as a result of adverse reactions. The most common event type leading to discontinuation was gastrointestinal adverse reactions reported in 3.4% with WEGOVY tablets and 2% with placebo.

In the clinical trial with WEGOVY 25 mg tablets, the types and frequency of common adverse reactions were similar to those listed in **Table 3**.

Adverse Reactions in a Clinical Trial of Pediatric Patients Aged 12 Years and Older with Obesity Treated with WEGOVY 2.4 mg Injection for Weight Reduction

WEGOVY injection was evaluated in a 68-week, double-blind, randomized, parallel group, placebo-controlled, multi-center trial in 201 pediatric patients aged 12 years and older with obesity *[see Clinical Studies (14.3)]*. Baseline characteristics included a mean age of 15.4 years; 38% of patients were male; 79% were White, 8% were Black or African American, 2% were Asian, and 11% were of other or unknown race; and 11% were of Hispanic or Latino ethnicity. The mean baseline body weight was 107.5 kg, and mean BMI was 37 kg/m^2.

Table 5 shows adverse reactions reported in greater than or equal to 3% of WEGOVY injection-treated pediatric patients and more frequently than in the placebo group from a trial in pediatric patients aged 12 years and older.

Table 5. Adverse Reactions (≥3% and Greater than Placebo) in WEGOVY 2.4 mg Injection-treated Pediatric Patients Aged 12 Years and Older with Obesity for Weight Reduction

	Placebo N=67 %	**WEGOVY Injection (2.4 mg Once Weekly) N=133 %**
Nausea	18	42
Vomiting	10	36
Diarrhea	19	22
Headache	16	17
Abdominal Pain	6	15
Nasopharyngitis	10	12
Dizziness	3	8
Gastroenteritis	3	7
Constipation	2	6
Gastroesophageal Reflux		

Gastroesophageal Reflux Disease	2	4
Sinusitis	2	4
Urinary tract infection	2	4
Ligament sprain	2	4
Anxiety	2	4
Hair Loss	0	4
Cholelithiasis	0	4
Eructation	0	4
Influenza	0	3
Rash	0	3
Urticaria	0	3

Adverse Reactions in Clinical Trials in Adults with MASH Treated with WEGOVY Injection

The safety of WEGOVY injection was evaluated in a randomized, double-blind, placebo-controlled trial (Study 11) that included 1,195 adult patients with MASH, including 800 patients who were exposed to WEGOVY for a median of 95.3 weeks and 395 patients who were exposed to placebo for a median of 83.1 weeks *[see Clinical Studies (14.4)].*

The most commonly reported adverse reactions were consistent with the other approved WEGOVY indications (see **Table 3**). There is limited information in patients with MASH and a BMI <25 kg/m^2. Additional information from the MASH trial is included in subsequent sections when notable. Unless indicated, the incidence of the adverse reactions in MASH patients was similar to other approved indications.

Other Adverse Reactions in Adults and/or Pediatric Patients Treated with WEGOVY Injection or WEGOVY Tablets

The safety of WEGOVY tablets and WEGOVY 7.2 mg injection have not been established in pediatric patients or patients with MASH.

Acute Gallbladder Disease

In WEGOVY clinical trials in adults for weight reduction, cholelithiasis was reported by 1.6% of WEGOVY injection-treated patients and 0.7% of placebo-treated patients and by 2.5% of WEGOVY tablet-treated patients and 1% of placebo tablet-treated patients. Cholecystitis was reported by 0.6% of WEGOVY injection-treated adult patients and 0.2% of placebo-treated patients. In a clinical trial in pediatric patients aged 12 years and older for weight reduction *[see Clinical Studies (14.3)]*, cholelithiasis was reported by 3.8% of WEGOVY injection-treated patients and 0% placebo-treated patients. Cholecystitis was reported by 0.8% of WEGOVY injection-treated pediatric patients and 0% placebo-treated patients.

Hypoglycemia

Patients with Type 2 Diabetes

In a trial of adult patients with type 2 diabetes and BMI greater than or equal to 27 kg/m^2 for weight reduction, clinically significant hypoglycemia (defined as a plasma glucose less than 54 mg/dL) was reported in 6.2% of WEGOVY injection-treated patients versus 2.5% of placebo-treated patients. A higher rate of clinically significant hypoglycemic episodes was reported with WEGOVY injection (semaglutide 2.4 mg) versus semaglutide 1 mg injection (10.7 vs. 7.2 episodes per 100 patient years of exposure, respectively); the rate in the placebo-treated group was 3.2 episodes per 100 patient years of exposure.

In addition, one episode of severe hypoglycemia requiring intravenous glucose was reported in a WEGOVY injection-treated patient versus none in placebo-treated patients. The risk of hypoglycemia was increased when WEGOVY was used with a sulfonylurea.

In a glycemic control trial evaluating a dose comparable to the 9 mg and 25 mg semaglutide tablet dose in patients with type 2 diabetes not on insulin, clinically significant hypoglycemia was reported in similar proportions of subjects in both treatment groups.

Patients without Type 2 Diabetes

Episodes of hypoglycemia have been reported with GLP-1 receptor agonists in adult patients without type 2 diabetes mellitus. In WEGOVY clinical trials in adult patients without type 2 diabetes mellitus for weight reduction, there was no systematic capturing or reporting of hypoglycemia.

In a CV outcomes trial in adult patients without type 2 diabetes, 3 episodes of serious hypoglycemia were reported in WEGOVY injection-treated patients versus 1 episode in placebo. Patients with a history of bariatric surgery (a risk factor for hypoglycemia) had more events of serious hypoglycemia while taking WEGOVY injection (2.3%, 2/87) than placebo (0%, 0/97).

Retinal Disorders in Patients with Type 2 Diabetes

In a trial of adult patients with type 2 diabetes and BMI greater than or equal to 27 kg/m^2 for weight reduction, retinal disorders were reported by 6.9% of patients treated with WEGOVY injection (semaglutide 2.4 mg), 6.2% of patients treated with semaglutide 1 mg, and 4.2% of patients treated with placebo. The majority of events were reported as diabetic retinopathy (4%, 2.7%, and 2.7%, respectively) and non-proliferative retinopathy (0.7%, 0%, and 0%, respectively).

In a glycemic control trial evaluating the 9 mg and 25 mg semaglutide tablet doses in patients with type 2 diabetes, a similar proportion of patients in each dose group reported diabetic retinopathy related adverse reactions during the trial; 1.3% and 1.9% of patients in the 9 mg and 25 mg semaglutide group, respectively, reported moderate-severe non-proliferative diabetic retinopathy events, and 0% and 0.4% reported proliferative retinopathy events, respectively.

In a 2-year trial with semaglutide 0.5 mg and 1 mg once-weekly injection in adult patients with type 2 diabetes and high cardiovascular risk, diabetic retinopathy complications (which was a 4-component adjudicated endpoint) occurred in patients treated with semaglutide injection (3%) compared to placebo (1.8%). The absolute risk increase for diabetic retinopathy complications was larger among patients with a history of diabetic retinopathy at baseline (semaglutide injection 8.2%, placebo 5.2%) than among patients without a known history of diabetic retinopathy (semaglutide injection 0.7%, placebo 0.4%).

Gastrointestinal Adverse Reactions

In clinical trials in adults for weight reduction, 73% of WEGOVY injection-treated patients and 47% of patients receiving placebo reported gastrointestinal adverse reactions, including severe reactions that were reported more frequently among patients receiving WEGOVY injection (4.1%) than placebo (0.9%). The most frequently reported reactions were nausea (44% vs. 16%), vomiting (25% vs. 6%), and diarrhea (30% vs. 16%). Other reactions that occurred at a higher incidence among WEGOVY injection-treated adult patients included dyspepsia, abdominal pain, abdominal distension, eructation, flatulence,

gastroesophageal reflux disease, gastritis, hemorrhoids, and hiccups. These reactions were most frequently reported during dosage escalation.

Severe gastrointestinal adverse reactions were reported in 2% of WEGOVY tablet-treated and 0% of placebo-treated patients, respectively.

In the pediatric clinical trial for weight reduction, 62% of WEGOVY injection-treated patients and 42% of placebo-treated patients reported gastrointestinal adverse reactions. The most frequently reported reactions were nausea (42% vs. 18%), vomiting (36% vs. 10%), and diarrhea (22% vs. 19%). Other gastrointestinal-related reactions that occurred at a higher incidence than placebo among WEGOVY injection-treated pediatric patients included abdominal pain, constipation, eructation, gastroesophageal reflux disease, dyspepsia, and flatulence.

Permanent discontinuation of treatment as a result of a gastrointestinal adverse reaction occurred in 4.3% of WEGOVY injection-treated adult patients versus 0.7% of placebo-treated patients. In a pediatric clinical trial for weight reduction, 2.3% of patients treated with WEGOVY injection versus 1.5% of patients who received placebo discontinued treatment as a result of gastrointestinal adverse reactions.

Dysesthesia

Dysesthesia, and related events of altered skin sensations including paresthesia, pain of skin, sensitive skin, and burning skin sensation, occurred in clinical trials with WEGOVY injection and tablet. The incidence of dysesthesia with WEGOVY increased with increasing dosage and drug levels in the blood.

In clinical trials evaluating patients with and without type 2 diabetes mellitus, WEGOVY 7.2 mg injection dysesthesia was reported at a higher rate in the WEGOVY 7.2 mg injection group (22%), compared to WEGOVY 2.4 mg injection (6%) and placebo (0.3%). Among 288 patients who experienced dysesthesia with WEGOVY 7.2 mg injection, 2% permanently discontinued treatment, 8% had temporary interruption, and 23% had dose reduction; most subjects with these actions taken to WEGOVY 7.2 mg injection (permanent discontinuation, temporary interruption, or dose reduction) recovered from the event. When an action was taken with WEGOVY, events resolved faster than the events where no action was taken with WEGOVY. Among patients who experienced dysesthesia with WEGOVY 7.2 mg injection, 18% did not report recovering during the trial duration; in most of the patients who did not recover, the WEGOVY dosage was unchanged and not discontinued. Of the patients who recovered after action taken with WEGOVY, 38 patients were re-escalated to 7.2 mg. Of these 38 patients, 17 (45%) subsequently reported recurrence of dysesthesia.

In a clinical trial with WEGOVY 25 mg tablets, 5% of WEGOVY-treated patients and no placebo-treated patients reported dysesthesia adverse reactions.

Other Adverse Reactions in Adults and/or Pediatric Patients

The below adverse reactions were reported in WEGOVY injection clinical trials and are applicable to both WEGOVY injection and WEGOVY tablets. The safety of WEGOVY tablets have not been established in pediatric patients or patients with MASH.

Acute Pancreatitis

In WEGOVY clinical trials in adults for weight reduction, acute pancreatitis was confirmed by adjudication in 4 WEGOVY-treated patients (0.2 cases per 100 patient years) and 1 in placebo-treated patients (less than 0.1 cases per 100 patient years). One additional case of acute pancreatitis was confirmed in a patient treated with WEGOVY in another clinical

trial.

Acute Kidney Injury

Acute kidney injury occurred in clinical trials for weight reduction in 7 adult patients (0.4 cases per 100 patient years) receiving WEGOVY versus 4 patients (0.2 cases per 100 patient years of exposure) receiving placebo. Some of these adverse reactions occurred in association with gastrointestinal adverse reactions or dehydration. In addition, 2 patients treated with WEGOVY had acute kidney injury with dehydration in other clinical trials. The risk of renal adverse reactions with WEGOVY was increased in adult patients with a history of renal impairment (trials included 65 patients with a history of moderate or severe renal impairment at baseline), and occurred more frequently during dose titration.

Increase in Heart Rate

Mean increases in resting heart rate of 1 to 4 beats per minute (bpm) were observed with routine clinical monitoring in WEGOVY-treated adult patients compared to placebo in clinical trials for weight reduction. In weight reduction trials in which adult patients were randomized prior to dose-escalation, more patients treated with WEGOVY, compared with placebo, had maximum changes from baseline at any visit of 10 to 19 bpm (41% versus 34%, respectively) and 20 bpm or more (26% versus 16%, respectively). In a clinical trial for weight reduction in pediatric patients aged 12 years and older with normal baseline heart rate, more patients treated with WEGOVY compared to placebo had maximum changes in heart rate of 20 bpm or more (54% versus 39%).

Hypotension and Syncope

Adverse reactions related to hypotension (hypotension, orthostatic hypotension, and decreased blood pressure) were reported in 1.3% of WEGOVY-treated adult patients versus 0.4% of placebo-treated patients and syncope was reported in 0.8% of WEGOVY-treated patients versus 0.2% of placebo-treated patients in clinical trials for weight reduction. Some reactions were related to gastrointestinal adverse reactions and volume loss associated with WEGOVY. Hypotension and orthostatic hypotension were more frequently seen in patients on concomitant antihypertensive therapy. In a clinical trial in pediatric patients aged 12 years and older for weight reduction, hypotension was reported in 2.3% of WEGOVY-treated patients versus 0% in placebo-treated patients.

Appendicitis

Appendicitis (including perforated appendicitis) occurred in 10 (0.5%) WEGOVY-treated adult patients and 2 (0.2%) patients receiving placebo in clinical trials for weight reduction.

Injection Site Reactions

In clinical trials in adults for weight reduction, 1.4% of WEGOVY-treated patients and 1% of patients receiving placebo experienced injection site reactions (including injection site pruritus, erythema, inflammation, induration, and irritation).

Hypersensitivity Reactions

Serious hypersensitivity reactions (e.g., anaphylaxis, angioedema) have been reported with WEGOVY.

In a pediatric clinical trial for weight reduction, rash was reported in 3% of WEGOVY-treated patients and 0% of placebo-treated patients, and urticaria was reported in 3% of WEGOVY-treated patients and 0% of placebo-treated patients.

In adult clinical trials for weight reduction, allergic reactions occurred in 16% (8/50) of WEGOVY-treated patients with anti-semaglutide antibodies and in 7% (114/1659) of WEGOVY-treated patients who did not develop anti-semaglutide antibodies *[see Clinical Pharmacology (12.6)].*

Fractures

In the CV outcomes trial in adults, more fractures of the hip and pelvis were reported on WEGOVY than on placebo in female patients: 1% (24/2,448) vs. 0.2% (5/2,424), and in patients ages 75 years and older: 2.4% (17/703) vs. 0.6% (4/663), respectively.

In a clinical trial in adults with MASH, fractures occurred in 4.4% of WEGOVY-treated patients (2.6 cases per 100 patient years) compared to 3.3% of placebo-treated patients (2 cases per 100 patient years). Fractures were reported in both males and females with a median age of 61 years (range, 44 to 75).

Urolithiasis

In a CV outcomes trial, 1.2% of WEGOVY-treated patients and 0.8% of patients receiving placebo reported urolithiasis, including serious reactions that were reported more frequently among patients receiving WEGOVY (0.6%) than placebo (0.4%).

Dysgeusia

In clinical trials in adults for weight reduction, 1.7% of WEGOVY-treated patients and 0.5% of placebo-treated patients reported dysgeusia.

Hair Loss

Hair loss adverse reactions in WEGOVY injection-treated patients were associated with weight reduction. In a pool of studies 2, 3, and 4, hair loss was reported in 3.3% of patients treated with WEGOVY 2.4 mg (4% female, 0.9% male) and in 1% of patients treated with placebo (2% female, 0 male). In a pediatric trial, hair loss was reported in 4% of patients treated with 2.4 mg of WEGOVY and no placebo-treated patients.

In clinical trials with WEGOVY 7.2 mg injection, hair loss was reported in 5.8% of patients treated with WEGOVY 7.2 mg (8.4% female, 0.2% male), 3.3% of patients treated with WEGOVY 2.4 mg (5.4% female, 0 male), and 1.0% of patients on placebo (1.5% female, 0 male). In the WEGOVY 7.2 mg group, 1 event led to permanent treatment discontinuation, 1 event led to temporary interruption, and 5 events led to dose reduction, versus none of these actions in the WEGOVY 2.4 mg or placebo groups.

Laboratory Abnormalities in Adults and/or Pediatric Patients Treated with WEGOVY Injection

The below laboratory abnormalities are applicable for both WEGOVY injection and WEGOVY tablets and include descriptions of WEGOVY injection clinical trial data, where relevant. The safety and effectiveness of WEGOVY tablets have not been established in pediatric patients or patients with MASH.

Amylase and Lipase

Adult and pediatric patients treated with WEGOVY had a mean increase from baseline in amylase of 15% to 16% and lipase of 39% in clinical trials for weight reduction. These changes were not observed in the placebo group.

In a clinical trial in adults with MASH, increases in lipase greater than 3 times the upper limit of normal (ULN) occurred in 4.7% (35/750) of WEGOVY-treated patients compared

with 1.3% (5/374) of placebo-treated patients. The clinical significance of elevations in lipase or amylase with WEGOVY is unknown in the absence of other signs and symptoms of pancreatitis.

Liver Tests

In a pediatric clinical trial for weight reduction, increases in alanine aminotransferase (ALT) greater than or equal to 5 times the ULN were observed in 4 (3%) WEGOVY-treated patients compared with 0% of placebo-treated patients. In some patients, increases in ALT and AST were associated with other confounding factors (such as gallstones). In the CV outcomes trial in adults, increases in total bilirubin greater than or equal to 3 times the ULN were observed in 0.3% (30/8,585) of WEGOVY-treated patients versus 0.2% (14/8,579) of placebo-treated patients.

6.2 Postmarketing Experience

The following adverse reactions have been reported during post-approval use of semaglutide, the active ingredient of WEGOVY. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Gastrointestinal Disorders: acute pancreatitis and necrotizing pancreatitis, sometimes resulting in death; ileus, intestinal obstruction, severe constipation including fecal impaction

Hypersensitivity: anaphylaxis, angioedema, rash, urticaria

Pulmonary: Pulmonary aspiration has occurred in patients receiving GLP-1 receptor agonists undergoing elective surgeries or procedures requiring general anesthesia or deep sedation

Renal and Urinary Disorders: acute kidney injury

7 DRUG INTERACTIONS

7.1 Concomitant Use with Insulin or an Insulin Secretagogue (e.g., Sulfonylurea)

WEGOVY lowers blood glucose and can cause hypoglycemia. The risk of hypoglycemia is increased when WEGOVY is used concomitantly with insulin or insulin secretagogues (e.g., sulfonylureas). The addition of WEGOVY in patients treated with insulin has not been evaluated.

When initiating WEGOVY, consider reducing the dose of concomitantly administered insulin secretagogue or insulin to reduce the risk of hypoglycemia *[see Warnings and Precautions (5.4), Adverse Reactions (6.1)]*.

7.2 Oral Medications

WEGOVY causes a delay of gastric emptying and thereby has the potential to impact the absorption of concomitantly administered oral medications. In clinical pharmacology trials with semaglutide 1 mg once weekly injection, semaglutide did not affect the absorption of orally administered medications *[see Clinical Pharmacology (12.3)]*. In a drug interaction study with the semaglutide tablet, levothyroxine exposure was increased 33% (90% CI: 1.25-1.42). Monitor the effects of oral medications concomitantly administered with WEGOVY. Consider increased clinical or laboratory

monitoring for medications that have a narrow therapeutic index or that require clinical monitoring.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Exposure Registry

There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to WEGOVY during pregnancy. Pregnant women exposed to WEGOVY and healthcare providers are encouraged to contact Novo Nordisk at 1-877-390-2760 or www.wegovypregnancyregistry.com.

Risk Summary

Based on animal reproduction studies, there may be potential risks to the fetus from exposure to semaglutide during pregnancy. Available pharmacovigilance data and data from clinical trials with WEGOVY use in pregnant patients are insufficient to establish a drug-associated risk of major birth defects, miscarriage or adverse maternal or fetal outcomes.

Weight loss offers no benefit to a pregnant patient and may cause fetal harm. When a pregnancy is recognized, advise the pregnant patient of the risk to a fetus. Discontinue WEGOVY in pregnant patients who are using it for weight reduction (see *Clinical Considerations*).

There may be risks to the mother and fetus related to underlying MASH with advanced liver fibrosis (see *Clinical Considerations*). Whether WEGOVY treatment during pregnancy reduces these risks is unknown. WEGOVY for the treatment of MASH with advanced liver fibrosis should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

In pregnant rats administered semaglutide during organogenesis, embryofetal mortality, structural abnormalities and alterations to growth occurred at clinically relevant maternal exposures at the maximum recommended human dose (MRHD) of WEGOVY subcutaneous injection 7.2 mg/week, based on AUC. In pregnant rabbits and cynomolgus monkeys administered semaglutide during organogenesis, early pregnancy losses and structural abnormalities were observed in rabbits and monkeys at clinically relevant exposures. These findings coincided with a marked maternal body weight loss in both animal species (see *Data*).

The background risk of major birth defects and miscarriage for the indicated populations are unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

Clinical Considerations

Disease-Associated Maternal and/or Embryo/fetal Risk: Appropriate weight gain based on pre-pregnancy weight is currently recommended for all pregnant patients, including those who already have overweight or obesity, because of the obligatory weight gain that occurs in maternal tissues during pregnancy.

There may be risks to the mother and fetus related to MASH with advanced liver fibrosis, such as increased risks of gestational diabetes, hypertensive complications, preterm

birth, and postpartum hemorrhage. The effect of WEGOVY on these risks is unknown.

Data

Animal Data: In a combined fertility and embryofetal development study in male and female rats, subcutaneous semaglutide doses of 0.01, 0.03, and 0.09 mg/kg/day (up to 0.1-fold the MRHD, based on AUC) were administered to male rats for 4 weeks prior to and throughout mating and to female rats for 2 weeks prior to mating, and throughout organogenesis to Gestation Day 17. In parental rats, pharmacologically mediated reductions in body weight gain and food consumption were observed at all dose levels. In the offspring, reduced growth and fetuses with visceral (heart blood vessels) and skeletal (cranial bones, vertebra, ribs) abnormalities were observed at clinically relevant exposures at the MRHD.

In an embryofetal development study in pregnant rabbits, subcutaneous semaglutide doses of 0.001, 0.0025, or 0.0075 mg/kg/day (up to 0.3-fold the MRHD) were administered throughout organogenesis from Gestation Day 6 to 19. Pharmacologically mediated reductions in maternal body weight gain and food consumption were observed at all dose levels. Early pregnancy losses and increased incidences of minor visceral (kidney, liver) and skeletal (sternebra) fetal abnormalities were observed at greater than or equal to 0.0025 mg/kg/day, at clinically relevant exposures at the MRHD.

In an embryofetal development study in pregnant cynomolgus monkeys, subcutaneous semaglutide doses of 0.015, 0.075, and 0.15 mg/kg twice weekly (up to 2-fold the MRHD) were administered throughout organogenesis, from Gestation Day 16 to 50. Pharmacologically mediated, marked initial maternal body weight loss and reductions in body weight gain and food consumption coincided with the occurrence of sporadic abnormalities (vertebra, sternebra, ribs) at greater than or equal to 0.075 mg/kg twice weekly, at clinically relevant exposures at the MRHD.

In a pre- and postnatal development study in pregnant cynomolgus monkeys, subcutaneous semaglutide doses of 0.015, 0.075, and 0.15 mg/kg twice weekly (up to 0.8-fold the MRHD) were administered from Gestation Day 16 to 140. Pharmacologically mediated marked initial maternal body weight loss and reductions in body weight gain and food consumption coincided with an increase in early pregnancy losses and led to delivery of slightly smaller offspring at greater than or equal to 0.075 mg/kg twice weekly, at clinically relevant exposures at the MRHD.

Salcaprozate sodium (SNAC), an absorption enhancer in WEGOVY tablets, crosses the placenta and reaches fetal tissues in rats. In a pre- and postnatal development study in pregnant Sprague Dawley rats, SNAC was administered orally at 1,000 mg/kg/day (exposure levels were not measured) on Gestation Day 7 through lactation Day 20. An increase in gestation length, an increase in the number of stillbirths and a decrease in pup viability were observed.

8.2 Lactation

Risk Summary

WEGOVY Oral Tablets

Data from a clinical lactation study with semaglutide oral tablet formulation reported semaglutide concentrations below the lower limit of quantification in human milk. However, SNAC and/or its metabolites are present in human milk. Since the activity of enzymes involved in SNAC clearance may be lower in infants compared to adults, higher SNAC plasma levels may occur in neonates and infants. Because of the unknown

potential for serious adverse reactions in the breastfed infant due to the possible accumulation of SNAC, and because there are alternative formulations of semaglutide that do not contain SNAC that can be used during lactation, advise patients that breastfeeding is not recommended during treatment with WEGOVY tablets.

WEGOVY Subcutaneous Injection

There are no data on the presence of subcutaneously administered semaglutide or its metabolites in human milk, the effects on the breastfed infant, or the effects on milk production. WEGOVY subcutaneous injection does not contain the SNAC metabolites.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for WEGOVY and any potential adverse effects on the breastfed infant from WEGOVY or from the underlying maternal condition.

8.3 Females and Males of Reproductive Potential

Because of the potential for fetal harm, discontinue WEGOVY in patients at least 2 months before they plan to become pregnant to account for the long half-life of semaglutide *[see Use in Specific Populations (8.1)]*.

8.4 Pediatric Use

The safety and effectiveness of WEGOVY injection in combination with a reduced-calorie diet and increased physical activity to reduce excess body weight and maintain weight reduction long term in pediatric patients aged 12 years and older with obesity have been established. Use of WEGOVY injection for this indication is supported by a 68-week, double-blind, placebo-controlled clinical trial in 201 pediatric patients aged 12 years and older with a BMI corresponding to ≥95th percentile for age and sex *[see Clinical Studies (14.3)]* and from trials in adult patients with obesity *[see Clinical Studies (14.2)]*. Use of the 1.7 mg once weekly maintenance dosage of WEGOVY injection in pediatric patients is also supported by additional exposure-efficacy and safety analyses in pooled adult and pediatric patients.

Adverse reactions with WEGOVY injection treatment in pediatric patients aged 12 years and older were generally similar to those reported in adults. Pediatric patients aged 12 years and older treated with WEGOVY injection had greater incidences of cholelithiasis, cholecystitis, hypotension, rash, and urticaria compared to adults treated with WEGOVY *[see Adverse Reactions (6.1)]*.

Although there was an increased frequency of hypoglycemia in adults with type 2 diabetes with obesity or overweight treated with WEGOVY injection, there are insufficient data to determine if the risk of hypoglycemia is higher in WEGOVY-treated pediatric patients with type 2 diabetes with obesity. Inform pediatric patients of the risk of hypoglycemia and educate them on the signs and symptoms of hypoglycemia. In pediatric patients aged 12 years and older with type 2 diabetes, monitor blood glucose prior to starting WEGOVY injection and during treatment. When initiating WEGOVY injection in pediatric patients aged 12 years and older with type 2 diabetes, consider reducing the dosage of concomitantly administered insulin secretagogue (such as sulfonylureas) or insulin to reduce the risk of hypoglycemia *[see Warnings and Precautions (5.4)]*.

The safety and effectiveness of WEGOVY injection have not been established in pediatric patients:

- to reduce the risk of major adverse CV events. Clinical trials for this indication are

highly impracticable because of the low prevalence of the condition in pediatric patients.
- to reduce excess body weight and maintain weight reduction long term with the 7.2 mg once weekly dosage.
- to reduce excess body weight and maintain weight reduction long term in those less than 12 years of age.
- for the treatment of noncirrhotic MASH.

The safety and effectiveness of WEGOVY tablets have not been established in pediatric patients.

8.5 Geriatric Use

In the WEGOVY injection clinical trials for weight reduction and long-term maintenance, 233 (9%) WEGOVY injection-treated patients were aged 65 to less than 75 years and 23 (1%) WEGOVY injection-treated patients were aged 75 years and older *[see Clinical Studies (14.2)]*. In a WEGOVY 25 mg tablets clinical trial for weight reduction and long-term maintenance, 16 (8%) WEGOVY tablets-treated patients were aged 65 to less than 75 years and 5 (2%) WEGOVY tablet-treated patients were aged 75 years and older *[see Clinical Studies (14.2)]*.

In a CV outcomes trial, 2,656 (30%) WEGOVY injection-treated patients were aged 65 to 75 years and 703 (8%) WEGOVY injection-treated patients were aged 75 years and older *[see Clinical Studies (14.1)]*. No overall difference in the effectiveness was observed between patients aged 65 years and older and younger adult patients. In the CV outcomes trial, patients aged 75 years and older reported more hip and pelvis fractures in the WEGOVY injection-treated patients than placebo-treated patients. Patients aged 75 years and older (WEGOVY injection-treated and placebo-treated) reported more serious adverse reactions overall compared to younger adult patients *[see Adverse Reactions (6.1)]*.

In the clinical trial in patients with MASH, of the 534 patients randomized to WEGOVY injection, 138 (26%) were aged 65 years and older and 13 (2%) were aged 75 years and older *[see Clinical Studies (14.4)]*. No overall differences in safety or effectiveness of WEGOVY injection have been observed between patients 65 years of age and older and younger adult patients with MASH.

8.6 Type 2 Diabetes Mellitus

WEGOVY tablets have not been studied for weight reduction in adults with type 2 diabetes and obesity or overweight *[see Clinical Studies (14.2)]*.

Administration of WEGOVY injection resulted in less weight reduction in patients with type 2 diabetes and obesity or overweight compared to those without type 2 diabetes and obesity or overweight *[see Clinical Studies (14.2)]*.

In patients without type 2 diabetes, the average semaglutide blood concentrations following administration of WEGOVY 25 mg tablets once daily or WEGOVY 2.4 mg injection once weekly were similar. In patients with type 2 diabetes, the average semaglutide blood concentrations following administration of WEGOVY 25 mg tablets once daily were lower than the average semaglutide blood concentrations following use of WEGOVY 2.4 mg injection once weekly *[see Clinical Pharmacology (12.3)]*. The lower semaglutide blood concentrations after administration of WEGOVY tablets in patients with type 2 diabetes may be associated with reductions in absolute bioavailability.

Given the differences in average semaglutide blood concentrations after administration of WEGOVY tablets and WEGOVY injection, and the higher variability in semaglutide blood concentrations after administration of WEGOVY tablets in all patient populations, some patients with type 2 diabetes taking WEGOVY tablets may experience subtherapeutic semaglutide blood concentrations. In patients with type 2 diabetes with an inadequate response to WEGOVY tablets, consider alternative therapies, including switching to WEGOVY injection *[see Dosage and Administration (2.5)]*.

10 OVERDOSAGE

Overdoses have been reported with other GLP-1 receptor agonists. Effects have included severe nausea, severe vomiting, and severe hypoglycemia. In the event of overdose, appropriate supportive treatment should be initiated according to the patient's clinical signs and symptoms. In the event of an overdose of WEGOVY, consider contacting the Poison Help line (1-800-222-1222) or a medical toxicologist for additional overdosage management recommendations. A prolonged period of observation and treatment for these symptoms may be necessary, taking into account the long half-life of WEGOVY of approximately 1 week.

11 DESCRIPTION

WEGOVY contains semaglutide, a human GLP-1 receptor agonist (or GLP-1 analog). The peptide backbone is produced by yeast fermentation. The main protraction mechanism of semaglutide is albumin binding, facilitated by modification of position 26 lysine with a hydrophilic spacer and a C18 fatty di-acid. Furthermore, semaglutide is modified in position 8 to provide stabilization against degradation by the enzyme dipeptidyl-peptidase 4 (DPP-4). A minor modification was made in position 34 to ensure the attachment of only one fatty di-acid. The molecular formula is $C_{187}H_{291}N_{45}O_{59}$ and the molecular weight is 4113.58 g/mol.

Figure 1. Structural Formula of Semaglutide



WEGOVY injection is a sterile, aqueous, clear, colorless solution. Each 0.5 mL single-dose pen contains a solution of WEGOVY containing 0.25 mg, 0.5 mg or 1 mg of semaglutide; and each 0.75 mL single-dose pen contains a solution of WEGOVY containing 1.7, 2.4 mg or 7.2 mg of semaglutide. Each 0.5 mL single-dose prefilled syringe contains a solution of WEGOVY containing 0.25 mg, 0.5 mg or 1 mg of semaglutide; and each 0.75

mL single-dose prefilled syringe contains a solution of WEGOVY containing 1.7, 2.4 mg of semaglutide. Each 1 mL of WEGOVY contains the following inactive ingredients: disodium phosphate dihydrate, 1.42 mg; sodium chloride, 8.25 mg; and water for injection. WEGOVY has a pH of approximately 7.4. Hydrochloric acid or sodium hydroxide may be added to adjust pH.

Each 3 mL prefilled single-patient-use pen contains semaglutide 9.6 mg (3.2 mg/mL), which provides 4 doses of 2.4 mg semaglutide per 0.75 mL. Each 1 mL of WEGOVY FlexTouch solution also contains the following inactive ingredients: disodium phosphate dihydrate, 1.42 mg; phenol, 5.5 mg; propylene glycol, 14 mg; and water for injections. WEGOVY has a pH of approximately 7.4. Hydrochloric acid or sodium hydroxide may be added to adjust pH.

Each single-patient-use WEGOVY FlexTouch Pen contains additional volume to allow for device priming.

WEGOVY tablets include semaglutide as a white to almost white hygroscopic powder. Each tablet of WEGOVY contains 1.5 mg, 4 mg, 9 mg or 25 mg of semaglutide and the following inactive ingredients: magnesium stearate and salcaprozate sodium (SNAC).

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Semaglutide is a GLP-1 analogue with 94% sequence homology to human GLP-1. Semaglutide acts as a GLP-1 receptor agonist that selectively binds to and activates the GLP-1 receptor, the target for native GLP-1.

GLP-1 is a physiological regulator of appetite and caloric intake, and the GLP-1 receptor is present in several areas of the brain involved in appetite regulation. Animal studies show that semaglutide distributed to and activated neurons in brain regions involved in regulation of food intake.

The exact mechanism of semaglutide in CV risk reduction in adults has not been established.

For treatment of MASH in humans, the precise mechanism of action of semaglutide is not fully understood and may involve multiple pathways mediated by weight loss and other factors. In a mouse model of diet-induced MASH, treatment with semaglutide resulted in histological improvements in steatosis, inflammation, and fibrosis in liver compared to baseline, which was associated with body weight loss, intermittent periods of reduced food intake, and improvements in relevant biomarkers. The relationship between the pathophysiology of MASH in animal models and humans has not been fully established.

12.2 Pharmacodynamics

Semaglutide lowers body weight with greater fat mass loss than lean mass loss. Semaglutide decreases calorie intake. The effects are likely mediated by affecting appetite.

Semaglutide stimulates insulin secretion and reduces glucagon secretion in a glucose-dependent manner. These effects can lead to a reduction of blood glucose.

Gastric Emptying

Semaglutide delays gastric emptying.

Cardiac Electrophysiology (QTc)

The effect of semaglutide on cardiac repolarization was tested in a thorough QTc trial. Semaglutide did not prolong QTc intervals at subcutaneous doses up to 1.5 mg at steady state.

Noninvasive Liver Disease Markers

Semaglutide decreases liver fat content measured by Magnetic Resonance Imaging Proton Density Fat Fraction (MRI-PDFF), liver stiffness assessed by transient elastography (TE), Enhanced Liver Fibrosis (ELF) score, and the levels of the pro-peptide of type III collagen biomarker (Pro-C3). The clinical relevance of these changes is yet to be confirmed.

12.3 Pharmacokinetics

Absorption

WEGOVY Injection

Absolute bioavailability of semaglutide is 89% following subcutaneous administration. Maximum concentration of semaglutide is reached 1 to 3 days post dose.

Similar exposure was achieved with subcutaneous administration of semaglutide in the abdomen, thigh, or upper arm.

The average semaglutide steady state concentration following subcutaneous administration of WEGOVY 2.4 mg once weekly was approximately 75 nmol/L in patients with obesity or overweight (BMI greater than or equal to 27 kg/m^2). The average semaglutide steady state concentration following subcutaneous administration of WEGOVY 7.2 mg once weekly injection was approximately 230 nmol/L in patients with obesity (BMI greater than or equal to 30 kg/m^2). The steady state exposure concentrations increased proportionally with doses up to 7.2 mg once weekly.

WEGOVY Tablets

WEGOVY tablets are co-formulated with SNAC which facilitates the absorption of semaglutide after oral WEGOVY tablet administration. The absorption of semaglutide predominantly occurs in the stomach.

Absolute bioavailability of semaglutide is estimated to be approximately 1% to 2% following oral tablet administration. Maximum concentration of semaglutide is reached 1 hour post dose.

The average semaglutide steady state concentration following 25 mg tablet oral administration was approximately 77 nmol/L in patients with obesity or overweight (BMI greater than or equal to 27 kg/m^2). The steady-state concentrations increased approximately proportionally with doses up to 25 mg once daily.

In patients with overweight or obesity without type 2 diabetes, semaglutide concentrations following once-daily administration of oral WEGOVY 25 mg tablet are predicted to be comparable to WEGOVY 2.4 mg once-weekly injection, with higher variability in semaglutide concentrations compared to subcutaneous administration (90% of patients had average concentrations between 27 and 186 nmol/L with WEGOVY 25 mg tablet versus 51 and 110 nmol/L with WEGOVY 2.4 mg once-weekly injection).

Effect of Volume and Timing of Water Consumption: Single doses of oral semaglutide

tablet were administered with 50 mL or 240 mL of water after an 8-hour overnight fast and a continued fast of 4 hours post-dose in healthy subjects. Semaglutide absorption [i.e., area under the curve (AUC) and peak concentrations (C_{max})] was higher following dosing with 50 mL water compared to that of 240 mL water.

For 10 days, healthy subjects received once daily doses of oral semaglutide with 50 mL or 120 mL of water under fasting conditions with post-dose fasting period of 15, 30, 60, or 120 minutes. In this study, semaglutide absorption (i.e., AUC and C_{max}) was higher after a longer post-dose fasting period. There were no clinically significant differences in semaglutide absorption with administration of 50 mL or 120 mL of water.

Distribution

The mean volume of distribution of semaglutide following subcutaneous administration in patients with obesity or overweight is approximately 12.5 L. Semaglutide is extensively bound to plasma albumin (greater than 99%) which results in decreased renal clearance and protection from degradation.

Elimination

The apparent clearance of semaglutide in patients with overweight or obesity is approximately 0.05 L/h. With an elimination half-life of approximately 1 week, semaglutide will be present in the circulation for about 5 to 7 weeks after the last injectable dose of 2.4 mg or 7.2 mg or oral dose of 25 mg.

Metabolism:

The primary route of elimination for semaglutide is metabolism following proteolytic cleavage of the peptide backbone and sequential beta-oxidation of the fatty acid sidechain.

Excretion:

The primary excretion routes of semaglutide-related material are via the urine and feces. Approximately 3% of the dose is excreted in the urine as intact semaglutide.

Specific Populations

Age, Sex, Race, Ethnicity, Body Weight, and Other Intrinsic Factors: No clinically significant differences in semaglutide pharmacokinetics were observed based on age (<65 years, 65 to 74, ≥75 years old), sex, race (White, Asian, or Black or African American), ethnicity, body weight, upper GI disease (i.e. chronic gastritis and/or gastroesophageal reflux disease), renal impairment [estimated glomerular filtration rate (eGFR) ≥30 to <90 mL/minute], or mild, moderate, and severe hepatic impairment (Child-Pugh Class A-C) or fibrosis stage (F2 or F3 in patients with MASH).

There is no difference observed in the exposure of semaglutide following subcutaneous administration of WEGOVY between patients with MASH and patients with overweight or obesity.

Patients with Type 2 Diabetes:

The average semaglutide steady state concentration following subcutaneous administration of WEGOVY 2.4 mg once weekly was approximately 68 nmol/L in patients with type 2 diabetes and obesity or overweight (BMI greater than or equal to 27 kg/m^2). The average semaglutide steady state concentration following 25 mg oral administration was approximately 44 nmol/L in patients with type 2 diabetes and obesity or overweight (BMI greater than or equal to 25 kg/m^2) with higher variability in semaglutide

concentrations compared to subcutaneous administration (90% of patients had average concentrations between 11 nmol/L and 134 nmol/L with WEGOVY 25 mg tablet versus 47 nmol/L and 99 nmol/L with WEGOVY 2.4 mg once-weekly injection) *[see Use in Specific Populations (8.6)]*.

Drug Interactions Studies

Subcutaneous Injection

The delay of gastric emptying with semaglutide may influence the absorption of concomitantly administered oral medications *[see Drug Interactions (7.2)].*

Oral Tablets

Clinical Studies and Model-Informed Approaches: The delay of gastric emptying with semaglutide may influence the absorption of concomitantly administered oral drugs *[see Drug Interactions (7.2)]*. Trials were conducted to study the potential effect of semaglutide on the absorption of oral drugs taken with semaglutide tablets administered orally at steady-state exposure.

- *Levothyroxine*: Total thyroxine (i.e., adjusted for endogenous levels) AUC of was increased by 33% following administration of a single dose of levothyroxine 600 mcg concomitantly administered with oral semaglutide. Maximum exposure (C_{max}) was unchanged.
- *Other Drugs*: No clinically significant differences in semaglutide pharmacokinetics were observed when used concomitantly with omeprazole. No clinically significant differences in the pharmacokinetics of the following drugs were observed when used concomitantly with semaglutide: lisinopril, S-warfarin, R-warfarin, metformin, digoxin, ethinyl estradiol, levonorgestrel, furosemide or rosuvastatin.

In Vitro Studies: Semaglutide has very low potential to inhibit or induce CYP enzymes, and to inhibit drug transporters.

12.6 Immunogenicity

The observed incidence of anti-drug antibodies is highly dependent on the sensitivity and specificity of the assay. Differences in assay methods preclude meaningful comparisons of the incidence of anti-drug antibodies in the studies described below with the incidence of anti-drug antibodies in other studies, including those of semaglutide or of other semaglutide products.

During the 68-week treatment periods in Studies 2 and 3 in adults *[see Clinical Studies (14.2)],* 50/1709 (3%) of patients treated with WEGOVY 2.4 mg injection developed anti-semaglutide antibodies (referred to as anti-drug-antibodies or ADA). A total of 28 (56% of the 50 patients treated with WEGOVY 2.4 mg injection who tested ADA positive) patients developed antibodies that cross-reacted with native GLP-1 in Studies 2 and 3.

Using a new assay, 202/1311 (15.4%) of patients treated with WEGOVY 7.2 mg injection and 34/304 (11.2%) of patients treated with WEGOVY 2.4 mg injection developed anti-semaglutide antibodies during 72-week treatment periods in Studies 8 and 9. A total of 36 (17.8% of 202 patients treated with WEGOVY 7.2 mg injection who tested ADA positive) and 3 (8.8% of 34 patients treated with WEGOVY 2.4 mg injection who tested ADA positive) patients developed antibodies that cross reacted with native GLP-1 in Studies 8 and 9. The clinical consequences of antibodies that cross react with native GLP-1 are unknown.

In adults with MASH treated with WEGOVY injection for 72 weeks in Study 11 *[see Clinical Studies (14.4)]*, 3/763 (0.4%) of patients developed ADA which were also cross-reactive to native GLP-1. No identified clinically significant effect of ADA on pharmacokinetics for WEGOVY injection was observed.

There is insufficient evidence to characterize the effects of ADA on pharmacodynamics, safety, or effectiveness of WEGOVY.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

In a 2-year carcinogenicity study in CD-1 mice, subcutaneous semaglutide doses of 0.3, 1 and 3 mg/kg/day (up to 7-fold the MRHD of WEGOVY subcutaneous injection 7.2 mg/week, based on AUC) were administered to the males, and 0.1, 0.3 and 1 mg/kg/day (up to 2-fold MRHD) were administered to the females. A statistically significant increase in thyroid C-cell adenomas and a numerical increase in C-cell carcinomas were observed in males and females at all dose levels, at clinically relevant exposures.

In a 2-year carcinogenicity study in Sprague Dawley rats, subcutaneous semaglutide doses of 0.0025, 0.01, 0.025 and 0.1 mg/kg/day were administered (up to 0.7-fold the exposure at the MRHD). A statistically significant increase in thyroid C-cell adenomas was observed in males and females at all dose levels, and a statistically significant increase in thyroid C-cell carcinomas was observed in males at greater than or equal to 0.01 mg/kg/day, at clinically relevant exposures.

Human relevance of thyroid C-cell tumors in rats is unknown and could not be determined by clinical studies or nonclinical studies *[see Warnings and Precautions (5.1)]*.

Semaglutide was not mutagenic or clastogenic in a standard battery of genotoxicity tests (i.e., bacterial mutagenicity [Ames], human lymphocyte chromosome aberration, rat bone marrow micronucleus).

In a combined fertility and embryo-fetal development study in rats, subcutaneous semaglutide doses of 0.01, 0.03 and 0.09 mg/kg/day (up to 0.1-fold the MRHD) were administered to male and female rats. Males were dosed for 4 weeks prior to mating, and females were dosed for 2 weeks prior to mating and throughout organogenesis until Gestation Day 17. No effects were observed on male fertility. In females, an increase in estrus-cycle length was observed at all dose levels, together with a small reduction in numbers of corpora lutea at greater than or equal to 0.03 mg/kg/day, at clinically relevant exposures. These effects were likely an adaptive response secondary to the pharmacological effect of semaglutide on food consumption and body weight.

13.2 Animal Toxicology and/or Pharmacology

WEGOVY tablets contain SNAC as an absorption enhancer. An increase in lactate levels and a decrease in glucose levels in plasma and cerebrospinal fluid (CSF) were observed in mechanistic rat studies with SNAC. Small but statistically significant increases in lactate levels (up to 2-fold) were observed in a few rats at approximately the clinical exposure. At higher exposures these findings were associated with moderate to marked adverse clinical signs (lethargy, abnormal respiration, ataxia, and reduced activity, body tone and reflexes) and marked decreases in plasma and CSF glucose levels. These findings are consistent with inhibition of cellular respiration and lead to mortality at SNAC

concentrations >100-times the clinical C_{max}.

14 CLINICAL STUDIES

14.1 Cardiovascular Outcomes Trial in Adult Patients with Cardiovascular Disease and Either Obesity or Overweight

Overview of Clinical Trial

Study 1 (NCT03574597) was a multi-national, multi-center, placebo-controlled, double-blind trial to determine the effect of WEGOVY injection relative to placebo on major adverse CV events (MACE) when added to current standard of care, which included management of CV risk factors and individualized healthy lifestyle counseling (including diet and physical activity) in adults with established CV disease and either obesity or overweight. The primary endpoint, MACE, was the time to first occurrence of a three-part composite outcome which included CV death, non-fatal myocardial infarction, and non-fatal stroke. During the 16-week dose escalation period, patients were titrated to WEGOVY 2.4 mg injected subcutaneously once weekly or placebo. Patients who could not tolerate the recommended escalation dosage could stay at a lower dose level.

All patients were 45 years or older, with an initial BMI of 27 kg/m^2 or greater and established CV disease (prior myocardial infarction, prior stroke, or peripheral arterial disease). Patients with a history of type 1 or type 2 diabetes were excluded. Concomitant CV therapies could be adjusted, at the discretion of the investigator, to ensure participants were treated according to the current standard of care for patients with established CV disease.

In this trial, 17,604 patients were randomized to WEGOVY injection or placebo. At baseline, the mean age was 62 years (range 45 to 93), 72% were male, 84% were White, 4% were Black or African American, and 8% were Asian, and 10% were Hispanic or Latino. Mean baseline body weight was 97 kg and mean BMI was 33 kg/m^2. At baseline, prior myocardial infarction was reported in 76% of randomized individuals, prior stroke in 23%, and peripheral arterial disease in 9%. Heart failure was reported in 24% of patients. At baseline, CV disease and risk factors were managed with lipid-lowering therapy (90%), platelet aggregation inhibitors (86%), angiotensin converting enzyme inhibitors or angiotensin II receptor blockers (74%), and beta blockers (70%). A total of 10% had moderate renal impairment (eGFR 30 to <60 mL/min/1.73m^2) and 0.4% had severe renal impairment eGFR <30 mL/min/1.73m^2.

Results

In total, 96.9% of patients completed the trial, and vital status was available for 99.4% of patients. The median follow-up duration was 41.8 months. A total of 31% of WEGOVY injection-treated patients and 27% of placebo-treated patients permanently discontinued study drug. Of those patients on treatment at 1 year, 76% were on the 2.4 mg dose, 8% were on the 1.7 mg dose, and 16% were on lower doses. Of those patients on treatment at 2 years, 77% were on the 2.4 mg dose, 7% were on the 1.7 mg dose, and 17% were on lower doses.

For the primary analysis, a Cox proportional hazards model was used to test for superiority. Type 1 error was controlled across multiple tests.

WEGOVY injection significantly reduced the risk for first occurrence of MACE. The estimated hazard ratio (95% CI) was 0.80 (0.72, 0.90) (see **Figure 2** and **Table 6**).

Figure 2. Cumulative Incidence Function: Time to First Occurrence of MACE in Study 1 with WEGOVY Injection



Data from the in-trial period. Cumulative incidence estimates are based on time from randomization to first EAC-confirmed cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke with non-CV death modeled as competing risk using the Aalen-Johansen estimator. Patients without events of interest were censored at the end of their in-trial observation period. Time from randomization to first cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke was analyzed using a Cox proportional hazards model with treatment as categorical fixed factor. The hazard ratio and confidence interval are adjusted for the group sequential design using the likelihood ratio ordering.

HR: Hazard ratio; CI: confidence interval; CV: cardiovascular

The treatment effect for the primary composite endpoint, its components, and other relevant endpoints in Study 1 are shown in **Table 6**.

Table 6. Treatment Effect for MACE and Other Events in Study 1 with WEGOVY Injection

	Patients with events n (%)		
	Placebo N=8,801	**WEGOVY Injection N=8,803**	**Hazard Ratio (95% CI)**

Primary composite endpoint			
Composite of CV death, non-fatal myocardial infarction, or non-fatal stroke[1]	701 (8%)	569 (6.5%)	0.8 (0.72; 0.9)[*,2]
Key secondary endpoints			
CV death[3]	262 (3%)	223 (2.5%)	0.85 (0.71; 1.01)
All-cause death[4]	458 (5.2%)	375 (4.3%)	0.81 (0.71; 0.93)
Other endpoints			
Fatal or non-fatal myocardial infarction[5]	334 (3.8%)	243 (2.8%)	0.72 (0.61; 0.85)
Fatal or non-fatal stroke[5]	178 (2%)	160 (1.8%)	0.89 (0.72; 1.11)
Hospitalization for heart failure or urgent heart failure visits[5,6]	122 (1.4%)	97 (1.1%)	0.79 (0.6; 1.03)

[*] p-value <0.001, one-sided p-value

[1] Primary endpoint

[2] Adjusted for group sequential design using the likelihood ratio ordering.

[3] CV death was the first confirmatory secondary endpoint in the testing hierarchy and superiority was not confirmed.

[4] Confirmatory secondary endpoint. Not statistically significant based on the prespecified testing hierarchy.

[5] Not included in the prespecified testing hierarchy for controlling type-I error.

[6] Effect on heart failure has not been established.

NOTE: Time to first event was analyzed in a Cox proportional hazards model with treatment as factor. For patients with multiple events, only the first event contributed to the composite endpoint.

Table 7. Mean Changes in Anthropometry and Cardiometabolic Parameters at Week 104 in Study 1[1,2]

	PLACEBO		**WEGOVY Injection**		
	Baseline	**Change from Baseline (LSMean)**	**Baseline**	**Change from Baseline (LSMean)**	**Difference from Placebo (LSMean)**
Body Weight (kg)	96.8	-0.9[3]	96.5	-9.4[3]	-8.5[3]
Waist Circumference (cm)	111.4	-1	111.3	-7.6	-6.5
Systolic Blood	131	-0.5	131	-3.8	-3.3

Pressure (mmHg)					
Diastolic Blood Pressure (mmHg)	79	-0.5	79	-1	-0.5
Heart Rate	69	0.7	69	3.8	3.1
HbA_{1c} (%)	5.8	0	5.8	-0.3	-0.3
	Baseline	**% Change from Baseline (LSMean)**	**Baseline**	**% Change from Baseline (LSMean)**	**Relative Difference from Placebo (%) (LSMean)**
Total Cholesterol (mg/dL)[4]	156	-1.9	155.5	-4.6	-2.8
LDL Cholesterol (mg/dL)[4]	78.5	-3.1	78.5	-5.3	-2.2
HDL Cholesterol (mg/dL)[4]	44.2	0.6	44.1	4.9	4.2
Triglycerides (mg/dL)[4]	139.5	-3.2	138.6	-18.3	-15.6

[1] Parameters listed in the table were not included in the pre-specified hierarchical testing.

[2] Responses were analyzed using an ANCOVA with treatment as fixed factor and baseline value as covariate. Before analysis, missing data were multiple imputed. The imputation model (linear regression) was done separately for each treatment arm and included baseline value as a covariate and was fitted to all subjects with a measurement regardless of treatment status at Week 104.

[3] For body weight the 'change from baseline' and 'difference to placebo' the unit is percentage change from baseline.

[4] Baseline value is the geometric mean.

The reduction of MACE with WEGOVY injection was not impacted by age, sex, race, ethnicity, BMI at baseline, or level of renal function impairment.

14.2 Weight Reduction and Long-term Maintenance Studies in Adults with Obesity or Overweight

Overview of Clinical Trials in Adults

The efficacy of WEGOVY once weekly subcutaneous injection for weight reduction and long-term maintenance of body weight in conjunction with a reduced-calorie diet and increased physical activity were studied in three 68-week, randomized, double-blind, placebo-controlled trials; one 68-week, randomized, double-blind, placebo withdrawal trial; one 68-week, randomized, double-blind trial that investigated two different doses of WEGOVY injection versus placebo and two 72-week randomized, double-blind trials that investigated two different doses of WEGOVY injection versus placebo. In Studies 2 (NCT#03548935), 3 (NCT#03552757), and 4 (NCT#03611582), WEGOVY or matching placebo was escalated to 2.4 mg subcutaneous injection weekly during a 16-week period followed by 52 weeks on maintenance dose. In Study 5 (NCT#03548987), WEGOVY injection was escalated during a 20-week run-in period, and patients who reached a WEGOVY 2.4 mg subcutaneous injection weekly dosage after the run-in period were

randomized to either continued treatment with WEGOVY injection or placebo for 48 weeks. In Study 6 (NCT#03811574), WEGOVY injection was escalated to 1.7 mg or 2.4 mg subcutaneous weekly dosages or placebo over 12 to 16 weeks followed by 52 weeks on either maintenance dose. In Study 8 (NCT#05646706) and Study 9 (NCT#05649137) WEGOVY injection was escalated to 2.4 mg or 7.2 mg subcutaneous weekly dosages or placebo over 20 weeks followed by 52 weeks on maintenance dose.

The efficacy of WEGOVY 25 mg oral once daily tablet for weight reduction and long-term maintenance of body weight in conjunction with a reduced-calorie diet and increased physical activity was evaluated in a 64-week randomized, double-blind, placebo-controlled trial [Study 7 (NCT#05564117)]. WEGOVY tablet or matching placebo was escalated to a 25 mg tablet once daily during a 12-week period followed by 52-week maintenance period.

In Studies 2, 3, 5, 7, 8 and 9, all patients received instruction for a reduced-calorie diet (approximately 500 kcal/day deficit) and increased physical activity counseling (recommended to a minimum of 150 min/week) that began with the first dose of study medication or placebo and continued throughout the trial. In Study 4, patients received an initial 8-week low-calorie diet (total energy intake 1,000 to 1,200 kcal/day) followed by 60 weeks of a reduced-calorie diet (1200 to 1800 kcal/day) and increased physical activity (100 mins/week with gradual increase to 200 mins/week).

Study 2 was a 68-week trial that enrolled 1,961 patients with obesity (BMI greater than or equal to 30 kg/m^2) or with overweight (BMI 27 to 29.9 kg/m^2) and at least one weight-related comorbid condition, such as treated or untreated dyslipidemia or hypertension; patients with type 2 diabetes mellitus were excluded. Patients were randomized in a 2:1 ratio to either WEGOVY injection or placebo. At baseline, mean age was 46 years (range 18 to 86), 74% were female, 75% were White, 13% were Asian and 6% were Black or African American. A total of 12% were Hispanic or Latino ethnicity. Mean baseline body weight was 105.3 kg and mean BMI was 37.9 kg/m^2.

Study 3 was a 68-week trial that enrolled 807 patients with type 2 diabetes and BMI greater than or equal to 27 kg/m^2. Patients included in the trial had HbA_{1c} 7-10% and were treated with either: diet and exercise alone or 1 to 3 oral anti-diabetic drugs (metformin, sulfonylurea, glitazone or sodium-glucose co-transporter 2 inhibitor). Patients were randomized in a 1:1 ratio to receive either WEGOVY injection or placebo. At baseline, the mean age was 55 years (range 19 to 84), 51% were female, 62% were White, 26% were Asian and 8% were Black or African American. A total of 13% were Hispanic or Latino ethnicity. Mean baseline body weight was 99.8 kg and mean BMI was 35.7 kg/m^2.

Study 4 was a 68-week trial that enrolled 611 patients with obesity (BMI greater than or equal to 30 kg/m^2) or with overweight (BMI 27 to 29.9 kg/m^2) and at least one weight-related comorbid condition such as treated or untreated dyslipidemia or hypertension; patients with type 2 diabetes mellitus were excluded. The patients were randomized in a 2:1 ratio to receive either WEGOVY injection or placebo. At baseline, the mean age was 46 years, 81% were female, 76% were White, 19% were Black or African American and 2% were Asian. A total of 20% were Hispanic or Latino ethnicity. Mean baseline body weight was 105.8 kg and mean BMI was 38 kg/m^2.

Study 5 was a 68-week trial that enrolled 902 patients with obesity (BMI greater than or equal to 30 kg/m^2) or with overweight (BMI 27 to 29.9 kg/m^2) and at least one weight-related comorbid condition such as treated or untreated dyslipidemia or hypertension; patients with type 2 diabetes mellitus were excluded. Mean body weight at baseline for the 902 patients was 106.8 kg and mean BMI was 38.3 kg/m^2. All patients received

WEGOVY injection during the run-in period of 20 weeks that included 16 weeks of dose escalation. Trial product was permanently discontinued before randomization in 99 of 902 patients (11%); the most common reason was adverse reactions (n=48, 5.3%); 803 patients reached WEGOVY 2.4 mg injection and were then randomized in a 2:1 ratio to either continue on WEGOVY injection or receive placebo. Among the 803 randomized patients, the mean age was 46 years, 79% were female, 84% were White, 13% were Black or African American, and 2% Asian. A total of 8% were Hispanic or Latino ethnicity. Mean body weight at randomization (Week 20) was 96.1 kg and mean BMI at randomization (Week 20) was 34.4 kg/m^2.

Study 6 was a 68-week trial that enrolled 401 East-Asian patients (Japan and South Korea) with BMI greater than or equal to 35 kg/m^2 and at least one weight-related comorbid condition or with BMI 27 to 34.9 kg/m^2 and at least two weight-related comorbid conditions. The patients were randomized 2:1:1 to receive subcutaneous injection of WEGOVY 2.4 mg, WEGOVY 1.7 mg, or placebo. At baseline, the mean age was 51 years, 63% were male, and all patients were Asian. Mean baseline body weight was 87.5 kg and mean BMI was 31.9 kg/m^2. At baseline, 24.7% of patients had type 2 diabetes mellitus.

Study 7 was a 64-week trial that enrolled 307 patients with obesity (BMI greater than or equal to 30 kg/m^2) or with overweight (BMI 27-29.9 kg/m^2) and at least one weight-related comorbid condition, such as treated or untreated dyslipidemia or hypertension; patients with type 2 diabetes mellitus were excluded. Patients were randomized in a 2:1 ratio to either WEGOVY 25 mg tablet or placebo. At baseline, mean age was 48 years, 79% were women, 92% were White, 1% were Asian and 7% were Black or African American. A total of 7.8% were Hispanic or Latino ethnicity. Mean baseline body weight was 106 kg and mean BMI was 38 kg/m^2.

Study 8 was a 72-week trial that enrolled 1,407 patients with obesity (BMI greater than or equal to 30 kg/m^2), patients with type 2 diabetes mellitus were excluded. Patients were randomized in a 5:1:1 ratio to either WEGOVY 7.2 mg, WEGOVY 2.4 mg or placebo once weekly. At baseline, mean age was 47 years (range 18 to 80), 74% were female, 86% were White, 4% were Asian and 9% were Black or African American. A total of 5% were Hispanic or Latino ethnicity. Mean baseline body weight was 113 kg and mean BMI was 39.9 kg/m^2.

Study 9 was a 72-week trial that enrolled 512 patients with obesity and type 2 diabetes. Patients included in the trial had HbA_{1c} 7 to 10% and were treated with either: diet and exercise alone or 1 to 3 oral antidiabetic drugs (metformin, sulfonylurea, glitazone or sodium-glucose co-transporter 2 inhibitor). Patients were randomized in a 3:1:1 ratio to receive either WEGOVY 7.2 mg, WEGOVY 2.4 mg or placebo once weekly. At baseline, the mean age was 56 years (range 19 to 79), 52% were female, 84% were White, 6% were Asian and 9% were Black or African American. A total of 6% were Hispanic or Latino ethnicity. Mean baseline body weight was 110.1 kg and mean BMI was 38.6 kg/m^2.

Results

In Studies 2, 3, and 4, 16% of WEGOVY injection-treated patients and 19% of the placebo-treated patients discontinued therapy. In these studies, 6.8% of WEGOVY injection-treated patients and 3.2% of placebo-treated patients discontinued their treatment due to an adverse reaction *[see Adverse Reactions (6.1)]*. In Study 5, the proportions of patients who discontinued study drug were 5.8% and 11.6% for WEGOVY injection and placebo, respectively. In Study 6, the proportions of patients who

discontinued study drug were 7.9%, 6.5%, and 3% for WEGOVY 1.7 mg injection, WEGOVY 2.4 mg injection, and placebo, respectively. In Study 7, the proportions of patients who discontinued study drug were 18% and 25.5% for WEGOVY 25 mg tablet and placebo, respectively. Adverse reactions led to treatment discontinuation in 6.8% of patients treated with WEGOVY 25 mg tablet and 5.9% patients receiving placebo. In Study 8, 5.4% of patients treated with WEGOVY 7.2 mg and 1% patients treated with placebo permanently discontinued treatment as a result of adverse reactions. In Study 9, 5.5% of patients treated with WEGOVY 7.2 mg and 2% patients treated with placebo permanently discontinued treatment as a result of adverse reactions.

For Studies 2, 3 and 4, the primary efficacy parameters were mean percent change in body weight and the percentages of patients achieving greater than or equal to 5% weight loss from baseline to Week 68.

After 68 weeks, treatment with WEGOVY injection resulted in a statistically significant reduction in body weight compared with placebo. Greater proportions of patients treated with WEGOVY injection achieved 5%, 10% and 15% weight loss than those treated with placebo as shown in **Table 8**.

Table 8. Changes in Body Weight at Week 68 in Studies 2, 3, and 4 with WEGOVY 2.4 mg Injection

	Study 2 (Obesity or Overweight with Comorbidity)		**Study 3 (Type 2 Diabetes with Obesity or Overweight)**		**Study 4 (Obesity or Overweight with Comorbidity Undergoing Intensive Lifestyle Therapy)**	
Intention-to-Treat[1]	**PLACEBO N=655**	**WEGOVY Injection N=1,306**	**PLACEBO N=403**	**WEGOVY Injection N=404**	**PLACEBO N=204**	**WEGOVY Injection N=407**
Body Weight						
Baseline mean (kg)	105.2	105.4	100.5	99.9	103.7	106.9
% change from baseline (LSMean)	-2.4	-14.9	-3.4	-9.6	-5.7	-16
% difference from placebo (LSMean) (95% CI)		-12.4 (-13.3; -11.6)*		-6.2 (-7.3; -5.2)*		-10.3 (-11.8; -8.7)*
% of Patients	31.1	83.5	30.2	67.4	47.8	84.8

losing greater than or equal to 5% body weight						
% difference from placebo (LSMean) (95% CI)		52.4 (48.1; 56.7)*		37.2 (30.7; 43.8)*		37 (28.9; 45.2)*
% of Patients losing greater than or equal to 10% body weight	12	66.1	10.2	44.5	27.1	73
% difference from placebo (LSMean) (95% CI)		54.1 (50.4; 57.9)*		34.3 (28.4; 40.2)*		45.9 (38; 53.7)*
% of Patients losing greater than or equal to 15% body weight	4.8	47.9	4.3	25.1	13.2	53.4
% difference from placebo (LSMean) (95% CI)		43.1 (39.8; 46.3)*		20.7 (15.7; 25.8)*		40.2 (33.1; 47.3)*
% of Patients losing greater than or equal to 20% body weight[2]	1.7	30.2	2.3	12.8	3.5	33.9
% difference from placebo (LSMean) (95% CI)[2]		28.6 (25,8; 31.3)		10.6 (6.4; 14.7)		30.4 (24.8; 36.1)

LSMean = least squares mean; CI = confidence interval

[1] The intent-to-treat population includes all randomized patients. In Study 2, at Week 68, the body weight was missing for 7.2% and 11.9% of patients randomized to WEGOVY injection and placebo, respectively. In Study 3, at Week 68, the body weight was missing for 4% and 6.7% of patients randomized to WEGOVY injection and placebo, respectively. In Study 4, at Week 68, the body weight was missing for 8.4% and 7.4% of patients randomized to WEGOVY injection and placebo, respectively. Missing data were imputed from retrieved subjects of the same randomized treatment arm (RD-MI).

[2] Not included in the pre-specified hierarchical testing.

[*] $p<0.0001$ (unadjusted 2-sided) for superiority.

For Study 5, the primary efficacy parameter was mean percent change in body weight from randomization (Week 20) to Week 68.

From randomization (week 20) to Week 68, treatment with WEGOVY injection resulted in a statistically significant reduction in body weight compared with placebo (**Table 9**). Because patients who discontinued WEGOVY injection during titration and those who did not reach the 2.4 mg weekly dose were not eligible for the randomized-treatment period, the results may not reflect the experience of patients in the general population who are first starting WEGOVY injection.

Table 9. Changes in Body Weight at Week 68 in Study 5 (Obesity or Overweight with Comorbidity After 20-week Run-in) with WEGOVY 2.4 mg Injection

	WEGOVY Injection N=803[1]	
Body Weight (only randomized patients)		
Mean at Week 0 (kg)	107.2	
	PLACEBO N=268	**WEGOVY Injection N=535**
Body Weight		
Mean at Week 20 (SD) (kg)	95.4 (22.7)	96.5 (22.5)
% change from Week 20 at Week 68 (LSMean)	6.9	-7.9
% difference from placebo (LSMean) (95% CI)		-14.8 (-16; -13.5)[*]

LSMean = least squares mean; CI = confidence interval

[1] 902 patients were enrolled at Week 0 with a mean baseline body weight of 106.8 kg. The intent-to-treat population includes all randomized patients. At Week 68, the body weight was missing for 2.8% and 6.7% of patients randomized to WEGOVY injection and placebo, respectively. Missing data were imputed from retrieved subjects of the same

randomized treatment arm (RD-MI).

* p<0.001 (unadjusted 2-sided) for superiority, controlled for multiplicity.

For Study 6, the primary efficacy parameters were mean percent change in body weight and the percentage of patients achieving greater than or equal to 5% weight loss from baseline to Week 68.

After 68 weeks, treatment with WEGOVY 1.7 mg and 2.4 mg injection resulted in a statistically significant reduction in body weight compared with placebo. Greater proportions of patients treated with WEGOVY injection achieved 5%, 10%, and 15% weight loss than those treated with placebo as shown in **Table 10**.

Table 10. Changes in Body Weight at Week 68 in Study 6 in East-Asian Patients with WEGOVY 1.7 and 2.4 mg Injection

	Study 6 (BMI ≥35 kg/m^2 with at Least One Comorbidity or BMI 27 – 34.9 kg/m^2 with at Least Two Comorbidities)		
Intention-to-treat[1]	**PLACEBO N=101**	**WEGOVY 1.7 mg Injection N=101**	**WEGOVY 2.4 mg Injection N=199**
Body Weight			
Baseline mean (kg)	90.2	86.1	86.9
% change from baseline (LSMean)	-2.1	-9.6	-13.2
% difference from placebo (LSMean) (95% CI)		-7.5 (-9.6; -5.4)*	-11.1 (-12.9; -9.2)*
% of Patients losing greater than or equal to 5% body weight	19.4	72.8	84
% difference from placebo (LSMean) (95% CI)		53.3 (41; 65.6)*	64.5 (54.8; 74.3)*
% of Patients losing greater than or equal to 10% body weight	4.5	39.1	59.9
% difference from placebo (LSMean) (95% CI)		34.5 (23.9; 45.1)*	55.4 (47.3; 63.6)*

% of Patients losing greater than or equal to 15% body weight	2.6	20.8	38.2
% difference from placebo (LSMean) (95% CI)		18.2 (9.8; 26.7)*	35.6 (27.9; 43.3)*

LSMean = least squares mean; CI = confidence interval

[1] The intent-to-treat population includes all randomized patients. At baseline, 24.7% of patients had type 2 diabetes mellitus. At Week 68, the body weight was missing for 3%, 3%, and 1% of patients randomized to WEGOVY 1.7 mg injection, WEGOVY 2.4 mg injection, and placebo, respectively. Missing data were imputed from retrieved subjects of the same randomized treatment arm (RD-MI).

* $p<0.0001$ (unadjusted 2-sided) for superiority.

For Study 7, the primary efficacy parameters were mean percent change in body weight and the percentages of patients achieving greater than or equal to 5% weight loss from baseline to Week 64.

After 64 weeks, treatment with WEGOVY 25 mg tablets resulted in a statistically significant reduction in body weight compared with placebo. Greater proportions of patients treated with WEGOVY 25 mg tablets achieved 5%, 10%, 15% and 20% weight loss than those treated with placebo as shown in **Table 11**.

Table 11. Changes in Body Weight at Week 64 in Study 7 with WEGOVY 25 mg Tablets

	Study 7 (Obesity or Overweight with Comorbidity)	
Intention-to-Treat[1]	**PLACEBO N = 102**	**WEGOVY 25 mg Tablet N = 205**
Body Weight		
Baseline mean (kg)	104.8	106.4
% change from baseline (LSMean)[3]	-2.4	-13.6
% difference from placebo (LSMean) (95% CI)[3]		-11.2 [-13.6; -8.8][2]
% of Patients losing greater than or equal to 5% body weight[3]	31.3	76.3
% difference from placebo (LSMean) (95% CI)[3]		45.1 [33.8; 56.3][2]

% of Patients losing greater than or equal to 10% body weight	14.4	59.8
% difference from placebo (LSMean) (95% CI)[3]		45.3 [35.5; 55.2][2]
% of Patients losing greater than or equal to 15% body weight[3]	5.5	47
% difference from placebo (LSMean) (95% CI)[3]		41.5 [33.2; 49.8][2]
% of Patients losing greater than or equal to 20% body weight[3]	3.1	27.9
% difference from placebo (LSMean) (95% CI)[3]		24.8 [17.7; 31.9][2]

LSMean = least squares mean; CI = confidence interval

[1] The intent-to-treat population includes all randomized patients. In Study 7, at Week 64, the body weight was missing for 6.3% and 11.8% of patients randomized to WEGOVY tablets and placebo, respectively.

[2] $p<0.0001$ (unadjusted 2-sided) for superiority.

[3] Model based estimates are based on an analysis of covariance (or logistic regression for the 5%, 10%, 15% or 20% achievement endpoints) including treatment as a factor and baseline value as a covariate. Missing data were multiple imputed from retrieved subjects of the same randomized treatment arm (RD-MI). The imputation model was a linear regression of the endpoint with gender as factor and baseline value of the endpoint, timing and the value of last available observation – on study (LAO-OS) of the endpoint as covariates, imputed by treatment status. For the 5%, 10%, 15% or 20% achievement endpoints, the missing body weight measurements were imputed before performing the logistic regression analyses.

RD-MI: Retrieved Drop-outs Multiple Imputation

For Studies 8 and 9, the primary efficacy parameters were mean percent change in body weight and the percentages of patients achieving greater than or equal to 5% weight loss from baseline to Week 72.

After 72 weeks of treatment, WEGOVY 7.2 mg injection resulted in a statistically significant reduction in body weight compared with placebo in Studies 8 and 9 (see **Table 12** and **Figures 6** and **10**).

Table 12.Changes in Body Weight at Week 72 in Studies 8 and 9 with WEGOVY 7.2 mg Injection

	Study 8 (Patients with Obesity)			**Study 9 (Type 2 Diabetes with Obesity)**		
Intention-to-	**PLACEBO**	**WEGOVY 2.4**	**WEGOVY**	**PLACEBO**	**WEGOVY**	**WEGOVY**

Treat[1]	N=201	mg Injection N=201	7.2 mg Injection N=1005	N=102	2.4 mg Injection N=103	7.2 mg Injection N=307
Body Weight						
Baseline mean (kg)	112.4	116.5	112.4	112.1	107	110.5
% change from baseline (LSMean)[a]	-3.9	-15.5	-18.8	-3.8	-10.4**	-13.2
% difference from placebo (LSMean) (95% CI)			-14.9 [-16.2; -13.6]*			-9.4 [-11.0; -7.8]*
Difference (%) from 2.4 mg [95% CI]			-3.3 [-4.9; -1.6]*			
% of Patients losing greater than or equal to 5% body weight	38.3	86.6**	88.7	34	74**	83.7
% difference from placebo (LSMean) (95% CI)			50.4 [42.5; 58.3]*			49.8 [39.6; 59.9]*
% of Patients losing greater than or equal to 10% body weight	20.3	72**	79.8	11.2	49.5**	60.7
% difference from placebo (LSMean) (95% CI)			59.5 [53.0; 66.0]*			49.5 [41.2; 57.7]*

% of Patients losing greater than or equal to 15% body weight	7.5	52.3**	63.8	6.9	27**	39.8
% difference from placebo (LSMean) (95% CI)			56.4 [51.5; 61.2]*			32.9 [25.5; 40.3]*
% of Patients losing greater than or equal to 20% body weight	2.8	32.3	45.5	2	13.5**	20.6
% difference from placebo (LSMean) (95% CI)			42.7 [38.8; 46.6]*			18.7 [13.4; 24.1]*

* $p<0.0001$ (unadjusted 2-sided) for superiority.

** This endpoint was not included in the hierarchy testing and is only for informative purposes.

1 The intent-to-treat population includes all randomized patients. In Study 8, at Week 72, the body weight was missing for 5.5%, 6.0% and 15.0% of patients randomized to WEGOVY 7.2 mg, WEGOVY 2.4 mg and placebo, respectively. In Study 9, at week 72, the body weight was missing for 5.2% and 6.9% of patients randomized to WEGOVY 7.2 mg and placebo, respectively.

a Model based estimates are based on an analysis of covariance (or logistic regression for the 5%, 10%, 15% or 20% achievement endpoints) including treatment as a factor and baseline value as a covariate. Missing data were multiple imputed from retrieved subjects of the same randomized treatment arm (RD-MI) in Study 8 or all retrieved subjects regardless of treatment arm assignment in Study 9. The imputation model was a linear regression of the endpoint with gender as factor and baseline value of the endpoint, timing and the value of last available observation – on study (LAO-OS) of the endpoint as covariates, imputed by treatment status. For the 5%, 10%, 15% or 20% achievement endpoints, the missing body weight measurements were imputed before performing the logistic regression analyses.

RD-MI: Retrieved Drop-outs Multiple Imputation

A reduction in body weight was observed with WEGOVY irrespective of age, sex, race, ethnicity, BMI at baseline, body weight (kg) at baseline, and level of renal function impairment.

The cumulative frequency distributions of change in body weight are shown in **Figure 3**, **Figure 4**, **Figure 5** and **Figure 6** for Studies 2, 3, 7 and 8. One way to interpret this figure is to select a change in body weight of interest on the horizontal axis and note the corresponding proportions of patients (vertical axis) in each treatment group who achieved at least that degree of weight loss. For example, note that the vertical line arising from -10% in Study 2 intersects the WEGOVY injection and placebo curves at approximately 66%, and 12%, respectively, which correspond to the values shown in **Table 8**.

Figure 3. Change in Body Weight (%) from Baseline to Week 68 (Study 2) with WEGOVY Injection



Observed data from in-trial period including imputed data for missing observations (RD-MI).

Figure 4. Change in Body Weight (%) from Baseline to Week 68 (Study 3) with WEGOVY Injection



Observed data from in-trial period including imputed data for missing observations (RD-MI).

Figure 5. Change in Body Weight (%) from Baseline to Week 64 (Study 7) with WEGOVY 25 mg Tablets



Observed data from in-trial period including imputed data for missing observations (RD-MI). The missing observation was the patient-level average from the 1,000 imputed datasets.

Figure 6. Change in Body Weight (%) from Baseline to Week 72 (Study 8) with WEGOVY 7.2 mg Injection



Observed data from in-trial period including imputed data for missing observations (RD-MI). Missing observations were multiply (x1000) imputed from retrieved participants of the same randomized treatment arm and missing value was replaced by the average across imputations.

The time courses of weight loss with WEGOVY injection and placebo from baseline through Week 68 are depicted in **Figures 7**, **8**,and **9**, and through Week 72 in **Figure 10**.

Figure 7. Change from Baseline (%) in Body Weight (Study 2 on Left and Study 3 on Right) with WEGOVY Injection



Observed values for patients completing each scheduled visit, and estimates with multiple imputations from retrieved dropouts (RD-MI).

Figure 8. Change from Baseline (%) in Body Weight (Study 4 on Left and Study 5[a] on Right) with WEGOVY Injection



Observed values for patients completing each scheduled visit, and estimates with multiple imputations from retrieved dropouts (RD-MI).

[a] Change from Week 0 was not a primary endpoint in Study 5. Dotted line indicates time of randomization. Randomized patients (shown) do not include 99 patients that discontinued during the 20-week run-in period.

Figure 9. Change in Body Weight (%) from Baseline to Week 68 (Study 6 in East-Asian Patients) with WEGOVY Injection



Observed values for patients completing each scheduled visit and estimates with multiple imputations from retrieved dropouts (RD-MI). At baseline, 24.7% of patients had type 2 diabetes mellitus.

Figure 10. Mean Change in Body Weight (%) from Baseline to Week 72 (WEGOVY 7.2 mg Injection– Study 8)



Observed values for patients completing each scheduled visit, and estimates with multiple imputations from retrieved dropouts (RD-MI).

Effect of WEGOVY Injection on Anthropometry and Cardiometabolic Parameters in Adults

Changes in waist circumference and cardiometabolic parameters with WEGOVY injection are shown in **Table 13** for Studies 2, 3, and 4; in **Table 14** for Study 5; in **Table 15** for Study 6; **Table 16** for Study 7, and in **Table 17** for Studies 8 and 9.

Table 13. Changes in Anthropometry and Cardiometabolic Parameters at Week 68 in Studies 2, 3 and 4 with WEGOVY Injection

	Study 2 (Obesity or Overweight with Comorbidity)		**Study 3 (Type 2 Diabetes with Obesity or Overweight)**		**Study 4 (Obesity or Overweight with Comorbidity Undergoing Intensive Lifestyle Therapy)**	
Intention-to-Treat	**PLACEBO N=655**	**WEGOVY Injection N=1,306**	**PLACEBO N=403**	**WEGOVY Injection N=404**	**PLACEBO N=204**	**WEGOVY Injection N=407**
Waist	114.8	114.6	115.5	114.5	111.8	113.6

Circumference (cm) Baseline Changes from baseline (LSMean[1]) Difference from placebo (LSMean)	-4.1	-13.5 -9.4	-4.5	-9.4 -4.9	-6.3	-14.6 -8.3
Systolic Blood Pressure (mmHg) Baseline Changes from baseline (LSMean[1]) Difference from placebo (LSMean)	127 -1.1	126 -6.2 -5.1	130 -0.5	130 -3.9 -3.4	124 -1.6	124 -5.6 -3.9
Diastolic Blood Pressure (mmHg)[2] Baseline Changes from baseline (LSMean[1]) Difference from placebo (LSMean)	80 -0.4	80 -2.8 -2.4	80 -0.9	80 -1.6 -0.7	81 -0.8	80 -3 -2.2
Heart Rate[2,3] Baseline Changes from baseline (LSMean[1]) Difference from placebo (LSMean)	72 -0.7	72 3.5 4.3	76 -0.2	75 2.5 2.7	71 2.1	71 3.1 1

HbA_{1c} (%)[2] Baseline Changes from baseline (LSMean[1]) Difference from placebo (LSMean)	5.7 -0.2	5.7 -0.4 -0.3	8.1 -0.4	8.1 -1.6 -1.2	5.8 -0.3	5.7 -0.5 -0.2
Total Cholesterol (mg/dL)[2,4] Baseline Percent Change from baseline (LSMean[1]) Relative difference from placebo (LSMean)	192.1 0.1	189.6 -3.3 -3.3	170.8 -0.5	170.8 -1.4 -0.9	188.7 2.1	185.4 -3.9 -5.8
LDL Cholesterol (mg/dL)[2,4] Baseline Percent Change from baseline (LSMean[1]) Relative difference from placebo (LSMean)	112.5 1.3	110.3 -2.5 -3.8	90.1 0.1	90.1 0.5 0.4	111.8 2.6	107.7 -4.7 -7.1
HDL (mg/dL)[2,4] Baseline Percent Change from baseline	49.5 1.4	49.4 5.2 3.8	43.8 4.1	44.7 6.9 2.7	50.9 5	51.6 6.5 1.5

(LSMean[1]) Relative difference from placebo (LSMean)						
Triglycerides (mg/dL)[2,4] Baseline Percent Change from baseline (LSMean[1]) Relative difference from placebo (LSMean)	127.9 -7.3	126.2 -21.9 -15.8	159.5 -9.4	154.9 -22 -13.9	110.9 -6.5	107.9 -22.5 -17

Missing data were imputed from retrieved subjects of the same randomized treatment arm (RD-MI).

[1] Model based estimates based on an analysis of covariance model including treatment (and stratification factors for Study 3 only) as a factor and baseline value as a covariate.

[2] Not included in the pre-specified hierarchical testing (except HbA_{1c} for Study 3).

[3] Model based estimates based on a mixed model for repeated measures including treatment (and stratification factors for Study 3 only) as a factor and baseline values as a covariate.

[4] Baseline value is the geometric mean.

Table 14. Mean Changes in Anthropometry and Cardiometabolic Parameters in Study 5 (Obesity or Overweight with Comorbidity After 20-week Run-in)[1] with WEGOVY Injection

	PLACEBO N=268		**WEGOVY Injection N=535**		
	Randomization (Week 20)	**Change from Randomization (Week 20) to Week 68 (LSMean[1])**	**Randomization (Week 20)**	**Change from Randomization (Week 20) to Week 68 (LSMean[1])**	**Difference from Placebo (LSMean)**
Waist Circumference (cm)	104.7	3.3	105.5	-6.4	-9.7
Systolic Blood Pressure	121	4.4	121	0.5	-3.9

(mmHg)					
Diastolic Blood Pressure (mmHg)[2]	78	0.9	78	0.3	-0.5
Heart Rate[2,3]	76	-5.3	76	-2	3.3
HbA_{1c} (%)[2]	5.4	0.1	5.4	-0.1	-0.2
	Randomization (Week 20)	**% Change from Randomization (Week 20) (LSMean[1])**	**Randomization (Week 20)**	**% Change from Randomization (Week 20) (LSMean[1])**	**Relative difference from Placebo (LSMean)**
Total Cholesterol (mg/dL)[2,4]	175.1	11.4	175.9	4.9	-5.8
LDL Cholesterol (mg/dL)[2,4]	109.1	7.6	108.7	1.1	-6.1
HDL Cholesterol (mg/dL)[2,4]	43.6	17.8	44.5	18.2	0.3
Triglycerides (mg/dL)[2,4]	95.3	14.8	98.1	-5.6	-17.8

Missing data were imputed from retrieved subjects of the same randomized treatment arm (RD-MI).

[1] Model based estimates based on an analysis of covariance model including treatment as a factor and baseline value as a covariate.

[2] Not included in the pre-specified hierarchical testing.

[3] Model based estimates based on a mixed model for repeated measures including treatment as a factor and baseline values as a covariate.

[4] Baseline value is the geometric mean.

Table 15. Mean Changes in Anthropometry and Cardiometabolic Parameters at Week 68 in Study 6 in East-Asian Patients with WEGOVY Injection

	Study 6 (BMI ≥35 kg/m^2 with at least one comorbidity or BMI 27 to 34.9 kg/m^2 with at least two comorbidities)		
Intention-to-treat	**PLACEBO N=101**	**WEGOVY 1.7 mg Injection N=101**	**WEGOVY 2.4 mg Injection N=199**
Waist circumference (cm) Baseline Change from baseline (LSMean[1]) Difference from placebo	103.8 -1.8	101.4 -7.7 -5.9	103.8 -11 -9.3

(LSMean)			
Systolic blood pressure (mmHg)[2] Baseline Change from baseline (LSMean[1]) Difference from placebo (LSMean)	133 -5.3	135 -10.8 -5.4	133 -10.8 -5.5
Diastolic blood pressure (mmHg)[2] Baseline Change from baseline (LSMean[1]) Difference from placebo (LSMean)	86 -2.2	85 -4.6 -2.4	83 -5.3 -3.1
Heart Rate[2, 3] Baseline Change from baseline (LSMean[1]) Difference from placebo (LSMean)	73 2.4	73 4.4 2	73 6.3 3.9
HbA_{1c} (%)[2] Baseline Change from baseline (LSMean[1]) Difference from placebo (LSMean)	6.4 0	6.4 -0.9 -0.9	6.4 -0.9 -0.9
Total Cholesterol (mg/dL)[2,4] Baseline Percent change from baseline (LSMean[1]) Relative difference from placebo (LSMean)	203.1 0.8	203.3 -6.6 -7.3	197.2 -8.7 -9.4
LDL Cholesterol (mg/dL)[2,4] Baseline Percent change from baseline (LSMean[1]) Relative difference from placebo (LSMean)	123.3 -3.8	120.1 -10.1 -6.5	116.5 -14.6 -11.2
HDL Cholesterol (mg/dL)[2,4] Baseline Percent change from baseline (LSMean[1]) Relative difference from placebo (LSMean)	48.7 5.9	50.2 6.7 0.7	50.8 9.2 3.1
Triglyceride (mg/dL)[2,4] Baseline Percent change from baseline (LSMean[1]) Relative difference from placebo (LSMean)	134.2 5.5	138.8 -19.5 -23.7	127.1 -21.2 -25.3

Missing data were imputed from retrieved subjects of the same randomized treatment arm (RD-MI). At baseline, 24.7% of patients had type 2 diabetes mellitus.

[1] Model based estimates based on an analysis of covariance model including treatment and type 2 diabetes status as factors and baseline value as a covariate.

[2] Not included in the pre-specified hierarchical testing.

[3] Model based estimates based on a mixed model for repeated measures including treatment and type 2 diabetes status as factors and baseline values as a covariate.

[4] Baseline value is the geometric mean.

Table 16. Mean Changes in Anthropometry and Cardiometabolic Parameters in Study 7 with WEGOVY 25 mg Tablets (Obesity or Overweight with Comorbidity)

	Study 7 (Obesity or overweight with comorbidity)	
Intention-to-treat	**PLACEBO N=102**	**WEGOVY Tablets 25 mg N=205**
Waist circumference (cm)[2] Baseline Change from baseline (LSMean[1]) Difference from placebo (LSMean)	 113.6 -2.8	 114 -12.2 -9.4
Systolic blood pressure (mmHg)[2] Baseline Change from baseline (LSMean[1]) Difference from placebo (LSMean)	 131 -5.4	 131 -7 -1.6
Diastolic blood pressure (mmHg)[2] Baseline Change from baseline (LSMean[1]) Difference from placebo (LSMean)	 83 -1.8	 83 -2.8 -1
Heart Rate[2] Baseline Change from baseline (LSMean[1]) Difference from placebo (LSMean)	 74 2.4	 72 2.7 0.4
HbA_{1c} (%)[2] Baseline Change from baseline (LSMean[1]) Difference from placebo (LSMean)	 5.7 -0.1	 5.7 -0.3 -0.2
Total Cholesterol (mg/dL)[2,3] Baseline Percent change from baseline (LSMean[1]) Relative difference from placebo (LSMean)	 190.5 -2.2	 191.4 -3.3 -1.2
LDL Cholesterol (mg/dL)[2,3] Baseline Percent change from baseline (LSMean[1]) Relative difference from placebo (LSMean)	 114.2 0.2	 111.1 -2.9 -3.1

HDL Cholesterol (mg/dL)[2,3] Baseline Percent change from baseline (LSMean[1]) Relative difference from placebo (LSMean)	50.1 -1.1	49.6 3.3 4.5
Triglyceride (mg/dL)[2,3] Baseline Percent change from baseline (LSMean[1]) Relative difference from placebo (LSMean)	117.5 -9.1	115.9 -18.1 -9.9

[1] Model based estimates are based on an analysis of covariance including treatment as a factor and baseline value as a covariate. Missing data were multiple imputed from retrieved subjects of the same randomized treatment arm (RD-MI). The imputation model was a linear regression of the endpoint with gender as factor and baseline value of the endpoint, timing and the value of last available observation – on study (LAO-OS) of the endpoint as covariates, imputed by treatment status.

RD-MI: Retrieved Drop-outs Multiple Imputation

[2] Not included in the pre-specified hierarchical testing.

[3] Baseline value is the geometric mean.

Table 17. Changes in Anthropometry and Cardiometabolic Parameters at Week 72 in Studies 8 and 9 (WEGOVY 7.2 mg Injection)

	Study 8 (Patients with obesity)			**Study 9 (Type 2 diabetes with obesity)**		
Intention-to-Treat	**PLACEBO N=201**	**WEGOVY 2.4 mg Injection N=201**	**WEGOVY 7.2 mg Injection N=1,005**	**PLACEBO N=102**	**WEGOVY 2.4 mg Injection N=103**	**WEGOVY 7.2 mg Injection N=307**
Waist Circumference (cm) Baseline Changes from baseline (LSMean[1]) Difference from placebo (LSMean [95% CI])	118.6 -5.6 -	120.3 -14.2	118.4 -17.5 -11.9 [-13.2; -10.6]*	123.9 -5.8 -	119.1 -10.7	121.8 -12.3 -6.6 [-9; -4.1]*
Systolic Blood Pressure (mmHg)[2] Baseline	130 -4.8 -	129 -8.5	131 -10.1 -5.3	132 -3.5 -	134 -5.6	134 -7.5 -4

Changes from baseline (LSMean[1]) Difference from placebo (LSMean)						
Diastolic Blood Pressure (mmHg)[2] Baseline Changes from baseline (LSMean[1]) Difference from placebo (LSMean)	83 -1.8 -	83 -3	84 -4.5 -2.7	82 -2.9	83 -2.3	82 -3.1 -0.2
Heart Rate (beats/min)[2] Baseline Changes from baseline (LSMean[1]) Difference from placebo (LSMean)	75 -0.6	76 0.3	75 0.5 1	77 -2.7	76 1	77 1.1 3.9
HbA_{1c} (%)[2] Baseline Changes from baseline (LSMean[1]) Difference from placebo (LSMean [95% CI])	5.7 0.0	5.6 -0.3	5.7 -0.3 -0.3	8.2 -0.2	8.1 -1.6	8 -1.7 -1.5 [-1.8; -1.2]*
Total Cholesterol (mg/dL)[2,3] Baseline	193.4 -4.0 -	190.4 -3.6	193.6 -3.4 0.6	176.8 3.5 -	172.6 -1.4	175.2 0.1 -3.4

Percent change from baseline (LSMean[1])						
Relative difference from placebo (LSMean)						
LDL Cholesterol (mg/dL)[2,3]	116.4	114.4	115.8	93.5	93.8	91.5
Baseline	-4.8	-4.8	-3.5	6.7	2.7	10.5
Percent change from baseline (LSMean[1])	-		1.3	-		3.8
Relative difference from placebo (LSMean)						
HDL (mg/dL)[2,3]	48.9	48.2	48.8	42.5	41.9	42.6
Baseline	-0.5	7.0	9.5	7.4	6	11.4
Percent change from baseline (LSMean[1])	-		10.1			4.0
Relative difference from placebo (LSMean)						
Triglycerides (mg/dL)[2,3]	121.9	121.6	124.2	180.5	161.1	175.5
Baseline	-0.9	-9.8	-16.4	17.2	-14.8	-7.8
Percent change from baseline (LSMean[1])	-		-15.4			-25.0
Relative difference						

from placebo (LSMean)						

* p<0.0001 (unadjusted 2-sided) for superiority.

The difference or relative difference from placebo was comparison between Wegovy 7.2 mg and the placebo arms.

[1] Model based estimates based on an analysis of covariance model including randomized treatment as factor and baseline value as a covariate
Missing data were multiple imputed from retrieved subjects of the same randomized treatment arm (RD-MI) in Study 8 or all retrieved subjects regardless of treatment arm assignment in Study 9. The imputation model was a linear regression of the endpoint with gender as factor and baseline value of the endpoint, timing and the value of last available observation – on study (LAO-OS) of the endpoint as covariates, imputed by treatment status.

RD-MI: Retrieved Drop-outs Multiple Imputation

[2] Not included in the pre-specified hierarchical testing (except the comparison between placebo and Wegovy 7.2 mg in Study 8 and Study 9 for waist circumference and HbA_{1c} for Study 9).

[3] Baseline value is the geometric mean.

14.3 Weight Reduction and Long-term Maintenance Study of WEGOVY Injection in Pediatric Patients Aged 12 Years and Older with Obesity

Overview of Clinical Trial in Pediatric Patients

WEGOVY injection was evaluated to reduce excess body weight in pediatric patients aged 12 years of age and older with obesity in a 68-week, double-blind, randomized, parallel group, placebo-controlled, multi-center trial in 201 pubertal pediatric patients aged 12 years and older with BMI corresponding to ≥95th percentile standardized for age and sex (Study 10) (NCT#04102189). After a 12-week lifestyle run-in period (including dietary recommendations and physical activity counseling), patients were randomized 2:1 to WEGOVY injection once weekly or placebo once weekly. WEGOVY injection or matching placebo was escalated to 2.4 mg or maximally tolerated dose during a 16-week period followed by 52 weeks on maintenance dose. Of WEGOVY injection-treated patients who completed the trial, 86.7% were on the 2.4 mg dosage at the end of the trial; for 5% of patients, 1.7 mg was the maximum tolerated dosage.

The mean age was 15 years; 38% of patients were male; 79% were White, 8% were Black or African American, 2% were Asian, and 11% were of other or unknown race; and 11% were of Hispanic or Latino ethnicity. The mean baseline body weight was 108 kg, and mean BMI was 37 kg/m^2.

Results

The proportions of patients who discontinued study drug were 10% for the WEGOVY injection-treated group and 10% for the placebo-treated group.

The primary endpoint was percent change in BMI from baseline to Week 68. After 68 weeks, treatment with WEGOVY injection resulted in a statistically significant reduction in

percent BMI compared with placebo. Greater proportions of patients treated with WEGOVY injection achieved ≥5% reduction in baseline BMI than those treated with placebo as shown in **Table 18**.

Table 18. Changes in Weight and BMI at Week 68 in Pediatric Patients with Obesity Aged 12 Years and Older in Study 10 with WEGOVY 2.4 mg Injection in Study 10

Intention-to-Treat[a]	**PLACEBO N=67**	**WEGOVY 2.4 mg Injection N=134**
BMI		
Baseline mean (kg/m^2)	35.7	37.7
% change from baseline in BMI (LSMean)	0.6	-16.1
% difference from placebo (LSMean) (95% CI)		-16.7 (-20.3; -13.2)*
% of Patients with greater than or equal to 5% reduction in baseline BMI[b]	19.7	77.1
% difference from placebo (LSMean)		57.4
% of Patients with greater than or equal to 10% reduction in baseline BMI[b]	7.7	65.1
% difference from placebo (LSMean)		57.5
% of Patients with greater than or equal to 15% reduction in baseline BMI[b]	4	57.8
% difference from placebo (LSMean)		53.9
Body Weight[b]		
Baseline mean (kg)	102.6	109.9
% change from baseline (LSMean)[a]	2.7	-14.7
		-17.4

% difference from placebo (LSMean)		

LSMean = least squares mean; CI = confidence interval

[a] The intention-to-treat population includes all randomized patients. Missing data were imputed using available data according to value and timing of last available observation on treatment and endpoint's baseline value from retrieved subjects (RD-MI). At Week 68, the BMI was missing for 2.2% and 7.5% of patients randomized to WEGOVY injection and placebo, respectively.

[b] Parameters not included in the pre-specified hierarchical testing.

* p<0.0001 (unadjusted 2-sided) for superiority.

The time course of change in BMI with WEGOVY injection and placebo from baseline through Week 68 is depicted in **Figure 11**. The cumulative frequency distribution of change in BMI is shown in **Figure 12**.

Figure 11. Change from Baseline (%) in BMI in Pediatric Patients with Obesity Aged 12 Years and Older in Study 10 with WEGOVY Injection



Observed values for patients completing each scheduled visit, and estimates with multiple imputations from retrieved dropouts (RD-MI).

Figure 12. Change in BMI (%) from Baseline to Week 68 in Pediatric Patients with Obesity Aged 12 Years and Older in Study 10 with WEGOVY Injection



Observed data from in-trial period including imputed data for missing observations (RD-MI).

Effect of WEGOVY Injection on Anthropometry and Cardiometabolic Parameters in Pediatric Patients with Obesity Aged 12 Years and Older

Changes in waist circumference and cardiometabolic parameters with WEGOVY injection are shown in **Table 19** for the study in pediatric patients aged 12 years and older.

Table 19. Mean Changes in Anthropometry and Cardiometabolic Parameters in Pediatric Patients with Obesity Aged 12 Years and Older in Study 10 with WEGOVY Injection

	PLACEBO N=67		**WEGOVY Injection N=134**		
	Baseline	**Change from Baseline (LSMean)**	**Baseline**	**Change from Baseline (LSMean)**	**Difference from placebo (LSMean)**
Waist Circumference	107.3	-0.6	111.9	-12.7	-12.1

(cm)[2]					
Systolic Blood Pressure (mmHg)[2]	120	-0.8	120	-2.7	-1.9
Diastolic Blood Pressure (mmHg)[2]	73	-0.8	73	-1.4	-0.6
Heart Rate[3]	76	-2.3	79	1.2	3.5
HbA_{1c} (%)[2,4]	5.4	-0.1	5.5	-0.4	-0.2
	Baseline	**% Change from Baseline (LSMean)**	**Baseline**	**% Change from Baseline (LSMean)**	**Relative difference from placebo (LSMean)**
Total Cholesterol (mg/dL)[2,5]	160.1	-1.3	159.4	-8.3	-7.1
LDL Cholesterol (mg/dL)[2,5]	91.7	-3.6	89.8	-9.9	-6.6
HDL Cholesterol (mg/dL)[2,5]	43.3	3.2	43.7	8	4.7
Triglycerides (mg/dL)[2,5]	108	2.6	111.3	-28.4	-30.2

[1] Parameters listed in the table were not included in the pre-specified hierarchical testing.

[2] Missing data were imputed using available data according to value and timing of last available observation on treatment and endpoint's baseline value from retrieved subjects (RD-MI). Model based estimates based on an analysis of covariance model including treatment and stratification groups (gender, Tanner stage group) and the interaction between stratification groups as factors and baseline value as a covariate.

[3] Model based estimates based on a mixed model for repeated measures including treatment as a factor and baseline value as a covariate all nested within visit.

[4] For patients without type 2 diabetes at randomization (N=129 for WEGOVY injection-treated patients and N=64 for placebo-treated patients).

[5] Baseline value is the geometric mean.

14.4 Noncirrhotic Metabolic Dysfunction-associated Steatohepatitis with Moderate to Advanced Liver Fibrosis in Adults Treated with WEGOVY Injection

Overview of Clinical Trial

The efficacy of WEGOVY injection was evaluated based on an efficacy analysis at Week 72 in Study 11 (NCT#04822181), a 240-week, randomized, double-blind, placebo-controlled trial. Enrolled patients had a baseline or recent liver biopsy showing clinically significant MASLD (metabolic dysfunction-associated steatotic liver disease), defined as MASH with fibrosis stage 2 or 3 and a non-alcoholic fatty liver disease (NAFLD) Activity Score (NAS) ≥4 with a score of 1 or more in steatosis, lobular inflammation, and hepatocyte ballooning. Efficacy determination was based on the effect of WEGOVY injection on resolution of steatohepatitis without worsening of liver fibrosis and on at least one stage improvement in liver fibrosis without worsening of steatohepatitis, on post-baseline liver biopsies collected at 72 weeks.

The Week 72 analysis included 800 F2 and F3 (at eligibility) patients randomized 1:2 to

receive placebo (n=266) or WEGOVY once weekly (n=534), in addition to standard of care for cardiometabolic comorbidities and healthy lifestyle counseling. WEGOVY injection or matching placebo was escalated to 2.4 mg once weekly during the initial 16 weeks of the treatment period. Dose escalation could be prolonged or patients could remain at a lower dose if 2.4 mg once weekly was not tolerable.

Demographic and baseline characteristics were balanced between treatment and placebo groups. Overall, the median (Q1 to Q3) age of patients at baseline was 57 (49 to 65) years, 57% were female, 18% were Hispanic, 68% were White, 27% were Asian, and 0.6% were Black or African American. Median (Q1 to Q3) body mass index (BMI) was 34 (30 to 38) kg/m^2 and median (Q1 to Q3) body weight was 93 (79 to 110) kg. Baseline characteristics are presented in **Table 20**.

Table 20. Baseline Characteristics in Adults Patients with Noncirrhotic MASH with Stage 2 to Stage 3 Fibrosis in Study 11

Characteristic	**Overall (N=800)**
Fibrosis stage, n (%) F2 F3	250 (31) 550 (69)
Body Mass Index (BMI, kg/m^2), n (%)[a] <25 25-30 30-35 ≥35	53 (7) 164 (21) 252 (32) 330 (41)
Lean MASH, n (%)[b]	22 (3)
Type 2 Diabetes, n (%)	447 (56)
Hypertension, n (%)	503 (63)
Dyslipidemia, n (%)	198 (25)
Statin use, n (%)	300 (38)
Fibrosis Index Based on 4 Factors (FIB-4), Median (Q1, Q3)[a]	1.6 (1.1, 2.3)
Enhanced Liver Fibrosis (ELF), Median (Q1, Q3)	9.9 (9.3, 10.5)

[a] Less than 5% missingness in the variable is omitted.

[b] Lean MASH defined as BMI <25 kg/m^2 for non-Asian patients and BMI <23 kg/m^2 for Asian patients.

Among the 79% of the patients with vibration-controlled transient elastography (VCTE) at baseline, median (Q1 to Q3) VCTE was 10.9 (8.6 to 15.5) kPa, which may not be representative of the entire study population. The 21% of patients with missing VCTE at baseline had higher percentages of being female and having baseline diabetes, hypertension, and dyslipidemia.

Results

Table 21 presents the Week 72 histopathology primary endpoint results comparing WEGOVY injection with placebo on:

1) the estimated percentage of patients with resolution of steatohepatitis and no

worsening of liver fibrosis and
2) the estimated percentage of patients with at least one stage improvement in liver fibrosis and no worsening of steatohepatitis. The secondary endpoint results of the estimated percentage of patients with resolution of steatohepatitis and improvement in liver fibrosis at Week 72 are also presented. Two pathologists independently read the liver biopsies for each patient; a third pathologist performed adjudication if consensus could not be reached between the two pathologists. WEGOVY injection demonstrated improvement on these histopathology endpoints at Week 72 compared to placebo.

Table 21. Efficacy Results at Week 72 in Adult Patients with Noncirrhotic MASH with Stage 2 or

Stage 3 Fibrosis in Study 11 of WEGOVY Injection

	PLACEBO N=266	**WEGOVY Injection N=534**
Resolution of steatohepatitis and no worsening of liver fibrosis		
Response Rate (%)	34	63
Difference in response rate vs. placebo (95% CI)		29 (21, 36)*
Improvement in liver fibrosis and no worsening of steatohepatitis		
Response Rate (%)	22	37
Difference in response rate vs. placebo (95% CI)		14 (8, 21)*
Resolution of steatohepatitis and improvement in liver fibrosis		
Response Rate (%)	16	33
Difference in response rate vs. placebo (95% CI)		17 (10, 23)*

* Results were statistically significant.

Endpoints were evaluated according to the MASH Clinical Research Network (CRN). Resolution of steatohepatitis is defined as a score of 0 to 1 for lobular inflammation, 0 for ballooning, and any value for steatosis. No worsening of steatohepatitis is defined as no increase from baseline in score for ballooning, lobular inflammation, or steatosis.

Estimated using pooled Mantel-Haenszel (MH) estimates stratified by baseline type 2 diabetes status (presence or absence) and baseline fibrosis stage (F2 or F3) with missing data handled by reference-based multiple imputation and 95% confidence intervals (CIs) calculated using Rubin’s rule to pool Sato’s estimate of standard errors across the imputed datasets.

Another secondary endpoint was the percent change in body weight from baseline to Week 72. Patients treated with WEGOVY injection (mean baseline body weight 95.4 kg) achieved an average of 10.5% weight loss from baseline at Week 72, and patients treated with placebo (mean baseline weight 97.6 kg) achieved an average of 2% weight loss from baseline at Week 72; treatment with WEGOVY injection resulted in an average of 8.5% greater weight loss from baseline compared to placebo (95% CI: 7.4% to 9.5%).

Starting at Week 12 and through Week 72, there was a trend of greater reductions from baseline in average ALT and AST in the WEGOVY injection group as compared to the placebo group.

16 HOW SUPPLIED/STORAGE AND HANDLING

WEGOVY Injection

How Supplied

WEGOVY injection is a clear, colorless solution available in cartons containing 4 single-dose pens, 4 single-dose syringes with integrated needles, or 1 single-patient-use pen (WEGOVY FlexTouch) as follows:

Device Type	Name	Total Strength per Total Volume	NDC
Single-dose Pen	WEGOVY	0.25 mg/0.5 mL	0169-4525-14
Single-dose Pen		0.5 mg/0.5 mL	0169-4505-14
Single-dose Pen		1 mg/0.5 mL	0169-4501-14
Single-dose Pen		1.7 mg/0.75 mL	0169-4517-14
Single-dose Pen		2.4 mg/0.75 mL	0169-4524-14
Single-dose Pen	WEGOVY HD	7.2 mg/0.75 mL	0169-4572-14
Single-patient-use Pen (with 4 weekly doses of 2.4 mg)	WEGOVY FlexTouch	9.6 mg/3 mL (3.2 mg/mL)	0169-4915-32
Single-dose Syringe	WEGOVY	0.25 mg/0.5 mL	0169-1602-14
Single-dose Syringe		0.5 mg/0.5 mL	0169-1605-14
Single-dose Syringe		1 mg/0.5 mL	0169-1610-14
Single-dose Syringe		1.7 mg/0.75 mL	0169-1617-14
Single-dose Syringe		2.4 mg/0.75 mL	0169-1624-14

Recommended Storage

- Do not store WEGOVY in the freezer or directly adjacent to the refrigerator cooling element.
- Do not freeze WEGOVY and do not use if it has been frozen.

Single-dose Pen or Single-dose Syringe Storage

Store the WEGOVY prefilled single-dose pen or single-dose syringe in the refrigerator from 2°C to 8°C (36°F to 46°F). If needed, prior to the pen or syringe cap removal, the pen can be kept from 8°C to 30°C (46°F to 86°F) up to 28 days. Protect WEGOVY injection from light. WEGOVY injection must be kept in the original carton until time of administration. Discard the WEGOVY pen or syringe after use.

Single-patient-use Pen (WEGOVY FlexTouch) Storage

Prior to first use, store WEGOVY FlexTouch single-patient-use penin a refrigerator between 2°C to 8°C (36°F to 46°F). After first use, WEGOVY FlexTouch single-patient-use pencan be stored for 56 days between 15°C to 30°C (59°F to 86°F) or in a refrigerator 2°C to 8°C (36°F to 46°F). WEGOVY FlexTouch single-patient-use penshould be protected from excessive heat and light. Keep the pen cap on when not in use. Always remove and safely discard the needle after each injection. Store the WEGOVY FlexTouch single-patient-use pen without an injection needle attached. Always use a new needle for each injection.

WEGOVY Tablets

How Supplied

WEGOVY tablets are available as:

Tablet strength	**Description**	**Package Configuration**	**NDC Number**
1.5 mg	White to light yellow, round shaped debossed with “1.5” on one side and “novo” on the other side	Bottle of 30 tablets	0169-4415-31
4 mg	White to light yellow, round shaped debossed with “4” on one side and “novo” on the other side	Bottle of 30 tablets	0169-4404-31
9 mg	White to light yellow, round shaped debossed with “9” on one side and “novo” on the other side	Bottle of 30 tablets	0169-4409-31
25 mg	White to light yellow, oval shaped debossed with “25” on one side and “novo” on the other side	Bottle of 30 tablets	0169-4425-31

Recommended Storage

Store at 20°C to 25°C (68°F to 77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Store and dispense WEGOVY tablets in the original bottle. Store WEGOVY tablet in the original bottle until use to protect tablets from moisture. Store WEGOVY tablets in a dry place away from moisture.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide and Instructions for Use).

Risk of Thyroid C-cell Tumors

Inform patients that semaglutide causes thyroid C-cell tumors in rodents and that the human relevance of this finding has not been determined. Counsel patients to report symptoms of thyroid tumors (e.g., a lump in the neck, hoarseness, dysphagia, or dyspnea) to their physician *[see Boxed Warning, Warnings and Precautions (5.1)]*.

Acute Pancreatitis

Inform patients of the potential risk for acute pancreatitis and its symptoms: severe abdominal pain that sometimes radiates to the back, and which may or may not be accompanied by nausea or vomiting. Instruct patients to discontinue WEGOVY promptly and contact their physician if pancreatitis is suspected *[see Warnings and Precautions (5.2)]*.

Acute Gallbladder Disease

Inform patients of the risk of acute gallbladder disease. Advise patients that substantial or rapid weight loss can increase the risk of gallbladder disease, but that gallbladder disease may also occur in the absence of substantial or rapid weight loss. Instruct patients to contact their healthcare provider for appropriate clinical follow-up if

gallbladder disease is suspected *[see Warnings and Precautions (5.3)].*

Hypoglycemia

Inform patients of the risk of hypoglycemia and educate patients on the signs and symptoms of hypoglycemia. Advise patients with diabetes mellitus on glycemic lowering therapy that they may have an increased risk of hypoglycemia when using WEGOVY and to report signs and/or symptoms of hypoglycemia to their healthcare provider *[see Warnings and Precautions (5.4)]*.

Acute Kidney Injury due to Volume Depletion

Inform patients of the potential risk of acute kidney injury due to dehydration associated with gastrointestinal adverse reactions. Advise patients to take precautions to avoid fluid depletion. Inform patients of the signs and symptoms of acute kidney injury and instruct them to promptly report any of these signs or symptoms or persistent (or extended) nausea, vomiting, and diarrhea to their healthcare provider *[see Warnings and Precautions (5.5)]*.

Severe Gastrointestinal Adverse Reactions

Inform patients of the potential risk of severe gastrointestinal adverse reactions. Instruct patients to contact their healthcare provider if they have severe or persistent gastrointestinal symptoms *[see Warnings and Precautions (5.6)].*

Hypersensitivity Reactions

Inform patients that serious hypersensitivity reactions have been reported during postmarketing use of semaglutide, the active ingredient in WEGOVY. Advise patients on the symptoms of hypersensitivity reactions and instruct them to stop taking WEGOVY and seek medical advice promptly if such symptoms occur *[see Warnings and Precautions (5.7)]*.

Diabetic Retinopathy Complications in Patients with Type 2 Diabetes

Inform patients with type 2 diabetes to contact their physician if changes in vision are experienced during treatment with WEGOVY *[see Warnings and Precautions (5.8)]*.

Heart Rate Increase

Instruct patients to inform their healthcare providers of palpitations or feelings of a racing heartbeat while at rest during WEGOVY treatment *[see Warnings and Precautions (5.9)]*.

Pulmonary Aspiration During General Anesthesia or Deep Sedation

Inform patients that WEGOVY may cause their stomach to empty more slowly which may lead to complications with anesthesia or deep sedation during planned surgeries or procedures. Instruct patients to inform healthcare providers prior to any planned surgeries or procedures if they are taking WEGOVY *[see Warnings and Precautions (5.10)].*

Never Share a WEGOVY FlexTouch Pen Between Patients

Advise patients that they must never share a WEGOVY FlexTouch with another person, even if the needle is changed, because doing so carries a risk for transmission of blood-borne pathogens *[see Warnings and Precautions (5.11)]*.

Pregnancy

WEGOVY may cause fetal harm. Advise patients to inform their healthcare provider of a known or suspected pregnancy. Advise patients who are exposed to WEGOVY during pregnancy to contact Novo Nordisk at 1-877-390-2760 or www.wegovypregnancyregistry.com *[see Use in Specific Populations (8.1)]*.

Administration

Instruct patients how to prepare and administer the correct dose of WEGOVY and assess their ability to inject subcutaneously to ensure the proper administration of WEGOVY. Inform patients and their caregiver(s) which WEGOVY presentation (e.g., single-dose pen or syringe, WEGOVY FlexTouch single-patient-use pen) they will receive and ensure they receive training appropriate for that specific presentation. If the prescribed WEGOVY presentation changes, ensure patients and caregivers receive appropriate training and instruct them to consult the Instructions for Use for the newly prescribed presentation. Advise caregivers that WEGOVY FlexTouch is not recommended for self-administration by those with visual impairment unless their healthcare provider determines the patient can properly administer *[see Dosage and Administration (2.1)].*

Missed Doses

Inform patients if a dose of WEGOVY injection is missed and the next scheduled dose is more than 2 days away (48 hours), administer WEGOVY injection as soon as possible. If one dose is missed and the next scheduled dose is less than 2 days away (48 hours), do not administer the dose. Inform patients to resume their regular once-weekly dosing schedule *[see Dosage and Administration (2.4)].*

Inform patients if a dose of WEGOVY tablets is missed, skip the missed dose and take the next dose the following day *[see Dosage and Administration (2.4)]*.

Lactation

Advise females not to breastfeed during treatment with WEGOVY tablets *[see Use in Specific Populations (8.2)]*.

Marketed by: Novo Nordisk Inc., Plainsboro, NJ 08536

For additional information about WEGOVY contact:

Novo Nordisk Inc.

800 Scudders Mill Road

Plainsboro, NJ 08536

1-833-934-6891

WEGOVY® and FlexTouch® are registered trademarks of Novo Nordisk A/S.

Patent Information: http://www.novonordisk-us.com/products/product-patents.html

© 2026 Novo Nordisk

Medication Guide

MEDICATION GUIDE	
WEGOVY® (wee-GOH-vee) (semaglutide)	**WEGOVY® (wee-GOH-vee) (semaglutide)**

injection, for subcutaneous use **tablets, for oral use**

Do not share your pen, Wegovy FlexTouch or needles with other people, even if the needle has been changed. You may give other people a serious infection, or get a serious infection from them.

Read this Medication Guide and Instructions for Use before you start using WEGOVY and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or your treatment.

What is the most important information I should know about WEGOVY?
WEGOVY may cause serious side effects, including:

- **Possible thyroid tumors, including cancer**. Tell your healthcare provider if you get a lump or swelling in your neck, hoarseness, trouble swallowing, or shortness of breath. These may be symptoms of thyroid cancer. In studies with rodents, WEGOVY and other medicines that work like WEGOVY caused thyroid tumors, including thyroid cancer. It is not known if WEGOVY will cause thyroid tumors or a type of thyroid cancer called medullary thyroid carcinoma (MTC) in people.
- Do not use WEGOVY if you or any of your family have ever had a type of thyroid cancer called medullary thyroid carcinoma (MTC), or if you have an endocrine system condition called Multiple Endocrine Neoplasia syndrome type 2 (MEN 2).

What is WEGOVY?

- WEGOVY injection is a prescription medicine used with a reduced-calorie diet and increased physical activity to:
  - reduce the risk of major cardiovascular events such as death, heart attack, or stroke in adults with known heart disease and with either obesity or overweight.
  - help adults and children aged 12 years and older with obesity, or some adults with excess weight (overweight) who also have weight-related medical problems to lose weight and keep the weight off.
  - treat adults with metabolic dysfunction-associated steatohepatitis (MASH) with moderate to advanced liver scarring (fibrosis), but not with cirrhosis of the liver.
- WEGOVY tablets are a prescription medicine used with a reduced-calorie diet and increased physical activity to:
  - reduce the risk of major cardiovascular events such as death, heart attack, or stroke in adults with known heart disease and with either obesity or overweight.
  - help adults with obesity, or some adults with excess weight (overweight) who also have weight-related medical problems to lose weight and keep the weight off.
- WEGOVY contains semaglutide and should not be used with other semaglutide-containing products or other GLP-1 receptor agonist medicines.
- It is not known if WEGOVY injection is safe and effective:
  - to reduce the risk of major cardiovascular events (death, heart attack, or stroke) in people under 18 years.
  - to help children under 12 years of age lose weight and keep the weight off.
  - for the treatment of MASH in people under 18 years of age.
- It is not known if WEGOVY tablets are safe and effective for use in people under 18 years of age.

Do not use WEGOVY if:

- you or any of your family have ever had a type of thyroid cancer called MTC or if you have an endocrine system condition called MEN 2.
- you have had a serious allergic reaction to semaglutide or any of the ingredients in WEGOVY injection or WEGOVY tablets. See the end of this Medication Guide for a complete list of ingredients in WEGOVY injection and WEGOVY tablets. See "**What are the possible side effects of WEGOVY?**" for symptoms of a serious allergic reaction.

Before using WEGOVY, tell your healthcare provider if you have any other medical conditions, including if you:

- have or have had problems with your pancreas or kidneys.
- have type 2 diabetes and a history of diabetic retinopathy.
- are scheduled to have surgery or other procedures that use anesthesia or deep sleepiness (deep sedation).
- are pregnant or plan to become pregnant. WEGOVY may harm your unborn baby. You should stop using WEGOVY 2 months before you plan to become pregnant.
  - **Pregnancy Exposure Registry:** There is a pregnancy exposure registry for women who use WEGOVY during pregnancy. The purpose of this registry is to collect information about the health of you and your baby. Talk to your healthcare provider about how you can take part in this registry or you may contact Novo Nordisk at 1-877-390-2760 or www.wegovypregnancyregistry.com.
- are breastfeeding or plan to breastfeed. Breastfeeding is not recommended during treatment with WEGOVY tablets. It is not known if WEGOVY when received through an injection passes into your breast milk. You should talk with your healthcare provider about the best way to feed your baby while using WEGOVY.
  Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. WEGOVY may affect the way some medicines work and some medicines may affect the way WEGOVY works. Tell your healthcare provider if you are taking other medicines to treat diabetes, including sulfonylureas or insulin. WEGOVY slows stomach emptying and can affect medicines that need to pass through the stomach quickly.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

How should I use WEGOVY?

- Use WEGOVY exactly as your healthcare provider tells you to.
- Use WEGOVY with a reduced-calorie diet and increased physical activity.
- If you take too much WEGOVY, you may have severe nausea, severe vomiting and severe low blood sugar. Call your healthcare provider or Poison Help line at 1-800-222-1222 or go to the nearest hospital emergency room right away. Advice is also available online at poisonhelp.org.

WEGOVY Injection

- Read the **Instructions for Use** that comes with WEGOVY.
- **Your healthcare provider should tell you which WEGOVY product you will be using (for example – single-dose pen or syringe, single-patient-use pen, or tablets).** Before you use WEGOVY injection for the first time, your

healthcare provider should show you how to use it. Once you have been trained, you may inject WEGOVY yourself if your healthcare provider determines that you are able to do so safely. If you are visually impaired, the WEGOVY FlexTouch pen is not recommended for self-injection.

- If you are prescribed a different WEGOVY product at any point, make sure you receive new training for the new product and read the Instructions for Use that come with it.
- WEGOVY is injected under the skin (subcutaneously) of your stomach (abdomen), thigh, or upper arm. **Do not** inject WEGOVY into a muscle (intramuscularly) or vein (intravenously).
- Change (rotate) your injection site with each injection. **Do not** use the same site for each injection.
- **Use WEGOVY 1 time each week, on the same day each week, at any time of the day.**
- If you need to change the day of the week, you may do so as long as your last dose of WEGOVY was given **2** or more days before.
- If you miss a dose of WEGOVY and the next scheduled dose is more than 2 days away (48 hours), take the missed dose as soon as possible. If you miss a dose of WEGOVY and the next scheduled dose is less than 2 days away (48 hours), do not administer the dose. Take your next dose on the regularly scheduled day.
- If you miss doses of WEGOVY for 2 or more consecutive weeks, contact your healthcare provider to talk about how to restart your treatment.
- You can take WEGOVY with or without food.

WEGOVY Tablets

- Take 1 WEGOVY tablet by mouth on an empty stomach in the morning with water (no more than 4 ounces). **Do not** take WEGOVY tablets with any other liquids besides water.
- **Do not** split, crush, chew or dissolve WEGOVY tablets. Swallow WEGOVY tablets whole.
- **Do not** take more than 1 WEGOVY tablet each day.
- After 30 minutes, you can eat, drink, or take other oral medicines.
- If you miss a dose of WEGOVY, skip the missed dose and go back to your regular schedule for the next dose.

What are the possible side effects of WEGOVY?
WEGOVY may cause serious side effects, including:

- **See "What is the most important information I should know about WEGOVY?"**
- **inflammation of your pancreas (pancreatitis).** Stop using WEGOVY and call your healthcare provider right away if you have severe pain in your stomach area (abdomen) that will not go away, with or without nausea or vomiting. Sometimes you may feel the pain from your abdomen to your back.
- **gallbladder problems.** WEGOVY may cause gallbladder problems including gallstones. Some gallbladder problems need surgery. Call your healthcare provider if you have any of the following symptoms:

o pain in your upper stomach (abdomen)

o yellowing of skin or eyes (jaundice)

- o fever
- o clay-colored stools

- **increased risk of low blood sugar (hypoglycemia) in patients with type 2 diabetes, especially those who also take medicines to treat type 2 diabetes mellitus such as an insulin or a sulfonylureas.** Low blood sugar in patients with type 2 diabetes who receive WEGOVY can be both a serious and common side effect. Talk to your healthcare provider about how to recognize and treat low blood sugar. You should check your blood sugar before you start taking WEGOVY and while you take WEGOVY. Signs and symptoms of low blood sugar may include:

- o dizziness or light-headedness
- o sweating
- o shakiness
- o blurred vision
- o slurred speech
- o weakness
- o anxiety
- o hunger
- o headache
- o irritability or mood changes
- o confusion or drowsiness
- o fast heartbeat
- o feeling jittery

- **dehydration leading to kidney problems.** Diarrhea, nausea, and vomiting may cause a loss of fluids (dehydration) which may cause kidney problems. It is important for you to drink fluids to help reduce your chance of dehydration. Tell your healthcare provider right away if you have nausea, vomiting, or diarrhea that does not go away.
- **severe stomach problems.** Stomach problems, sometimes severe, have been reported in people who use WEGOVY. Tell your healthcare provider if you have stomach problems that are severe or will not go away.
- **serious allergic reactions.** Stop using WEGOVY and get medical help right away, if you have any symptoms of a serious allergic reaction including:

- o swelling of your face, lips, tongue or throat
- o severe rash or itching
- o very rapid heartbeat
- o problems breathing or swallowing
- o fainting or feeling dizzy

- **change in vision in people with type 2 diabetes.** Tell your healthcare provider if you have changes in vision during treatment with WEGOVY.
- **increased heart rate.** WEGOVY can increase your heart rate while you are at rest. Your healthcare provider should check your heart rate while you take

WEGOVY. Tell your healthcare provider if you feel your heart racing or pounding in your chest and it lasts for several minutes.

- **food or liquid getting into the lungs during surgery or other procedures that use anesthesia or deep sleepiness (deep sedation).** WEGOVY may increase the chance of food getting into your lungs during surgery or other procedures. Tell all your healthcare providers that you are taking WEGOVY before you are scheduled to have surgery or other procedures.

The most common side effects of WEGOVY in adults or children aged 12 years and older may include:

- nausea
- stomach (abdomen) pain
- dizziness
- gas
- diarrhea
- changes in skin sensations
- feeling bloated
- vomiting
- headache
- belching
- stomach flu
- constipation
- tiredness (fatigue)
- upset stomach
- low blood sugar in people with type 2 diabetes
- heartburn
- hair loss

Talk to your healthcare provider about any side effect that bothers you or does not go away. These are not all the possible side effects of WEGOVY.
Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store WEGOVY?
WEGOVY injection

- Do not store WEGOVY in the freezer or directly beside the refrigerator cooling element.
- **Do not freeze WEGOVY and do not use WEGOVY if it has been frozen.**
- **WEGOVY single-dose pen and single-dose syringe:**
  - Store in the refrigerator between 36°F to 46°F (2°C to 8°C).
  - If needed, before removing the pen or syringe cap, **WEGOVY single-dose pen** and **single-dose syringe** can be stored from 46°F to 86°F (8°C to 30°C) in the original carton for up to 28 days.
  - Keep **WEGOVY single-dose pen and syringe** in the original carton to protect it from light.
  - Throw away the **WEGOVY single-dose pen or syringe** if it has been frozen, has been exposed to light or temperatures above 86°F (30°C), has been out of the refrigerator for 28 days or longer, or after use.
- **WEGOVY FlexTouch single-patient-use pen:**
  - Store in the refrigerator between 36°F to 46°F (2°C to 8°C).
  - After first use, store WEGOVY FlexTouch pen between 59°F to 86°F (15°C to 30°C) or in the refrigerator between 36°F to 46°F (2°C to 8°C) for up to 56

days.
- Protect WEGOVY FlexTouch pen from excessive heat and light.
- Keep the pen cap on when not in use.
- Always remove and safely discard the needle after each injection.
- Store WEGOVY FlexTouch pen without an injection needle attached.
- Always use a new needle for each injection.
- Throw away if the WEGOVY FlexTouch pen if it has been frozen, has been exposed to light or temperatures above 86°F (30°C), or has been out of the refrigerator for 56 days or longer.

WEGOVY tablets

- Store WEGOVY tablets at room temperature between 68°F to 77°F (20°C to 25°C).
- Store tablets in the original closed WEGOVY bottle until you are ready to take one. Do not store in any other container.
- Store in a dry place away from moisture.

Keep WEGOVY and all medicines out of the reach of children.

General information about the safe and effective use of WEGOVY.
Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use WEGOVY for a condition for which it was not prescribed. Do not give WEGOVY to other people, even if they have the same condition that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about WEGOVY that is written for health professionals.

What are the ingredients in WEGOVY?
Active Ingredient: semaglutide
Inactive Ingredients in WEGOVY injection:disodium phosphate dihydrate, sodium chloride, water for injection. Phenol and propylene glycol are also inactive ingredients in the single-patient-use WEGOVY FlexTouch. Hydrochloric acid or sodium hydroxide may be added to adjust pH.
Inactive Ingredients in WEGOVY tablets: salcaprozate sodium (SNAC) and magnesium stearate

Marketed by: Novo Nordisk Inc., Plainsboro, NJ 08536
WEGOVY® and FlexTouch® are registered trademarks of Novo Nordisk A/S.
Patent Information: http://novonordisk-us.com/products/product-patents.html
© 2026 Novo Nordisk
For more information, go to startWegovy.com or call 1-833-Wegovy-1.

This Medication Guide has been approved by the U.S. Food and Drug Administration.
Revised: 06/2026

Instructions for Use – Pen

Instructions for Use
WEGOVY®
(semaglutide) injection
WEGOVY comes in five strengths:

0.25 mg / 0.5 mL | 0.5 mg / 0.5 mL | 1 mg / 0.5 mL | 1.7 mg / 0.75 mL | 2.4 mg / 0.75 mL

Before you use your WEGOVY pen for the first time, talk to your healthcare provider or your caregiver about how to prepare and inject WEGOVY correctly.



Important information
Read this Instructions for Use before you start using WEGOVY. This information does not replace talking to your healthcare provider about your medical condition or treatment.

- **Your WEGOVY pen is for 1 time use only**. The WEGOVY pen is for subcutaneous (under the skin) use only.
- **The dose of WEGOVY is already set on your pen.**
- **The needle is covered by the needle cover and the needle will not be seen.**
- **Do not** remove the pen cap until you are ready to inject.
- Do not touch or push on the needle cover. You could get a needle stick injury.
- Your WEGOVY injection will start when the needle cover is pressed firmly against your skin.
- **Do not** remove the pen from your skin before the yellow bar in the pen window has stopped moving. The medicine may appear on the skin or squirt from the needle and you may not get your full dose of WEGOVY if:
  - o the pen is removed too early **or**
  - o you have not pressed the pen firmly against the skin for the entire injection.
- If the yellow bar does not start moving or stops during the injection, contact your healthcare provider or Novo Nordisk at startWegovy.com or call Novo Nordisk Inc. at 1-833-934-6891.
- The needle cover will lock when the pen is removed from your skin. **You cannot stop the injection and restart it later.**
- People who are blind or have vision problems should not use the WEGOVY pen without help from a person trained to use the WEGOVY pen.

How do I store WEGOVY?

- Store the WEGOVY pen in the refrigerator between 36°F to

46°F (2°C to 8°C).

- If needed, before removing the pen cap, WEGOVY can be stored from 46°F to 86°F (8°C to 30°C) in the original carton for up to 28 days.
- Keep WEGOVY in the original carton to protect it from light.
- **Do not freeze**.
- Throw away the pen if WEGOVY has been frozen, has been exposed to light or temperatures above 86°F (30°C), or has been out of the refrigerator for 28 days or longer.

Keep WEGOVY and all medicines out of the reach of children.

How to use your WEGOVY pen	**Do not use your WEGOVY pen without receiving training from your healthcare provider.** Make sure that you or your caregiver know how to give an injection with the pen before you start your treatment.

Read and follow the instructions so that you use your WEGOVY pen

correctly:

Preparation

Step 1. Prepare for your injection.

- **Supplies you will need to give your WEGOVY injection:**
  - WEGOVY pen
  - 1 alcohol swab or soap and water
  - 1 gauze pad or cotton ball
  - 1 sharps disposable container for used WEGOVY pens



- **Wash your hands.**
- **Check your WEGOVY pen.**
  Do not use your WEGOVY pen if:
  - The pen appears to have been used or any part of the pen appears broken, for example if it has been dropped.
  - The WEGOVY medicine is not clear and colorless through the pen window.
  - The expiration date (EXP) has passed.

Contact Novo Nordisk at 1-833-934-6891 if your WEGOVY pen fails any of these checks.

Step 2. Choose your injection site.

- Your healthcare provider can help you choose the injection site that is best for you
  - You may inject into your upper leg (front of the thighs), lower stomach (keep 2 inches away from your belly button) or upper arm.
- Do not inject into an area where the skin is tender, bruised, red, or hard. Avoid injecting into areas with scars or stretch marks.
- You may inject in the same body area each week, but make sure it is not in the same spot each time.

Clean the injection site with an alcohol swab or soap and water. Do not touch the injection site after cleaning. Allow the skin to dry before injecting.



Injection
Step 3. Remove pen cap.

- **Pull the pen cap straight off your pen.**

Step 4. Inject WEGOVY.

- **Push the pen firmly against your skin and keep applying pressure until the yellow bar has stopped moving.**
  If the yellow bar does not start moving, press the pen more firmly against your skin.
- **You will hear 2 clicks during the injection.**
- Click 1: the injection has started.
- Click 2: the injection is ongoing.
- **If you have problems with the injection, refer to the "Troubleshooting" section.**



Throw away pen

Step 5. Throw away (dispose of) pen.

Safely dispose of the WEGOVY pen right away after each use. See **"How do I throw away (dispose of) WEGOVY pens?"**

- **What if blood appears after injection?**
  If blood appears at the injection site, press the site lightly with a gauze pad or cotton ball.

Troubleshooting

- **If you have problems injecting, change to a more firm injection site, such as upper leg, or upper arm or consider standing up while injecting into the lower stomach.**
- **If medicine appears on the skin or squirts from the needle, make sure the next time you inject to keep applying pressure until the yellow bar has stopped moving. Then you can lift the pen slowly from your skin.**



How do I throw away (dispose of) WEGOVY pens?

Put the used WEGOVY pen in an FDA-cleared sharps disposal container right away after use. **Do not throw away (dispose of)** the pen in your household trash.

If you do not have an FDA-cleared sharps disposal container, you may use a household container that is:

- made of a heavy-duty plastic,
- able to be closed with a tight-fitting, puncture-resistant lid, without sharps being able to come out,
- upright and stable during use,
- leak-resistant, and
- properly labeled to warn of hazardous waste inside the container.

When your sharps disposal container is almost full, you will need to follow your community guidelines for the right way to dispose of your sharps disposal container. There may be state or local laws about how you should throw away used needles and syringes. For more information about safe sharps disposal, and for specific sharps disposal in the state that you live in, go to the FDA's website at http://www.fda.gov/safesharpsdisposal.

- Do not reuse the pen.
- Do not recycle the pen or sharps disposal container, or throw them into household trash.

Important: Keep your WEGOVY pen, sharps disposal container and all medicines out of the reach of children.

How do I care for my pen?

Protect your pen

- Do not drop your pen or knock it against hard surfaces.
- Do not expose your pen to any liquids.
- If you think that your pen may be damaged, do not try to fix it. Use a new one.
- Keep the pen cap on until you are ready to inject. Your pen will no longer be sterile if you store an unused pen without the cap, if you pull the pen cap off and put it on again, or if the pen cap is missing. This could lead to an infection.

If you have any questions about WEGOVY, go to startWegovy.com or call Novo Nordisk Inc. at 1-833-Wegovy-1

Marketed by:
Novo Nordisk Inc.

Plainsboro, NJ 08536
For information about WEGOVY, go to startWegovy.com or contact:
Novo Nordisk Inc.
800 Scudders Mill Road
Plainsboro, NJ 08536
1-833-Wegovy-1
WEGOVY ® is a registered trademark of Novo Nordisk A/S.
Patent Information: http://novonordisk-us.com/products/product-patents.html
© 2025 Novo Nordisk

This Instructions for Use has been approved by the U.S. Food and Drug Administration. Revised: November 2025

Instructions for Use – WEGOVY HD

Instructions for Use
WEGOVY® HD
(semaglutide) injection
WEGOVY HD comes in the following strength:

7.2 mg/0.75 mL

Before you use your WEGOVY pen for the first time, talk to your healthcare provider or your caregiver about how to prepare and inject WEGOVY correctly.

Pull out to get started



Important information
Read this Instructions for Use before you start using WEGOVY. This information does not replace talking to your healthcare provider about your medical condition or treatment.

- **Your WEGOVY pen is for 1 time use only.** The WEGOVY pen is for subcutaneous (under the skin) use only.
- **The dose of WEGOVY is already set on your pen.**
- **The needle is covered by the needle cover and the needle will not be seen.**
- Do not remove the pen cap until you are ready to inject.
- Do not touch or push on the needle cover. You could get a needle stick injury.
- Your WEGOVY injection will start when the needle cover is pressed firmly against your skin.
- **Do not** remove the pen from your skin before the yellow bar in the pen window has stopped moving. The medicine may appear on the skin or squirt from the needle and you may not get your full dose of WEGOVY if:
- the pen is removed too early **or**
- you have not pressed the pen firmly against the skin for the entire injection.
- If the yellow bar does not start moving or stops during the injection, contact your healthcare provider or Novo Nordisk at startWegovy.com or call Novo Nordisk Inc. at 1-833-934-6891.
- The needle cover will lock when the pen is removed from your skin. **You cannot stop the injection and restart it later.**
- People who are blind or have vision problems should not use the WEGOVY pen without help from a person trained to use the WEGOVY pen.

How do I store WEGOVY?

- Store the WEGOVY pen in the refrigerator between 36°F to 46°F (2°C to 8°C).
- If needed, before removing the pen cap, WEGOVY can be stored from 46°F to 86°F (8°C to 30°C) in the original carton for up to 28 days.
- Do not store WEGOVY in the freezer or directly beside the refrigerator cooling element.
- **Do not freeze WEGOVY and do not use if it has been frozen.**
- Keep WEGOVY in the original carton to protect it from light.
- Throw away the pen if WEGOVY has been frozen, has been exposed to light or temperatures above 86°F (30°C), or has been out of the refrigerator for 28 days or longer or after use.

Keep WEGOVY and all medicines out of the reach of children.

WEGOVY pen parts



How to use your WEGOVY pen	**Do not use your WEGOVY pen without receiving training from your healthcare provider.** Make sure that you or your caregiver know how to give an injection with the pen before you start your treatment.

Read and follow the instructions so that you use your WEGOVY pen correctly:

Preparation

Step 1. Prepare for your injection.

- **Supplies you will need to give your WEGOVY injection:**
  - WEGOVY pen

- 1 alcohol swab or soap and water
- 1 gauze pad or cotton ball
- 1 sharps disposable container for used WEGOVY pens



- **Wash your hands.**
- **Check your WEGOVY pen.**
  Do not use your WEGOVY pen if:
- The pen appears to have been used or any part of the pen appears broken, for example if it has been dropped.
- The WEGOVY medicine is not clear and colorless through the pen window.
- The expiration date (EXP) has passed.

Contact Novo Nordisk at 1-833-934-6891 if your WEGOVY pen fails any of these checks.

Step 2. Choose your injection site.

- Your healthcare provider can help you choose the injection site that is best for you.
- You may inject into your upper leg (front of the thighs), lower stomach (keep 2 inches away from your belly button) or upper arm.
- Do not inject into an area where the skin is tender, bruised, red, or hard. Avoid injecting into areas with scars or stretch marks.
- You may inject in the same body area each week, but make sure it is not in the same spot each time.

Clean the injection site with an alcohol swab or soap and water. Do not touch the injection site after cleaning. Allow the skin to dry before injecting.



Injection
Step 3. Remove pen cap.

- **Pull the pen cap straight off your pen.**

Step 4. Inject WEGOVY.

- **Push the pen firmly against your skin and keep applying pressure until the yellow bar has stopped moving.**
  If the yellow bar does not start moving, press the pen more firmly against your skin.
- **You will hear 2 clicks during the injection.**
- Click 1: the injection has started.
- Click 2: the injection is ongoing.
- **If you have problems with the injection, refer to the "Troubleshooting" section.**



Throw away pen

Step 5. Throw away (dispose of) pen.

Safely dispose of the WEGOVY pen right away after each use. See **"How do I throw away (dispose of) WEGOVY pens?"**

- **What if blood appears after injection?**

If blood appears at the injection site, press the site lightly with a gauze pad or cotton

ball.

Troubleshooting

- **If you have problems injecting, change to a more firm injection site, such as upper leg, or upper arm or consider standing up while injecting into the lower stomach.**
- **If medicine appears on the skin or squirts from the needle, make sure the next time you inject to keep applying pressure until the yellow bar has stopped moving. Then you can lift the pen slowly from your skin.**



How do I throw away (dispose of) WEGOVY pens?
Put the used WEGOVY pen in an FDA-cleared sharps disposal container right away after use. **Do not throw away (dispose of)** the pen in your household trash.
If you do not have an FDA-cleared sharps disposal container, you may use a household container that is:

- made of a heavy-duty plastic,
- able to be closed with a tight-fitting, puncture-resistant lid, without sharps being able to come out,
- upright and stable during use,
- leak-resistant, and
- properly labeled to warn of hazardous waste inside the container.

When your sharps disposal container is almost full, you will need to follow your community guidelines for the right way to dispose of your sharps disposal container. There may be state or local laws about how you should throw away used needles and syringes. For more information about safe sharps disposal, and for specific sharps disposal in the state that you live in, go to the FDA's website at http://www.fda.gov/safesharpsdisposal.

- Do not reuse the pen.
- Do not recycle the pen or sharps disposal container, or throw them into household trash.

Important: Keep your WEGOVY pen, sharps disposal container and all medicines out of the reach of children.

⚠ **How do I care for my pen?**
Protect your pen

- Do not drop your pen or knock it against hard surfaces.
- Do not expose your pen to any liquids.
- If you think that your pen may be damaged, do not try to fix it. Use a new one.
- Keep the pen cap on until you are ready to inject. Your pen will no longer be sterile if you store an unused pen without the cap, if you pull the pen cap off and put it on again, or if the pen cap is missing. This could lead to an infection.

✆ If you have any questions about WEGOVY, go to startWegovy.com or call Novo Nordisk Inc. at 1-833-Wegovy-1

Marketed by:

Novo Nordisk Inc.
Plainsboro, NJ 08536
For information about WEGOVY, go to startWegovy.com
or contact:
Novo Nordisk Inc.
800 Scudders Mill Road
Plainsboro, NJ 08536
1-833-Wegovy-1
Wegovy® is a registered trademark of Novo Nordisk A/S.
Patent Information: http://novonordisk-us.com/products/product-patents.html
© 2026 Novo Nordisk

This Instructions for Use has been approved by the U.S. Food and Drug Administration. Revised: 03/2026

Instructions for Use – Manual Prefilled Syringe

INSTRUCTIONS FOR USE
WEGOVY® [wee-GOH-vee]
(semaglutide)
injection, subcutaneous only
single-dose prefilled syringe

Before using your prefilled syringe, always read these instructions carefully.
This leaflet instructs how to perform an injection under the skin (subcutaneous).
The prefilled syringe might be different than any other device you have used previously.
Before you use your prefilled syringe, talk to your healthcare provider about how to inject your prefilled syringe correctly.

Get to Know Your Prefilled Syringe
You should check the label on the carton and prefilled syringe to ensure it contains the correct medicine. Please make sure you have the correct dose prescribed by your healthcare provider.





Important Information You Need to Know Before Injecting with your Prefilled Syringe

- Use the prefilled syringe exactly the way your healthcare provider tells you to.

- The prefilled syringe contains 1 complete dose. It is for one-time use only.
- **Do not** use the prefilled syringe if the carton is damaged or if the tamper-evident is broken.
- **Do not** warm the prefilled syringe using any heat sources such as hot water or a microwave.
- **Do not** use the prefilled syringes if the expiration date on the label or carton has expired.
- **Do not** use the prefilled syringe if it has been frozen or thawed.

People who are blind or have vision problems should not use the prefilled syringe without help from a person who knows how to use the prefilled syringe.

Gather the Following Materials Before Your Injection

- 1 WEGOVY prefilled syringe
- alcohol wipe
- cotton ball or gauze
- sharps disposal container for used prefilled syringes

Not Included in Your Carton



Frequently Asked Questions

Do I inject the full volume of the prefilled syringe?

Yes, the prefilled syringe is for one-time (single) use and contains one complete dose.

What if the prefilled syringe is damaged?

Do not use a dropped, damaged or broken prefilled syringe. Take a new prefilled syringe for your injection.

What if the needle is broken?

Do not use a needle that appears to be bent, dropped, damaged or broken. Take a new prefilled syringe for your injection.

If you have any questions about WEGOVY, go to startWEGOVY.com or call Novo Nordisk Inc. at 1-833-Wegovy-1.

Prepare Your Prefilled Syringe

1. **Take One Prefilled Syringe**

- Wash and dry your hands.
- Open the carton.
- Gently take out one prefilled syringe by holding the glass barrel.
- Check the syringe label. Confirm you have the dose prescribed by your healthcare provider.



Do not hold or touch the prefilled syringe by the plunger rod or needle cap.

Keep the needle cap on until you are ready to inject.

2. **Inspect your Prefilled Syringe**
- Check the expiration date (EXP) on the label has not passed.
- **Check that the medicine in the prefilled syringe is clear and colorless.** If the medicine looks cloudy or contains particles, **do not** use the prefilled syringe.

It is normal to see air bubbles; they do not affect your dose. **Do not** try to remove bubbles.
Do not use the prefilled syringe if it appears to have been used, dropped, damaged or broken.



3. **Choose Your Injection Site**

Inject into the skin of:

- stomach area (abdomen) at least 2 inches from the belly button.
- thighs.
- back of upper arm (only with help from someone else).

Clean the injection site with an alcohol wipe and allow to dry. Your healthcare provider can help you choose the injection site that is best for you.
Avoid using the exact same spot as the last time between injections.



Inject Your Dose

4. **Pull Off the Needle Cap**
- Pull the needle cap straight off and throw it in a sharps disposal container.

You might see drops of medicine from the needle tip. This is normal and does not affect your dose. Continue with your injection right away.
Do not put the needle cap back on, to avoid needle stick injuries.



5. **Insert Your Prefilled Syringe**
- Pinch and hold a fold of skin at the injection site between your thumb and fingers. Hold the pinch throughout the injection.
- Fully insert the needle into the pinched skin at a 45-degree angle.

This is to inject under the skin (subcutaneous) and not the muscle.



6. **Inject Your Dose**
- Keep a 45-degree angle. With a constant pressure, gently push the plunger rod all the way down until the plunger stops.

When the plunger is at the bottom and the prefilled syringe is empty, your dose is complete.

7. **Remove Your Syringe**
- Pull out the empty syringe at a 45-degree angle. Your injection is complete.

If blood appears at the injection site, lightly press with a cotton ball or gauze over it, without rubbing.
Do not put the needle cap back on the syringe.



After Injection

8. **Dispose of (Throw Away) Your Used Syringe**
- Put the used syringe in an FDA-cleared sharps disposal container right away after use.
- If you do not have an FDA-cleared sharps disposal container, you may use a household container that is:
  - made of a heavy-duty plastic,
  - can be closed with a tight-fitting, puncture-resistant lid, without sharps being able to come out,
  - upright and stable during use,
  - leak-resistant, and
  - properly labeled to warn of hazardous waste inside the container.
- When your sharps disposal container is almost full, you will need to follow your community guidelines for the right way to dispose of your sharps disposal container. There may be state or local laws about how you should throw away used needles and syringes. For more information about safe sharps disposal, and for specific information about sharps disposal in the state that you live in, go to the FDA's website at: http://www.fda.gov/safesharpsdisposal.
- **Do not** dispose of your used sharps disposal container in your household trash unless your community guidelines permit this. **Do not** recycle your used sharps disposal container.

Safely dispose of your syringe that is out of date or no longer needed.
Keep your syringe and all medicines out of the reach of children.

How Should I Store My WEGOVY Prefilled Syringe?

- Store the unused WEGOVY prefilled syringes in the refrigerator between 36°F to 46°F (2°C to 8°C).
- If needed, the prefilled syringes can be stored out of the refrigerator between 46°F to 86°F (8°C to 30°C) for up to 28 days. Record the date your prefilled syringe was removed from the refrigerator and dispose of it if not used within 28 days.
- **Do not** freeze.
- Always keep the prefilled syringes in the original carton to protect from light.

Keep the prefilled syringe and all medicines out of the reach of children.

Marketed by: Novo Nordisk Inc. Plainsboro, NJ 08536 For information about WEGOVY, contact: Novo Nordisk Inc. 800 Scudders Mill Road Plainsboro, NJ 08536 1-833-Wegovy-1 *Wegovy® is a registered trademark of Novo Nordisk A/S.* Patent Information: http://www.novonordisk-us.com/products/product-patents.html © 2026 Novo Nordisk	

The Instructions for Use has been approved by the U.S Food and Drug Administration. Approved: 05/2026

Instructions for Use - FlexTouch

INSTRUCTIONS FOR USE
WEGOVY® (wee-GOH-vee) FlexTouch®
(semaglutide) injection, for subcutaneous use
2.4 mg dose single-patient-use pen
Each pen contains 4 fixed doses, one dose taken weekly

Unfold the whole leaflet and read these instructions carefully before using your WEGOVY FlexTouch single-patient-use pen.
This single-patient-use pen contains four doses of WEGOVY 2.4 mg. If you have been using a single-dose WEGOVY pen until now, please get familiar with your WEGOVY FlexTouch single-patient-use pen and needles.
Do not use your pen without proper training from your healthcare provider. Make sure that you know how to give yourself an injection with the pen before you start your treatment.
Do not share your WEGOVY pen with other people, even if the needle has been changed. You may give other people a serious infection or get a serious infection from them.
If you are blind or have poor eyesight and cannot read the dose counter on the pen, do not use this pen without help. Get help from a person with good eyesight who is trained to use the WEGOVYpen.

Please use the table inside the lid of the carton to keep track of how many injections you have used and how many doses remain in your pen.

Start by checking your pen to make sure that it contains WEGOVY, then look at the pictures in this 'Instructions for Use' to get to know the different parts of your pen and needle.

Your pen is a prefilled, single-patient-use, dial-a-dose pen. One pen contains 4 doses of 2.4 mg. You can only select doses of 2.4 mg . NovoFine® Plus 32G 4 mm disposable needles are included with your WEGOVY FlexTouch pen.

Always use a new needle for each injection.

If you need additional needles, NovoFine® Plus, NovoFine® or other compatible disposable needles with a length up to 8 mm can be used. For needle compatibility information, talk to your healthcare professional.

Supplies you will need to give your WEGOVY injection:

- WEGOVY FlexTouch pen 2.4 mg dose
- a new NovoFine Plus or NovoFine needle or other compatible needle
- alcohol swab
- cotton ball or gauze
- an FDA-cleared sharps disposal container for throwing away used WEGOVY pens and needles. See **Step 5 'After your injection'** for information concerning disposal.



Step 1. Prepare your pen with a new needle

Wash your hands with soap and water.
Check the name and colored label of your pen, to make sure that it contains your 2.4 mg dose of WEGOVY. This is especially important if you take more than 1 type of medicine.
Pull off the pen cap. See **Figure A**.



Check that the medicinein your pen is clear and colorless.
Look through the pen window. If the medicinelooks cloudy, do not use the pen. See **Figure B**.



Do not attach a new needle to your pen until you are ready to take your injection.
Take a new needle, and tear off the paper tab. See **Figure C**.



Push the needle straight onto the pen. Turn until it is on tight. See **Figure D**.



The needle is covered by 2 caps. You must remove both caps. If you forget to remove both caps, you will not inject any medicine.
Pull off the outer needle cap. **Do not** throw it away. See **Figure E**.
Pull off the inner needle cap and throw it away. See **Figure F**.
A drop of medicinemay appear at the needle tip. This is normal, but you must still check the medicine flow, if you use a new pen for the first time.
Never use a bent or damaged needle.
Always use a new needle for each injection.
This will prevent contamination, infection, leakage of medicine, and blocked needles leading to the wrong dose.



Do not reuse or share your needles with other people. You may give other people a serious infection or get a serious infection from them.



Step 2. Check the flow with each new pen

Before you inject your first dose with your new pen, check the medicineflow.
For your second, third and fourth dose with this pen, go to **Step 3**. **Set your dose.**

Turn the dose selector **until the dose counter shows the flow check symbol (" —)**.
See **Figure G**.
Make sure that the flow check symbol lines up with the dose pointer.



Hold the pen with the needle pointing up.
Press and hold in the dose button until the dose counter shows 0. The 0 must line up with the dose pointer. See **Figure H**.
A drop of medicinewill appear at the needle tip.
If no drop appears, check the flow again. **This should only be done twice.** If there is still no drop of medicine, change the needle and check the flow 1 more time.
Do not use the pen if a drop of medicinestill does not appear.
Always make sure that a drop of medicine appears at the needle tip before you use a new pen for the first time. This makes sure that the medicineflows.
If no drop of medicine appears, you will **not** inject any medicine, even though the dose counter may move.
This may mean that there is a blocked or damaged needle.
A small drop of medicine may remain at the needle tip, but it will not be injected.
Only check the medicineflow before your first injection with each new pen.



Step 3. Set your dose

Turn the dose selector until the **dose counter stops**, and it **shows your 2.4 mg dose**. See **Figure I**.



The dashed line ⋮ in the dose counter will guide you to your 2.4 mg dose. See **Figure J**.
The dose selector clicks differently when turned forward or backwards. You will hear a 'click' every time you turn the dose selector. **Do not set the dose by counting the number of clicks you hear.**
You can only select 2.4 mg for each dose. When your pen contains less than 2.4 mg, the dose counter stops before 2.4 mg is shown. There is not enough medicine for your dose injection.
When your prescribed dose of 2.4 mg lines up with the dose pointer, you have selected your dose. See **Figure I**.
If the dose counter stops before you reach your prescribed dose, see the section **'Do you have enough medicine?'**.



Do you have enough medicine?

If the dose counter stops before you reach your prescribed dose of 2.4 mg, there is not enough medicine left for a full dose. Throw away (dispose of) the pen and use a new WEGOVY pen.



Choose your injection site

Choose an injection site on your upper arms, stomach or upper legs. Keep a distance of 2 inches (5 cm) from your belly button. See image for locations.
You may inject in the same body area each week, but make sure it is not in the same spot used for the last injection.
Wipe the skin with an alcohol swab. Let the injection site dry before you inject your dose.



Step 4. Inject your dose

Insert the needle into your skin as your healthcare provider has shown you. See **Figure L**.
Make sure you can see the dose counter. Do not cover it with your fingers. This could stop the injection.



Press and hold down the dose button until the dose counter shows -0-. The -0- must line up with the dose pointer. You may then hear or feel a click. See **Figure M**.



Continue pressing the dose button while keeping the needle in your skin and count slowly to 6. See **Figure N**.
If the needle is removed earlier, you may see a stream of medicinecoming from the needle tip. If this happens, the full dose will not be delivered.



Remove the needle from your skin. You can then release the dose button. See **Figure O**.
If blood appears at the injection site, press the site lightly with a cotton ball or gauze. Do not rub the area.
You may see a drop of medicineat the needle tip after injecting. This is normal and does not affect your dose.
Always watch the dose counter to make sure you have injected your complete dose. Hold the dose button down until the dose counter shows -0-.
How to identify a blocked or damaged needle?
If -0- does not appear in the dose counter after continuously pressing the dose button, you may have used a blocked or damaged needle.
If this happens you have **not** received **any** medicine even though the dose counter has moved from the



original dose that you have selected.
How to handle a blocked needle?
Change the needle as described in **Step 1**: "**Prepare your pen with a new needle**". Make sure you select the full dose you need.
Never touch the dose counter when you inject. This can stop the injection.

Step 5. After your injection

Carefully unscrew the needle from the pen. Do not put the needle caps back on the needle, to avoid needle sticks. See **Figure P.**



Place the needle in an FDA-cleared sharps container right away to reduce the risk of needle sticks. See **Figure Q**.



Put the pen cap on your pen after each use to protect the medicinefrom light. See **Figure R**.



If you do not have an FDA-cleared sharps disposal container, follow a one-handed needle recapping method. Carefully slip the needle into the outer needle cap. Push the capped needle against a firm, vertical surface until you hear a "click." Twist off the capped needle from the pen and dispose of the needle in an FDA-cleared sharps disposal container as soon as possible or in a household container as listed in '**Disposing of used pens and needles'**. See **Figure S**.
Never try to put the inner needle cap back on the needle. You may stick yourself with the needle.
Always remove the needle from your pen. This prevents contamination, infection, leakage of medicine, and blocked needles leading to the wrong dose. If the needle is blocked, you will not inject any medicine.
Always throw away (dispose of) the needle after each injection.



Throwing away (disposing of) used pens and needles:

- After having injected all four doses, there may still be medicine left in the pen. The pen should be thrown away (disposed of).
- Put your used WEGOVY pen and needle in an FDA-cleared sharps disposal container right away after use.
- If you do not have an FDA-cleared sharps disposal container, you may use a household container that is:
    - made of a heavy-duty plastic
    - can be closed with a tight-fitting, puncture-resistant lid, without sharps being able to come out
    - upright and stable during use
    - leak-resistant
    - properly labeled to warn of hazardous waste inside the container.

When your sharps disposal container is almost full, you will need to follow your community guidelines for the right way to dispose of your sharps disposal container. There may be state or local laws about how you should throw away used needle and syringes. For more information about the safe sharps disposal, and for specific information about sharps disposal in the state that you live in, go to the FDA's website at: http://www.fda.gov/safesharpsdisposal.

- **Do not** dispose of your used sharps disposal container in your household trash unless your community guidelines permit this.
- **Do not** recycle your used sharps disposal container.
- Safely dispose of medicine that is out of date or no longer needed.

Important

Caregivers must **be very careful when handling used needles** to prevent needle sticks and cross infection.
Never use a syringe to withdraw medicinefrom your pen.
Always keep your pen and needles **out of reach of others**, **especially children**.
Do not share your pen or needles with anyone else. You may give an infection to them or get an infection from them.
Do not leave it in a car or other place where it can

get too hot or too cold.

Caring for your pen

Do not drop your pen or knock it against hard surfaces. If you drop it or suspect a problem, attach a new needle and check the medicine flow before you inject.
Do not try to repair your pen or pull it apart.
Do not expose your pen to dust, dirt, or liquid.
Do not wash, soak, or lubricate your pen. If necessary, clean it with mild detergent on a moistened cloth.
How should I store my WEGOVY pen?

- Store in the refrigerator between 36°F to 46°F (2°C to 8°C).
- After first use, store WEGOVY FlexTouch between 68°F to 77°F (20°C to 25°C) or in a refrigerator 36°F to 46°F (2°C to 8°C) for up to 56 days.
- WEGOVY should be protected from excessive heat and light.
- Keep the pen cap on when not in use.
- Always remove and safely discard the needle after each injection.
- Store the **single-patient-use** WEGOVY FlexTouch pen without an injection needle attached.
- Throw away if the **single-patient-use** WEGOVY FlexTouch pen has been frozen, has been exposed to light or temperatures above 86°F (30°C), or has been out of the refrigerator for 56 days or longer.

Marketed by:
Novo Nordisk Inc.
Plainsboro, NJ 08536
For information about WEGOVY, contact:
Novo Nordisk Inc.
800 Scudders Mill Road
Plainsboro, NJ 08536
1-833-wegovy-1
(se habla español)
© 2026 Novo Nordisk
WEGOVY®, FlexTouch®, NovoFine® and NovoFinePlus® are registered trademarks of Novo Nordisk A/S.
Patent Information: http://novonordisk-us.com/products/product-patents.html

This Instructions for Use has been approved by the U.S. Food and Drug Administration.
Revised 06//2026

Package/Label Display Panel – 0.25 mg Pen

NDC 0169-**4525**-14 List 452514

0.25 mg

wegovy®

(semaglutide) injection

0.25 mg/0.5 mL

Use Wegovy 1 time a week

4 Single-Dose Prefilled Pens

Each pen delivers a single dose of 0.25 mg semaglutide

For subcutaneous use only

Single-Dose only

Rx only

Contains: 4 Wegovy pens, Product Literature.

Dispense the enclosed Medication Guide to each patient.



Package/Label Display Panel – 0.5 mg Pen

NDC 0169-**4505**-14 List 450514

0.5 mg

wegovy®

(semaglutide) injection

0.5 mg/0.5 mL

Use Wegovy 1 time a week

4 Single-Dose Prefilled Pens

Each pen delivers a single dose of 0.5 mg semaglutide

For subcutaneous use only

Single-Dose only

Rx only

Contains: 4 Wegovy pens, Product Literature.

Dispense the enclosed Medication Guide to each patient.



Package/Label Display Panel – 1 mg Pen

NDC 0169-**4501**-14 List 450114

1 mg

wegovy®

(semaglutide) injection

1 mg/0.5 mL

Use Wegovy 1 time a week

4 Single-Dose Prefilled Pens

Each pen delivers a single dose of 1 mg semaglutide

For subcutaneous use only

Single-Dose only

Rx only

Contains: 4 Wegovy pens, Product Literature.

Dispense the enclosed Medication Guide to each patient.



Package/Label Display Panel – 1.7 mg Pen

NDC 0169-**4517**-14 List 451714

1.7 mg

wegovy®

(semaglutide) injection

1.7 mg/0.75 mL

Use Wegovy 1 time a week

4 Single-Dose Prefilled Pens

Each pen delivers a single dose of 1.7 mg semaglutide

For subcutaneous use only

Single-Dose only

Rx only

Contains: 4 Wegovy pens, Product Literature.

Dispense the enclosed Medication Guide to each patient.



Package/Label Display Panel – 2.4 mg Pen

NDC 0169-**4524**-14 List 452414

2.4 mg

wegovy®

(semaglutide) injection

2.4 mg/0.75 mL

Use Wegovy 1 time a week

4 Single-Dose Prefilled Pens

Each pen delivers a single dose of 2.4 mg semaglutide

For subcutaneous use only

Single-Dose only

Rx only

Contains: 4 Wegovy pens, Product Literature.

Dispense the enclosed Medication Guide to each patient.



PACKAGE/LABEL PRINCIPAL DISPLAY PANEL – WEGOVY HD – 7.2 mg Pen

NDC 0169-4572-14 List 457214

wegovy® 7.2 mg HD

(semaglutide) injection

7.2 mg/0.75 mL

Use Wegovy 1 time a week

Each pen delivers a single-dose of 7.2 mg semaglutide

For subcutaneous use only

Rx only

Dispense the enclosed Medication Guide to each patient.

Contains: 4 Single-Dose Prefilled Pens

NDC 0169-4572-14

List 457214

wegovy® HD 7.2 mg

(semaglutide) injection
7.2 mg/0.75 mL

Use Wegovy 1 time a week

Each pen delivers a single-dose of 7.2 mg semaglutide

For subcutaneous use only

Rx only

Dispense the enclosed Medication Guide to each patient.

Contains: 4 Single-Dose Prefilled Pens

novo nordisk®

Get support at **startWegovy.com**

7.2 mg

Push to open

wegovy® HD 7.2 mg
(semaglutide) injection
7.2 mg/0.75 mL
4 Single-Dose Prefilled Pens

wegovy® HD 7.2 mg
(semaglutide) injection
7.2 mg/0.75 mL
4 Single-Dose Prefilled Pens

GTIN/Serial No./EXPLOT:

00301694572140

wegovy® HD 7.2 mg
(semaglutide) injection
7.2 mg/0.75 mL
4 Single-Dose Prefilled Pens

Use as directed by your healthcare professional.
Recommended Dosage: See prescribing information.
Subcutaneous use only

REFRIGERATE – DO NOT FREEZE
Storage: Store in the refrigerator from 36°F to 46°F (2°C to 8°C) in original carton to protect from light. If needed, prior to cap removal, each single-dose pen can be kept unrefrigerated from 46°F to 86°F (8°C to 30°C) up to 28 days. **Protect from light. Do Not Freeze.**

Not a child resistant container. Keep out of reach of children.
If the seal is broken before first use, contact your pharmacist.

Use Wegovy 1 time a week
Write the **day** of the week you choose to inject

© 2026 Novo Nordisk A/S
Novo Nordisk Inc.
Plainsboro, NJ 08536
1-833-Wegovy-1
(Se habla español)
startWegovy.com

Marketed by:
Novo Nordisk Inc.
Plainsboro, NJ 08536

Product of Denmark

Ingredients: Each 0.75 mL of Wegovy contains 7.2 mg of semaglutide and the following inactive ingredients per 1 mL: disodium phosphate dihydrate, 1.42 mg; sodium chloride, 8.25 mg and water for injection. Hydrochloric acid or sodium hydroxide may be added to adjust pH.
Sterile.

3 01694 57214 0

Patent Information:
http://www.novonordisk-us.com/products/product-patents.html

Wegovy® is a registered trademark of Novo Nordisk A/S

Get support at
startWegovy.com

Or scan the QR code

Call 1-833-Wegovy-1

novo nordisk®

8-9535-31-301-1

22-004-30

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL – 0.25 mg Manual Prefilled Syringe

NDC 0169-1602-14 List 160214

wegovy® 0.25 mg

(semaglutide) injection

0.25 mg/0.5 mL

Use Wegovy 1 time a week

For subcutaneous use only

Each prefilled syringe delivers 0.25 mg semaglutide

Rx only

Dispense the enclosed Medication Guide to each patient.

Dispense in this sealed carton.

Contains: 4 x 0.5 mL Single-Dose Prefilled Syringes

Push to open



PACKAGE/LABEL PRINCIPAL DISPLAY PANEL – 0.5 mg Manual Prefilled Syringe

NDC 0169-1605-14 List 160514

wegovy® 0.5 mg

(semaglutide) injection

0.5 mg/0.5 mL

Use Wegovy 1 time a week

For subcutaneous use only

Each prefilled syringe delivers 0.5 mg semaglutide

Rx only

Dispense the enclosed Medication Guide to each patient.

Dispense in this sealed carton.

Contains: 4 x 0.5 mL Single-Dose Prefilled Syringes

Push to open



PACKAGE/LABEL PRINCIPAL DISPLAY PANEL – 1 mg Manual Prefilled Syringe

NDC 0169-1610-14 List 161014

wegovy® 1 mg

(semaglutide) injection

1 mg/0.5 mL

Use Wegovy 1 time a week

For subcutaneous use only

Each prefilled syringe delivers 1 mg semaglutide

Rx only

Dispense the enclosed Medication Guide to each patient.

Dispense in this sealed carton.

Contains: 4 x 0.5 mL Single-Dose Prefilled Syringes

Push to open

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL – 1.7 mg Manual Prefilled Syringe

NDC 0169-1617-14 List 161714

wegovy® 1.7 mg

(semaglutide) injection

1.7 mg/0.5 mL

Use Wegovy 1 time a week

For subcutaneous use only

Each prefilled syringe delivers 1.7 mg semaglutide

Rx only

Dispense the enclosed Medication Guide to each patient.

Dispense in this sealed carton.

Contains: 4 x 0.5 mL Single-Dose Prefilled Syringes

Push to open

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL – 2.4 mg Manual Prefilled Syringe

NDC 0169-1624-14 List 162414

wegovy® 2.4 mg

(semaglutide) injection

2.4 mg/0.5 mL

Use Wegovy 1 time a week

For subcutaneous use only

Each prefilled syringe delivers 2.4 mg semaglutide

Rx only

Dispense the enclosed Medication Guide to each patient.

Dispense in this sealed carton.

Contains: 4 x 0.5 mL Single-Dose Prefilled Syringes

Push to open

Package/Label Display Panel - FlexTouch

NDC 0169-4915-32

wegovy® FlexTouch® Rx only

(semaglutide) injection

9.6 mg/3 mL (3.2 mg/mL)

Use Wegovy FlexTouch Once Weekly

For Single-Patient-Use Only

For subcutaneous use only

Dispense the enclosed Medication Guide to each patient.

Contains: 1 single-patient-use pen, 4 NovoFine Plus 32G needles



Package/Label Display Panel – 1.5 mg Tablets

NDC 0169-4415-31 LIST 441531
wegovy® 1.5 mg

(semaglutide) Tablets

1.5 mg

Rx only

DISPENSE WITH MEDICATION GUIDE

30 tablets

Package/Label Display Panel – 4 mg Tablets

NDC 0169-4404-31 LIST 440431
wegovy® 4 mg

(semaglutide) Tablets

4 mg

Rx only

DISPENSE WITH MEDICATION GUIDE

30 tablets

Package/Label Display Panel – 9 mg Tablets

NDC 0169-4409-31 LIST 440931
wegovy® 9 mg

(semaglutide) Tablets

9 mg

Rx only

DISPENSE WITH MEDICATION GUIDE

30 tablets

Package/Label Display Panel – 25 mg Tablets

NDC 0169-4425-31 LIST 442531
wegovy® 25 mg

(semaglutide) Tablets

25 mg

Rx only

DISPENSE WITH MEDICATION GUIDE

30 tablets

WEGOVY

semaglutide injection, solution

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0169-4525
Route of Administration	SUBCUTANEOUS		

Active Ingredient/Active Moiety		
Ingredient Name	**Basis of Strength**	**Strength**
SEMAGLUTIDE (UNII: 53AXN4NNHX) (SEMAGLUTIDE - UNII:53AXN4NNHX)	SEMAGLUTIDE	0.25 mg in 0.5 mL

Inactive Ingredients	
Ingredient Name	**Strength**
SODIUM PHOSPHATE, DIBASIC, DIHYDRATE (UNII: 94255I6E2T)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
WATER (UNII: 059QF0KO0R)	

Other Ingredients		
Ingredient Kind	**Ingredient Name**	**Quantity**
May contain	**HYDROCHLORIC ACID** (UNII: QTT17582CB)	
May contain	**SODIUM HYDROXIDE** (UNII: 55X04QC32I)	

Packaging				
#	**Item Code**	**Package Description**	**Marketing Start Date**	**Marketing End Date**
1	NDC:0169-4525-14	4 in 1 CARTON	06/05/2021	
1	NDC:0169-4525-01	0.5 mL in 1 SYRINGE, PLASTIC; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		
2	NDC:0169-4525-94	4 in 1 CARTON	06/05/2021	
2	NDC:0169-4525-90	0.5 mL in 1 SYRINGE, PLASTIC; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA215256	06/05/2021	

WEGOVY

semaglutide injection, solution

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0169-4505
Route of Administration	SUBCUTANEOUS		

Active Ingredient/Active Moiety		
Ingredient Name	**Basis of Strength**	**Strength**
SEMAGLUTIDE (UNII: 53AXN4NNHX) (SEMAGLUTIDE - UNII:53AXN4NNHX)	SEMAGLUTIDE	0.5 mg in 0.5 mL

Inactive Ingredients	
Ingredient Name	**Strength**
SODIUM PHOSPHATE, DIBASIC, DIHYDRATE (UNII: 94255I6E2T)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
WATER (UNII: 059QF0KO0R)	

Other Ingredients		
Ingredient Kind	**Ingredient Name**	**Quantity**
May contain	**HYDROCHLORIC ACID** (UNII: QTT17582CB)	

May contain	**SODIUM HYDROXIDE** (UNII: 55X04QC32I)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0169-4505-14	4 in 1 CARTON	06/05/2021	
1	NDC:0169-4505-01	0.5 mL in 1 SYRINGE, PLASTIC; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA215256	06/05/2021	

WEGOVY

semaglutide injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0169-4501
Route of Administration	SUBCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SEMAGLUTIDE (UNII: 53AXN4NNHX) (SEMAGLUTIDE - UNII:53AXN4NNHX)	SEMAGLUTIDE	1.0 mg in 0.5 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM PHOSPHATE, DIBASIC, DIHYDRATE (UNII: 94255I6E2T)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
WATER (UNII: 059QF0KO0R)	

Other Ingredients

Ingredient Kind	Ingredient Name	Quantity
May contain	**HYDROCHLORIC ACID** (UNII: QTT17582CB)	
May contain	**SODIUM HYDROXIDE** (UNII: 55X04QC32I)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0169-4501-14	4 in 1 CARTON	06/05/2021	

1	NDC:0169-4501-01	0.5 mL in 1 SYRINGE, PLASTIC; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA215256	06/05/2021	

WEGOVY

semaglutide injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0169-4517
Route of Administration	SUBCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SEMAGLUTIDE (UNII: 53AXN4NNHX) (SEMAGLUTIDE - UNII:53AXN4NNHX)	SEMAGLUTIDE	1.7 mg in 0.75 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM PHOSPHATE, DIBASIC, DIHYDRATE (UNII: 94255I6E2T)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
WATER (UNII: 059QF0KO0R)	

Other Ingredients

Ingredient Kind	Ingredient Name	Quantity
May contain	**HYDROCHLORIC ACID** (UNII: QTT17582CB)	
May contain	**SODIUM HYDROXIDE** (UNII: 55X04QC32I)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0169-4517-14	4 in 1 CARTON	06/05/2021	
1	NDC:0169-4517-01	0.5 mL in 1 SYRINGE, PLASTIC; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA215256	06/05/2021	

WEGOVY

semaglutide injection, solution

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0169-4524
Route of Administration	SUBCUTANEOUS		

Active Ingredient/Active Moiety		
Ingredient Name	**Basis of Strength**	**Strength**
SEMAGLUTIDE (UNII: 53AXN4NNHX) (SEMAGLUTIDE - UNII:53AXN4NNHX)	SEMAGLUTIDE	2.4 mg in 0.75 mL

Inactive Ingredients	
Ingredient Name	**Strength**
SODIUM PHOSPHATE, DIBASIC, DIHYDRATE (UNII: 94255I6E2T)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
WATER (UNII: 059QF0KO0R)	

Other Ingredients		
Ingredient Kind	**Ingredient Name**	**Quantity**
May contain	**HYDROCHLORIC ACID** (UNII: QTT17582CB)	
May contain	**SODIUM HYDROXIDE** (UNII: 55X04QC32I)	

Packaging				
#	**Item Code**	**Package Description**	**Marketing Start Date**	**Marketing End Date**
1	NDC:0169-4524-14	4 in 1 CARTON	06/05/2021	
1	NDC:0169-4524-01	0.75 mL in 1 SYRINGE, PLASTIC; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA215256	06/05/2021	

WEGOVY

semaglutide injection, solution

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0169-4572

Route of Administration	SUBCUTANEOUS		

Active Ingredient/Active Moiety		
Ingredient Name	**Basis of Strength**	**Strength**
SEMAGLUTIDE (UNII: 53AXN4NNHX) (SEMAGLUTIDE - UNII:53AXN4NNHX)	SEMAGLUTIDE	7.2 mg in 0.75 mL

Inactive Ingredients	
Ingredient Name	**Strength**
SODIUM PHOSPHATE, DIBASIC, DIHYDRATE (UNII: 94255I6E2T)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
WATER (UNII: 059QF0KO0R)	

Other Ingredients		
Ingredient Kind	**Ingredient Name**	**Quantity**
May contain	**HYDROCHLORIC ACID** (UNII: QTT17582CB)	
May contain	**SODIUM HYDROXIDE** (UNII: 55X04QC32I)	

Packaging				
#	**Item Code**	**Package Description**	**Marketing Start Date**	**Marketing End Date**
1	NDC:0169-4572-14	4 in 1 CARTON	03/19/2026	
1	NDC:0169-4572-01	0.75 mL in 1 SYRINGE, PLASTIC; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA215256	06/05/2021	

WEGOVY

semaglutide injection, solution

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0169-1602
Route of Administration	SUBCUTANEOUS		

Active Ingredient/Active Moiety		
Ingredient Name	**Basis of Strength**	**Strength**
SEMAGLUTIDE (UNII: 53AXN4NNHX) (SEMAGLUTIDE - UNII:53AXN4NNHX)	SEMAGLUTIDE	0.5 mg in 1 mL

Inactive Ingredients	
Ingredient Name	**Strength**
SODIUM PHOSPHATE, DIBASIC, DIHYDRATE (UNII: 94255I6E2T)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
WATER (UNII: 059QF0KO0R)	

Other Ingredients		
Ingredient Kind	**Ingredient Name**	**Quantity**
May contain	**HYDROCHLORIC ACID** (UNII: QTT17582CB)	
May contain	**SODIUM HYDROXIDE** (UNII: 55X04QC32I)	

Packaging				
#	**Item Code**	**Package Description**	**Marketing Start Date**	**Marketing End Date**
1	NDC:0169-1602-14	4 in 1 CARTON	12/31/2030	
1	NDC:0169-1602-01	0.5 mL in 1 SYRINGE, GLASS; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		
2	NDC:0169-1602-94	4 in 1 CARTON	12/31/2030	
2	NDC:0169-1602-90	0.5 mL in 1 SYRINGE, GLASS; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA215256	06/05/2021	

WEGOVY

semaglutide injection, solution

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0169-1605
Route of Administration	SUBCUTANEOUS		

Active Ingredient/Active Moiety		
Ingredient Name	**Basis of Strength**	**Strength**
SEMAGLUTIDE (UNII: 53AXN4NNHX) (SEMAGLUTIDE - UNII:53AXN4NNHX)	SEMAGLUTIDE	1 mg in 1 mL

Inactive Ingredients	
Ingredient Name	**Strength**
SODIUM PHOSPHATE, DIBASIC, DIHYDRATE (UNII: 94255I6E2T)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	

WATER (UNII: 059QF0KO0R)	

Other Ingredients		
Ingredient Kind	**Ingredient Name**	**Quantity**
May contain	**HYDROCHLORIC ACID** (UNII: QTT17582CB)	
May contain	**SODIUM HYDROXIDE** (UNII: 55X04QC32I)	

Packaging				
#	**Item Code**	**Package Description**	**Marketing Start Date**	**Marketing End Date**
1	NDC:0169-1605-14	4 in 1 CARTON	12/31/2030	
1	NDC:0169-1605-01	0.5 mL in 1 SYRINGE, GLASS; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA215256	06/05/2021	

WEGOVY

semaglutide injection, solution

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0169-1610
Route of Administration	SUBCUTANEOUS		

Active Ingredient/Active Moiety		
Ingredient Name	**Basis of Strength**	**Strength**
SEMAGLUTIDE (UNII: 53AXN4NNHX) (SEMAGLUTIDE - UNII:53AXN4NNHX)	SEMAGLUTIDE	2 mg in 1 mL

Inactive Ingredients	
Ingredient Name	**Strength**
SODIUM PHOSPHATE, DIBASIC, DIHYDRATE (UNII: 94255I6E2T)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
WATER (UNII: 059QF0KO0R)	

Other Ingredients		
Ingredient Kind	**Ingredient Name**	**Quantity**
May contain	**HYDROCHLORIC ACID** (UNII: QTT17582CB)	
May contain	**SODIUM HYDROXIDE** (UNII: 55X04QC32I)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0169-1610-14	4 in 1 CARTON	12/31/2030	
1	NDC:0169-1610-01	0.5 mL in 1 SYRINGE, GLASS; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA215256	06/05/2021	

WEGOVY

semaglutide injection, solution

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0169-1617
Route of Administration	SUBCUTANEOUS		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
SEMAGLUTIDE (UNII: 53AXN4NNHX) (SEMAGLUTIDE - UNII:53AXN4NNHX)	SEMAGLUTIDE	2.27 mg in 1 mL

Inactive Ingredients	
Ingredient Name	Strength
SODIUM PHOSPHATE, DIBASIC, DIHYDRATE (UNII: 94255I6E2T)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
WATER (UNII: 059QF0KO0R)	

Other Ingredients		
Ingredient Kind	Ingredient Name	Quantity
May contain	**HYDROCHLORIC ACID** (UNII: QTT17582CB)	
May contain	**SODIUM HYDROXIDE** (UNII: 55X04QC32I)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0169-1617-14	4 in 1 CARTON	12/31/2030	
1	NDC:0169-1617-01	0.75 mL in 1 SYRINGE, GLASS; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA215256	06/05/2021	

WEGOVY

semaglutide injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0169-1624
Route of Administration	SUBCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SEMAGLUTIDE (UNII: 53AXN4NNHX) (SEMAGLUTIDE - UNII:53AXN4NNHX)	SEMAGLUTIDE	3.2 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM PHOSPHATE, DIBASIC, DIHYDRATE (UNII: 94255I6E2T)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
WATER (UNII: 059QF0KO0R)	

Other Ingredients

Ingredient Kind	Ingredient Name	Quantity
May contain	**HYDROCHLORIC ACID** (UNII: QTT17582CB)	
May contain	**SODIUM HYDROXIDE** (UNII: 55X04QC32I)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0169-1624-14	4 in 1 CARTON	12/31/2030	
1	NDC:0169-1624-01	0.75 mL in 1 SYRINGE, GLASS; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA215256	06/05/2021	

WEGOVY

semaglutide injection, solution

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0169-4915
Route of Administration	SUBCUTANEOUS		

Active Ingredient/Active Moiety		
Ingredient Name	**Basis of Strength**	**Strength**
SEMAGLUTIDE (UNII: 53AXN4NNHX) (SEMAGLUTIDE - UNII:53AXN4NNHX)	SEMAGLUTIDE	3.2 mg in 1 mL

Inactive Ingredients	
Ingredient Name	**Strength**
SODIUM PHOSPHATE, DIBASIC, DIHYDRATE (UNII: 94255I6E2T)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
WATER (UNII: 059QF0KO0R)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Other Ingredients		
Ingredient Kind	**Ingredient Name**	**Quantity**
May contain	**SODIUM HYDROXIDE** (UNII: 55X04QC32I)	
May contain	**PHENOL** (UNII: 339NCG44TV)	
May contain	**PROPYLENE GLYCOL** (UNII: 6DC9Q167V3)	

Packaging				
#	**Item Code**	**Package Description**	**Marketing Start Date**	**Marketing End Date**
1	NDC:0169-4915-03	1 in 1 CARTON	12/31/2026	
1	NDC:0169-4915-32	3 mL in 1 SYRINGE, GLASS; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA215256	06/05/2021	

WEGOVY

semaglutide tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0169-4404
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
SEMAGLUTIDE (UNII: 53AXN4NNHX) (SEMAGLUTIDE - UNII:53AXN4NNHX)	SEMAGLUTIDE	4 mg

Inactive Ingredients	
Ingredient Name	Strength
SALCAPROZATE SODIUM (UNII: 1YTW0422YU)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics			
Color	WHITE (Light Yellow)	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	4;novo
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0169-4404-31	30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/06/2026	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA218316	01/06/2026	

WEGOVY

semaglutide tablet

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0169-4415
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
SEMAGLUTIDE (UNII: 53AXN4NNHX) (SEMAGLUTIDE - UNII:53AXN4NNHX)	SEMAGLUTIDE	1.5 mg

Inactive Ingredients	
Ingredient Name	**Strength**
SALCAPROZATE SODIUM (UNII: 1YTW0422YU)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics			
Color	WHITE (Light Yellow)	**Score**	no score
Shape	ROUND	**Size**	7mm
Flavor		**Imprint Code**	1;5;novo
Contains			

Packaging				
#	**Item Code**	**Package Description**	**Marketing Start Date**	**Marketing End Date**
1	NDC:0169-4415-31	30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/06/2026	
2	NDC:0169-4415-92	30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/06/2026	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA218316	01/06/2026	

WEGOVY

semaglutide tablet

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0169-4409
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	**Basis of Strength**	**Strength**
SEMAGLUTIDE (UNII: 53AXN4NNHX) (SEMAGLUTIDE - UNII:53AXN4NNHX)	SEMAGLUTIDE	9 mg

Inactive Ingredients	
Ingredient Name	**Strength**
SALCAPROZATE SODIUM (UNII: 1YTW0422YU)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics			
Color	WHITE (Light Yellow)	**Score**	no score

Shape	ROUND	**Size**	7mm
Flavor		**Imprint Code**	9;novo
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0169-4409-31	30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/06/2026	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA218316	01/06/2026	

WEGOVY

semaglutide tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0169-4425
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SEMAGLUTIDE (UNII: 53AXN4NNHX) (SEMAGLUTIDE - UNII:53AXN4NNHX)	SEMAGLUTIDE	25 mg

Inactive Ingredients

Ingredient Name	Strength
SALCAPROZATE SODIUM (UNII: 1YTW0422YU)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics

Color	WHITE (Light Yellow)	**Score**	no score
Shape	OVAL	**Size**	12mm
Flavor		**Imprint Code**	25;novo
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0169-4425-31	30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/06/2026	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA218316	01/06/2026	

Labeler - Novo Nordisk Pharmaceutical Industries, LP (622920320)

Establishment

Name	Address	ID/FEI	Business Operations
Novo Nordisk A/S - 25a-B		309477891	MANUFACTURE(0169-4915)

Revised: 6/2026 Novo Nordisk Pharmaceutical Industries, LP