

**URIMAR-T- methenamine, sodium phosphate, monobasic, monohydrate, phenyl salicylate, methylene blue, and hyoscyamine sulfate tablet**  
**Scite Pharma, LLC**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*

-----  
**URIMAR-T™**

**Rx Only**

**DESCRIPTION**

URIMAR-T™ for oral administration.

---

**Each tablet contains:**

Methenamine	120 mg
Sodium Phosphate Monobasic	40.8 mg
Phenyl Salicylate	36.2 mg
Methylene Blue	10.8 mg
Hyoscyamine Sulfate	0.12 mg

---

**Inactive ingredients:** colloidal silicon dioxide, lactose, polyethylene glycol, crospovidone, magnesium stearate, FD&C Lake Blue #1.

**METHENAMINE**

[100-97-0] 1,3,5,7-Tetraazatricyclo [3.3.1.-1<sup>3,7</sup>] decane; hexamethylenetetramine; HMT; HMTA; hexamine; 1,3,5,7-tetraazaadamantane hexamethylenimine; Uritone; Urotropin. C<sub>6</sub>H<sub>12</sub>N<sub>4</sub>; mol wt 140.19; C 51.40%, H 8.63%, N 39.96%. Methenamine (hexamethylenetetramine) exists as colorless, lustrous crystals or white crystalline powder. Its solutions are alkaline to litmus. Freely soluble in water, soluble in alcohol and in chloroform.

**SODIUM PHOSPHATE MONOBASIC**

[7558-80-7] Phosphoric acid sodium salt (1:1); Sodium biphosphate; sodium dihydrogen phosphate; acid sodium phosphate; monosodium orthophosphate; primary sodium phosphate; H<sub>2</sub>NaO<sub>4</sub>P; mol wt 119.98, H 1.68%, Na 19.16%, O 53.34%, P 25.82%. Monohydrate, white, odorless slightly deliquesce crystals or granules. At 100° C loses all its water; when ignited it converts to metaphosphate. It is freely soluble in water and practically insoluble in alcohol. The aqueous solution is acid. pH of 0.1 molar aqueous solution at 25° C: 4.5.

**PHENYL SALICYLATE**

[118-55-8] 2-Hydroxybenzoic acid phenyl ester; Salol.  $C_{13}H_{10}O_3$ ; mol wt 214.22, C 72.89%, H 4.71%, O 22.41%. Made by the action of phosphorus oxy-chloride on a mixture of phenol and salicylic acid. Phenyl Salicylate exists as white crystals with a melting point of 41°-43° C. It is very slightly soluble in water and freely soluble in alcohol.

## **METHYLENE BLUE**

[61-73-4] 3,7-Bis(dimethylamino) phenothiazin-5-ium chloride; C.I. Basic Blue 9; methylthioninium chloride; tetramethylthionine chloride; 3,7-bis(dimethylamino) phenazathionium chloride.  $C_{16}H_{18}ClN_3S$ ; mol wt 319.85, C 60.08%, H 5.67%, Cl 11.08%, N 13.14%, S 10.03%. Methylene Blue (Methylthionine chloride) exists as dark green crystals. It is soluble in water and in chloroform; sparingly soluble in alcohol.

## **HYOSCYAMINE SULFATE**

[620-61-1] [3(S)-endo]- $\alpha$ -(Hydroxymethyl)-benzeneacetic acid 8-methyl-8-azabicyclo[3.2.1]oct-3-yl ester sulfate(2:1)(salt); 1 $\alpha$ H,5 $\alpha$  H-tropan-3  $\alpha$ -ol(-)-tropate (ester) sulfate(2:1)(salt); 3 $\alpha$ -tropanyl S-(-)-tropate; l-tropic acid ester with tropine; l-tropine tropate.  $C_{34}H_{48}N_2O_{10}S$ . Hyoscyamine Sulfate is an alkaloid of belladonna. Exists as a white crystalline powder. Its solutions are alkaline to litmus. Affected by light, it is slightly soluble in water; freely soluble in alcohol; sparingly soluble in ether.

## **CLINICAL PHARMACOLOGY**

METHENAMINE degrades in an acidic urine environment releasing formaldehyde which provides bactericidal or bacteriostatic action. It is well absorbed from the gastrointestinal tract. 70 to 90% reaches the urine unchanged at which point it is hydrolyzed if the urine is acidic. Within 24 hours it is almost completely (90%) excreted; of this amount at pH 5, approximately 20% is formaldehyde. Protein binding - some formaldehyde is bound to substances in the urine and surrounding tissues. Methenamine is freely distributed to body tissue and fluids but is not clinically significant as it does not hydrolyze at a pH greater than 6.8.

SODIUM PHOSPHATE MONOBASIC an acidifier, helps to maintain an acid pH in the urine necessary for the degradation of methenamine.

PHENYL SALICYLATE releases salicylate, a mild analgesic for pain.

METHYLENE BLUE possesses weak antiseptic properties. It is well absorbed by the gastrointestinal tract and is rapidly reduced to leukomethylene blue which is stabilized in some combination form in the urine. 75% is excreted unchanged.

HYOSCYAMINE SULFATE is a parasympatholytic drug which relaxes smooth muscles and thus produces an antispasmodic effect. It is well absorbed from the gastrointestinal tract and is rapidly distributed throughout the body tissues. Most is excreted in the urine within 12 hours, 13% to 50% being unchanged. Protein binding for hyoscyamine sulfate is moderate and biotransformation is hepatic.

## **INDICATIONS AND USAGE**

URIMAR-T™ is indicated for the treatment of symptoms of irritative voiding. Indicated for the relief of local symptoms, such as inflammation, hypermotility, and pain, which accompany lower urinary tract infections. Indicated for the relief of urinary tract symptoms caused by diagnostic procedures.

## **CONTRAINDICATIONS**

Urimar-T™ is contraindicated in patients with a hypersensitivity to any of the ingredients. Risk-benefit should be considered when the following medical problems exist: Cardiac disease (especially cardiac arrhythmias, congestive heart failure, coronary heart disease, and mitral stenosis); gastrointestinal tract obstructive disease; glaucoma; myasthenia gravis; acute urinary retention may be precipitated in obstructive uropathy (such as bladder neck obstruction due to prostatic hypertrophy).

## **WARNINGS**

Do not exceed recommended dosage. If rapid pulse, dizziness, or blurring of vision occurs, **discontinue use immediately.**

**KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.**

**IN CASE OF ACCIDENTAL OVERDOSE, SEEK PROFESSIONAL ASSISTANCE OR CONTACT A POISON CONTROL CENTER IMMEDIATELY.**

## **PRECAUTIONS**

**Urimar-T contains methylene blue and should NOT be taken with serotonergic psychiatric medications.**

### **Cross sensitivity and/or related problems**

Patients intolerant of other belladonna alkaloids or other salicylates may be intolerant of this medication also. Delay in gastric emptying could complicate the management of gastric ulcers.

### **Pregnancy**

Reproduction (FDA Pregnancy Category C)

Hyoscyamine and methenamine cross the placenta. Studies have not been done in either animals or humans. It is not known whether URIMAR-T™ tablets can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. URIMAR-T™ tablets should be given to a pregnant woman only if clearly needed.

### **Nursing mothers**

Methenamine and traces of hyoscyamine are excreted in breast milk. Caution should be exercised when URIMAR-T™ tablets are administered to a nursing mother.

### **Prolonged use**

There have been no studies to establish the safety of prolonged use in humans. No

known long-term animal studies have been performed to evaluate carcinogenic potential.

## **Pediatric**

Infants and young children are especially susceptible to the toxic effect of the belladonna alkaloids.

## **Geriatric**

Use with caution in elderly patients as they may respond to the usual doses of the belladonna alkaloids with excitement, agitation, drowsiness, or confusion.

## **DRUG INTERACTIONS**

### **Drug Interactions**

Although the exact mechanism of this drug interaction is unknown, methylene blue inhibits the action of monoamine oxidase A— an enzyme responsible for breaking down serotonin in the brain. It is believed that when methylene blue is given to patients taking serotonergic psychiatric medications, high levels of serotonin can build up in the brain, causing toxicity. This is referred to as Serotonin Syndrome. Signs and symptoms of Serotonin Syndrome include mental changes (confusion, hyperactivity, memory problems), muscle twitching, excessive sweating, shivering or shaking, diarrhea, trouble with coordination, and/or fever.

### **Additional Information for Healthcare Professionals**

Methylene blue can interact with serotonergic psychiatric medications and cause serious CNS toxicity.

In emergency situations requiring life-threatening or urgent treatment with methylene blue (as described above), the availability of alternative interventions should be considered and the benefit of methylene blue treatment should be weighed against the risk of serotonin toxicity. If methylene blue must be administered to a patient receiving a serotonergic drug, the serotonergic drug must be immediately stopped, and the patient should be closely monitored for emergent symptoms of CNS toxicity for two weeks (five weeks if fluoxetine [Prozac] was taken), or until 24 hours after the last dose of methylene blue, whichever comes first.

In non-emergency situations when non-urgent treatment with methylene blue is contemplated and planned, the serotonergic psychiatric medication should be stopped to allow its activity in the brain to dissipate. Most serotonergic psychiatric drugs should be stopped at least 2 weeks in advance of methylene blue treatment. Fluoxetine (Prozac), which has a longer half-life compared to similar drugs, should be stopped at least 5 weeks in advance.

Treatment with the serotonergic psychiatric medication may be resumed 24 hours after the last dose of methylene blue.

Serotonergic psychiatric medications should not be started in a patient receiving methylene blue. Wait until 24 hours after the last dose of methylene blue before starting the antidepressant.

Educate your patients to recognize the symptoms of serotonin toxicity or CNS toxicity and advise them to contact a healthcare professional immediately if they experience any symptoms while taking serotonergic psychiatric medications or methylene blue.

As a result of hyoscyamine's effects on gastrointestinal motility and gastric emptying, absorption of other oral medications may be decreased during concurrent use with this combination medication.

#### Urinary alkalizers and thiazide diuretics

May cause the urine to become alkaline reducing the effectiveness of methenamine by inhibiting its conversion to formaldehyde.

#### Antimuscarinics

Concurrent use may intensify antimuscarinic effects of hyoscyamine because of secondary antimuscarinic activities of these medications.

#### Antacids/antidiarrheals

Concurrent use may reduce absorption of hyoscyamine resulting in decreased therapeutic effectiveness. Concurrent use with antacids may cause urine to become alkaline reducing the effectiveness of methenamine by inhibiting its conversion to formaldehyde. Doses of these medications should be spaced 1 hour apart from doses of hyoscyamine.

#### Antimyasthenics

Concurrent use with hyoscyamine may further reduce intestinal motility, therefore, caution is recommended.

Ketoconazole and hyoscyamine may cause increased gastrointestinal pH.

Concurrent administration with hyoscyamine may result in marked reduction in the absorption of ketoconazole. Patients should be advised to take this combination at least 2 hours after ketoconazole.

#### Monoamine oxidase (MAO) inhibitors

Concurrent use with hyoscyamine may intensify antimuscarinic side effects.

Opioid (narcotic) analgesics may result in increased risk of severe constipation.

#### Sulfonamides

These drugs may precipitate with formaldehyde in the urine increasing the danger of crystalluria.

Patients should be advised that the urine and/or stools may become blue to blue-green as a result of the excretion of methylene blue.

## **ADVERSE REACTIONS**

Cardiovascular - rapid pulse, flushing

Central Nervous System - blurred vision, dizziness, drowsiness

Respiratory - shortness of breath or troubled breathing

Genitourinary - difficult micturition, acute urinary retention

Gastrointestinal - dry mouth, nausea and vomiting

Serious allergic reactions to this drug are rare. Seek immediate medical attention if you notice symptoms of a serious allergic reaction, including itching, rash, severe dizziness, swelling or trouble breathing.

This medication can cause urine and sometimes stools to turn blue to blue-green. This effect is harmless and will subside after medication is stopped.

**Call your doctor or physician for medical advice about side effects. To report SUSPECTED ADVERSE REACTIONS, contact Scite Pharma, LLC at 1-866-633-9033, or FDA at 1-800-FDA-1088, [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

### **Drug Abuse And Dependence**

A dependence on the use of URIMAR-T™ has not been reported and due to the nature of its ingredients, abuse of URIMAR T™ is not expected.

### **OVERDOSAGE**

Emesis or gastric lavage. Slow intravenous administration of physostigmine in doses of 1 to 4 mg (0.5 to 1 mg in children) repeated as needed in one to two hours to reverse severe antimuscarinic symptoms.

Administration of small doses of diazepam to control excitement and seizures.

Artificial respiration with oxygen if needed for respiratory depression.

Adequate hydration.

Symptomatic treatment as necessary.

If overdose is suspected, contact the poison control center at 1-800-222-1222, or your local emergency room immediately.

### **DOSAGE AND ADMINISTRATION**

**Adults:** One tablet orally 4 times per day followed by liberal fluid intake.

**Pediatric:** Dosage must be individualized by physician for older children. Not recommended for use in children six years of age or younger.

### **HOW SUPPLIED**

**Urimar-T™** are blue to blue-violet speckled, capsule-shaped tablets imprinted with 334 on one side and plain on the other, available in bottles of 100 tablets, (NDC: 79043-334-01), and sample tablets of 1, (NDC: 79043-334-99).

## **STORAGE**

Store in a cool, dry place at controlled room temperature 15° to 30°C (59° to 86°F). Keep container tightly closed. Protect from moisture and direct sunlight.

Dispense in a tight, light-resistant container as defined in the USP/NF with a child resistant closure.

**WARNING: Keep this and all drugs out of reach of children.**

Rx Only

Manufactured for:  
Scite Pharma, LLC  
Canton, MS 39046

Rev 02/2021

## **PRINCIPAL DISPLAY PANEL - 100 Tablet Bottle Label**

NDC 79043-334-01

URIMAR-T™

URINARY ANTISEPTIC

Each tablet contains:

Methenamine 120 mg

Sodium Phosphate Monobasic 40.8 mg

Phenyl Salicylate 36.2 mg

Methylene Blue 10.8 mg

Hyoscyamine Sulfate 0.12 mg

scite

PHARMA

Rx Only

Contents: 100 Tablets

**Dosage:** Adults-one tablet orally 4 times per day followed by liberal fluid intake. Pediatric-Dosage must be individualized by physician for older children. Not recommended for use in children six years of age or younger.

**SEE PRODUCT LITERATURE FOR FULL PRESCRIBING INFORMATION**

**Storage:** Store in a cool, dry place at controlled room temperature 15° to 30°C (59° to 86°F). Dispense in a tight, light-resistant container as defined in the USP/NF with a child resistant closure.

**KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.**

**NDC 79043-334-01**

**URIMAR-T<sup>™</sup>**  
URINARY ANTISEPTIC

**Each tablet contains:**  
Methenamine ..... 120 mg  
Sodium Phosphate Monobasic 40.8 mg  
Phenyl Salicylate ..... 36.2 mg  
Methylene Blue ..... 10.8 mg  
Hyoscyamine Sulfate ..... 0.12 mg

**scite PHARMA** Rx Only

**Contents: 100 Tablets**

**Precaution:** Contains Methylene Blue and should NOT be taken with serotonergic psychiatric medications.

**Note:** Patient should be advised that urine may be colored blue to blue-green while taking this medication.

Tamper-evident seal under cap. **Do not use if seal is broken or missing.**

Manufactured for:  
Scite Pharma, LLC  
Canton, MS 39046  
Rev 02/2021

**N 3 79043 33401 5**

## URIMAR-T

methenamine, sodium phosphate, monobasic, monohydrate, phenyl salicylate, methylene blue, and hyoscyamine sulfate tablet

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:79043-334
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Methenamine</b> (UNII: J50OIX95QV) (Methenamine - UNII:J50OIX95QV)	Methenamine	120 mg
<b>SODIUM PHOSPHATE, MONOBASIC, MONOHYDRATE</b> (UNII: 593YOG76RN) (PHOSPHATE ION - UNII:NK08V8K8HR, SODIUM CATION - UNII:LYR4M0NH37)	SODIUM PHOSPHATE, MONOBASIC, MONOHYDRATE	40.8 mg
<b>PHENYL SALICYLATE</b> (UNII: 28A37T47QO) (PHENYL SALICYLATE - UNII:28A37T47QO)	PHENYL SALICYLATE	36.2 mg
<b>METHYLENE BLUE</b> (UNII: T42P99266K) (Methylene Blue Cation - UNII:ZMZ79891ZH)	METHYLENE BLUE	10.8 mg
<b>HYOSCYAMINE SULFATE</b> (UNII: F2R8V82B84) (HYOSCYAMINE - UNII:PX44XO846X)	HYOSCYAMINE SULFATE	0.12 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>LACTOSE, UNSPECIFIED FORM</b> (UNII: J2B2A4N98G)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6130)	
<b>CROSPVIDONE</b> (UNII: 2S7830E561)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	

**Product Characteristics**

<b>Color</b>	BLUE	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	15mm
<b>Flavor</b>		<b>Imprint Code</b>	334
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79043-334-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	03/22/2021	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
Unapproved drug other		07/22/2005	

**Labeler** - Scite Pharma, LLC (117555106)

Revised: 3/2021

Scite Pharma, LLC