# ACETAMINOPHEN- acetaminophen tablet Cardinal Health 107, LLC

-----

### Acetaminophen 325 mg

#### **Drug Facts**

### Active ingredient (in each tablet)

Acetaminophen 325 mg

#### Purpose

Pain reliever/ Fever reducer

#### Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold
- toothache
- premenstrual and menstrual cramps
- temporarily reduces fever

**Liver warning:** This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg) in 24 hours for adults or 5 tablets (1,625 mg) in 24 hours for children. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### **Do not use** $\square$

 with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

#### Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

### **Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away.(1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. **This Unit Dose package is not child** 

resistant and is Intended for Institutional Use Only.

#### **Directions**

- do not take more than directed (see overdose warning)
- adults and children 12 years and over:
- take 2 tablets every 4 to 6 hours while symptoms last
- do not take more than 10 tablets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor
- children 6 years to under 12 years
  - o take 1 tablet every 4 to 6 hours while symptoms last
- do not take more than 5 tablets in 24 hours
- do not use for more than 5 days unless directed by a doctor
- children under 6 years: ask a doctor

#### Other information

Store in a dry place at  $15^{\circ}$  -  $30^{\circ}$ C ( $59^{\circ}$  -  $86^{\circ}$ F).

Povidone, pregelatinized starch, sodium starch glycolate, stearic acid.

### Questions or comments?

#### 1-800-231-4670

### Keep this carton for complete product information

Tamper Evident: Do not use if sealed blister units are broken or damaged.

Distributed by: MAJOR® PHARMACEUTICALS

Indianapolis, IN 46268

(800) 616-2471

www.majorpharmaceuticals.com

**Distributed By:** 

**Cardinal Health** 

Dublin, OH 43017

L5509849-10524

L5509849-20524

### **Principal Display Panel**

Acetaminophen

Regular Strength Tablets 325mg

Pain Reliever/ Fever Reducer

10 Tablets



X119

#### **ACETAMINOPHEN** Regular Strength Tablets 325mg Pain Reliever/Fever Reducer

10 TABLETS

**Drug Facts** 

Active ingredient (in each tablet)

Acetaminophen 325 mg

Pain reliever/ Fever

Inactive ingredients Povidone, pregelatinized starch, sodium starch glycolate, stearic acid.

- Uses Temporarily relieves minor aches and pains due to:

   headache muscular aches backache minor pain of arthritis
- the common cold toothache premenstrual and menstrual cramps

temporarily reduces fever

Directions

- do not take more than directed (see overdose warning)

- adults and children 12 years and over:

  take 2 tablets every 4 to 6 hours while symptoms last

  do not take more than 10 tablets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor
- children 6 years to under 12 years

  take 1 tablet every 4 to 6 hours while symptoms last
- do not take more than 5 tablets in 24 hours
- do not use for more than 5 days unless directed by a doctor

children under 6 years: ask a doctor

Other information

Store in a dry place at 15° - 30°C (59° - 86°F).

WARNINGS: This Unit Dose package is not child resistant and is Intended for Institutional Use Only. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Tamper Evident: Do not use if sealed blister units are broken or damaged.

Questions or comments? 1-800-231-4670 Distributed By: MAJOR® PHARMACEUTICALS Indianapolis, IN 46268 (800) 616-2471 www.majorpharmaceuticals.com

Distributed by Cardinal Health Dublin, OH 43017

L5509849-10524

Exp: Lot:

Distributed by Cardinal Health Dublin, OH 43017 L5509849-20524

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg) in 24 hours for adults or 5 tablets (1,625 mg) in 24 hours for children.

Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
   child takes more than 5 doses in 24 hours,
- which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day

while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening blisters rash
If a skin reaction occurs, stop use and seek medical help right

Do not use with any other drug containing acetamino (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a docto or pharmacist. If you are allergic to acetaminophen or any of the inactive ingredients in this product Ask a doctor before use if you have liver disease Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
   pain gets worse or lasts more than 5 days in children
- under 12 years me fever gets worse or lasts more than 3 days
- new symptoms occur redness or swelling is present These could be signs of a serious condition

### **ACETAMINOPHEN**

acetaminophen tablet

#### **Product Information**

**Product Type** HUMAN OTC DRUG Item Code (Source) NDC:55154-7899(NDC:0904-6773)

**Route of Administration ORAL** 

#### **Active Ingredient/Active Moiety**

**Basis of Strength Strength Ingredient Name** ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII: 36209ITL9D) **ACETAMINOPHEN** 325 mg

#### **Inactive Ingredients** Strength Ingredient Name

**SODIUM STARCH GLYCOLATE TYPE A POTATO** (UNII: 5856|3G2A2)

STARCH, CORN (UNII: O8232NY3SJ) POVIDONE K30 (UNII: U725QWY32X)

STEARIC ACID (UNII: 4ELV7Z65AP)

Product Characteristics							
Color	white	Score	2 pieces				
Shape	ROUND	Size	10mm				
Flavor		Imprint Code	CPC;220				
Contains							

P	Packaging						
#	Item Code	Package Description	Marketing Start Date	Marketing End Date			
1	NDC:55154- 7899-0	1 in 1 BAG	10/23/2018				
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product					

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC Monograph Drug	M013	10/23/2018			

## Labeler - Cardinal Health 107, LLC (118546603)

Revised: 5/2024 Cardinal Health 107, LLC