

**MAXIMUM STRENGTH NON DROWSY DAYTIME SEVERE COLD AND FLU-  
acetaminophen, dextromethorphan hydrobromide, guaifenesin,  
phenylephrine hydrochloride capsule, liquid filled  
DOLGENCORP, INC.**

-----

**Maximum Strength Non Drowsy Daytime Severe Cold and Flu**

***Drug Facts***

***Active ingredients (in each caplet)***

Acetaminophen  
Dextromethorphan HBr 10 mg  
Guaifenesin 200 mg  
Phenylephrine HCl 5 mg

***Purpose***

Pain reliever/fever reducer  
Cough suppressant  
Expectorant  
Nasal decongestant

***Uses***

- for the temporary relief of the following cold/flu symptoms: • minor aches and pains • headache • sore throat • nasal congestion • cough
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive • temporarily reduces fever

***Warnings***

**Liver warning:** This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours • with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. • if you have ever had an allergic reaction to this product or any of its ingredients.

**Ask a doctor before use if you have**

• liver disease • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to an enlarged prostate gland • persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema • cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

**When using this product do not exceed recommended dosage**

**Stop use and ask a doctor if** • nervousness, dizziness, or sleeplessness occur • pain, nasal congestion or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms

**Directions**

- do not take more than directed

---

adults and children • take 2 caplets every 4 hours • swallow whole; do not crush, chew  
12 years & over or dissolve • do not take more than 10 caplets in 24 hours  
children under 12 years ask a doctor

---

**Other information**

store at 25°C (77°F); excursions permitted between 15°–30°C (59°–86°F)

**Inactive Ingredients**

FD&C Yellow # 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan, titanium dioxide

## Questions or comments?

1-888-333-9792

## Principal Display Panel

Compare to Vicks® DayQuil® Severe Cold & Flu active ingredients†

Maximum Strength Cold & Flu

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER

DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT

GUAIFENESIN / EXPECTORANT

PHENYLEPHRINE HCl / NASAL DECONGESTANT

Relief of:

Relieves All Headache, Body Pain,  
Fever, Sore Throat, Chest Congestion,  
Cough, Nasal Congestion, Sinus  
Congestion, Sinus Pressure

20 Softgels

**Compare to the active ingredients of Maximum Strength Mucinex® Fast-Max® Cold & Flu ALL IN ONE\***

**For Ages 12 & Over**  
**Maximum Strength**  
**Mucus Relief**  
**Cold & Flu**  
Acetaminophen/Pain Reliever/Fever Reducer  
Dextromethorphan HBr/Cough Suppressant  
Guaifenesin/Expectorant  
Phenylephrine HCl/Nasal Decongestant

**TEMPORARILY RELIEVES**

- Headache
- Sore Throat
- Nasal Congestion
- Body Pain
- Chest Congestion
- Sinus Congestion
- Fever
- Cough
- Sinus Pressure

**Dissolves Quickly**

**20 Softgels**  
Each Softgel Box

**Drug Facts**  
**Active ingredients (in each softgel)**

Active ingredients (in each softgel)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 500 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

**Drug Facts (continued)**

**Uses**

- Relieves these common cold and flu symptoms:
  - Headache
  - Sore throat
  - Nasal congestion
  - Body pain
  - Chest congestion
  - Sinus congestion
  - Fever
  - Cough
  - Sinus pressure
- Helps loosen phlegm (mucus) and thin bronchial secretions to help the natural passageways of the lungs breathe easier and make coughs more productive.

**Warnings**

**Live warning:** This product contains acetaminophen. Severe liver damage may occur if you take:
 

- more than 12 softgels in 24 hours, which is the maximum daily amount;
- with other drugs containing acetaminophen;
- if or when alcohol is taken daily while using this product.

**Other live warning:** Acetaminophen may cause severe skin reactions. Symptoms may include:
 

- skin redness;
- skin rash;
- skin sores;
- skin blisters;
- skin peeling.

 If you have had a skin reaction to acetaminophen, stop use and seek medical help right away.

**See doctor warning:** If sore throat is severe, persists for more than 3 days, is accompanied or followed by fever, headache, rash, loss of voice, or swelling, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription narcotic medicine called MAOI (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have:**

- liver disease;
- kidney disease;
- high blood pressure;
- throat disease;
- any condition that is an enlarged prostate gland;
- history of chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema.

**Drug Facts (continued)**

**Directions**

- Take 2 softgels every 4 hours.
- Do not take more than directed (see Warnings section).
- Do not take more than 12 softgels in any 24-hour period.
- Adults and children 12 years of age and over: take 2 softgels every 4 hours.
- Children under 12 years of age: do not take.

**Other information**

- Store between 20° to 25°C (68° to 77°F).
- Keep out of reach of children.
- Excipients include: gelatin, croscarmellose sodium, polyethylene glycol, polyethylene glycol, croscarmellose sodium, croscarmellose sodium, croscarmellose sodium.

**Contains inactive ingredients:** gelatin, croscarmellose sodium, polyethylene glycol, polyethylene glycol, croscarmellose sodium, croscarmellose sodium.

**Questions or comments?** 1-888-333-9792

# MAXIMUM STRENGTH NON DROWSY DAYTIME SEVERE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride

capsule, liquid filled

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:75712-007
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	341;A13
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:75712-007-20	2 in 1 CARTON	03/28/2023	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

OTC Monograph Drug	M012	03/28/2023	
--------------------	------	------------	--

**Labeler** - DOLGENCORP, INC. (006946172)

Revised: 12/2023

DOLGENCORP, INC.