

**BUPROPION HYDROCHLORIDE - bupropion hydrochloride tablet, film coated**  
**Alembic Pharmaceuticals Limited**

-----

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use BUPROPION HYDROCHLORIDE TABLETS safely and effectively. See full prescribing information for BUPROPION HYDROCHLORIDE TABLETS, BUPROPION HYDROCHLORIDE tablets, for oral use

**Initial U.S. Approval: 1985**

<p><b>WARNING: SUICIDAL THOUGHTS AND BEHAVIORS</b> <i>See full prescribing information for complete boxed warning.</i></p> <ul style="list-style-type: none"><li>• <b>Increased risk of suicidal thinking and behavior in children, adolescents, and young adults taking antidepressants. (5.1)</b></li><li>• <b>Monitor for worsening and emergence of suicidal thoughts and behaviors. (5.1)</b></li></ul>
--

----- **RECENT MAJOR CHANGES** -----

Boxed Warning 05/2017  
Warnings and Precautions, Neuropsychiatric Adverse Events and Suicide Risk in Smoking Cessation Treatment (5.2)  
05/2017

----- **INDICATIONS AND USAGE** -----

Bupropion hydrochloride tablets USP are an amine ketone antidepressant, indicated for the treatment of major depressive disorder (MDD) (1)(1)

----- **DOSAGE AND ADMINISTRATION** -----

- Starting dose: 200 mg per day given as 100 mg twice daily. (2.1)
- General: Increase dose gradually to reduce seizure risk. (2.1, 5.3)
- After 3 days, may increase the dose to 300 mg per day, given as 100 mg 3 times daily at an interval of at least 6 hours between doses. (2.1)
- Usual target dose: 300 mg per day as 100 mg 3 times daily. (2.1)
- Maximum dose: 450 mg per day given as 150 mg 3 times daily. (2.1)
- Periodically reassess the dose and need for maintenance treatment. (2.1)
- Moderate to severe hepatic impairment: 75 mg once daily. (2.2, 8.7)
- Mild hepatic impairment: Consider reducing the dose and/or frequency of dosing. (2.2, 8.7)
- Renal impairment: Consider reducing the dose and/or frequency. (2.3, 8.6) (2)

----- **DOSAGE FORMS AND STRENGTHS** -----

• Tablets: 75 mg and 100 mg (3) (3)

----- **CONTRAINDICATIONS** -----

- Seizure disorder. (4, 5.3)
- Current or prior diagnosis of bulimia or anorexia nervosa. (4, 5.3)
- Abrupt discontinuation of alcohol, benzodiazepines, barbiturates, antiepileptic drugs. (4, 5.3)
- Monoamine Oxidase Inhibitors (MAOIs): Do not use MAOIs intended to treat psychiatric disorders with bupropion

hydrochloride tablets or within 14 days of stopping treatment with bupropion hydrochloride tablets. Do not use bupropion hydrochloride tablets within 14 days of stopping an MAOI intended to treat psychiatric disorders. In addition, do not start bupropion hydrochloride tablets in a patient who is being treated with linezolid or intravenous methylene blue. (4, 7.6)

- Known hypersensitivity to bupropion or other ingredients of bupropion hydrochloride tablets. (4, 5.8) (4)

#### ----- **WARNINGS AND PRECAUTIONS** -----

- Neuropsychiatric adverse events during smoking cessation: Postmarketing reports of serious or clinically significant neuropsychiatric adverse events have included changes in mood (including depression and mania), psychosis, hallucinations, paranoia, delusions, homicidal ideation, aggression, hostility, agitation, anxiety, and panic, as well as suicidal ideation, suicide attempt, and completed suicide. Observe patients attempting to quit smoking with bupropion for the occurrence of such symptoms and instruct them to discontinue bupropion and contact a healthcare provider if they experience such adverse events. (5.2)
- Seizure risk: The risk is dose-related. Can minimize risk by gradually increasing the dose and limiting daily dose to 450 mg. Discontinue if seizure occurs. (4, 5.3, 7.3)
- Hypertension: Bupropion hydrochloride tablets can increase blood pressure. Monitor blood pressure before initiating treatment and periodically during treatment. (5.4)
- Activation of mania/hypomania: Screen patients for bipolar disorder and monitor for these symptoms. (5.5)
- Psychosis and other neuropsychiatric reactions: Instruct patients to contact a healthcare professional if such reactions occur. (5.6)
- Angle-closure glaucoma: Angle-closure glaucoma has occurred in patients with untreated anatomically narrow angles treated with antidepressants. (5.7) (5)

#### ----- **ADVERSE REACTIONS** -----

Most common adverse reactions (incidence  $\geq 5\%$  and  $\geq 1\%$  more than placebo rate) are: agitation, dry mouth, constipation, headache/migraine, nausea/vomiting, dizziness, excessive sweating, tremor, insomnia, blurred vision, tachycardia, confusion, rash, hostility, cardiac arrhythmias, and auditory disturbance. (6.1) (6)

**To report SUSPECTED ADVERSE REACTIONS, contact FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).** (6)

#### ----- **DRUG INTERACTIONS** -----

- CYP2B6 inducers: Dose increase may be necessary if coadministered with CYP2B6 inducers (e.g., ritonavir, lopinavir, efavirenz, carbamazepine, phenobarbital, and phenytoin) based on clinical response, but should not exceed the maximum recommended dose. (7.1)
- Drugs metabolized by CYP2D6: Bupropion inhibits CYP2D6 and can increase concentrations of: antidepressants (e.g., venlafaxine, nortriptyline, imipramine, desipramine, paroxetine, fluoxetine, sertraline), antipsychotics (e.g., haloperidol, risperidone, thioridazine), beta-blockers (e.g., metoprolol), and Type 1C antiarrhythmics (e.g., propafenone, flecainide). Consider dose reduction when using with bupropion. (7.2)
- Digoxin: May decrease plasma digoxin levels. Monitor digoxin levels.
- Drugs that lower seizure threshold: Dose bupropion hydrochloride tablets with caution. (5.3, 7.3)
- Dopaminergic drugs (levodopa and amantadine): CNS toxicity can occur when used concomitantly with bupropion hydrochloride tablets. (7.4)
- MAOIs: Increased risk of hypertensive reactions can occur when used concomitantly with bupropion hydrochloride tablets. (7.6)
- Drug-laboratory test interactions: bupropion hydrochloride tablets can cause false-positive urine test results for amphetamines. (7.7) (7)

#### ----- **USE IN SPECIFIC POPULATIONS** -----

- Pregnancy: Use only if benefit outweighs potential risk to the fetus. (8.1) (8)

**See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.**

**Revised: 9/2019**

---

**FULL PRESCRIBING INFORMATION: CONTENTS\***  
**WARNING: SUICIDAL THOUGHTS AND BEHAVIORS**  
**1 INDICATIONS AND USAGE**  
**2 DOSAGE AND ADMINISTRATION**  
2.1 General Instructions for Use  
2.2 Dose Adjustment in Patients with Hepatic Impairment  
2.3 Dose Adjustment in Patients with Renal Impairment

- 2.4 Switching a Patient to or from a Monoamine Oxidase Inhibitor (MAOI) Antidepressant
- 2.5 Use of Bupropion Hydrochloride Tablets with Reversible MAOIs Such as Linezolid or Methylene Blue

### **3 DOSAGE FORMS AND STRENGTHS**

### **4 CONTRAINDICATIONS**

### **5 WARNINGS AND PRECAUTIONS**

- 5.1 Suicidal Thoughts and Behaviors in Children, Adolescents, and Young Adults
- 5.2 Neuropsychiatric Adverse Events and Suicide Risk in Smoking Cessation Treatment
- 5.3 Seizure
- 5.4 Hypertension
- 5.5 Activation of Mania/Hypomania
- 5.6 Psychosis and Other Neuropsychiatric Reactions
- 5.7 Angle-Closure Glaucoma
- 5.8 Hypersensitivity Reactions

### **6 ADVERSE REACTIONS**

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

### **7 DRUG INTERACTIONS**

- 7.1 Potential for Other Drugs to Affect Bupropion Hydrochloride Tablets
- 7.2 Potential for Bupropion Hydrochloride Tablets to Affect Other Drugs
- 7.3 Drugs that Lower Seizure Threshold
- 7.4 Dopaminergic Drugs (Levodopa and Amantadine)
- 7.5 Use with Alcohol
- 7.6 MAO Inhibitors
- 7.7 Drug-Laboratory Test Interactions

### **8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment

### **9 DRUG ABUSE AND DEPENDENCE**

- 9.1 Controlled Substance
- 9.2 Abuse

### **10 OVERDOSAGE**

- 10.1 Human Overdose Experience
- 10.2 Overdosage Management

### **11 DESCRIPTION**

### **12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics

### **13 NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

### **14 CLINICAL STUDIES**

### **16 HOW SUPPLIED/STORAGE AND HANDLING**

### **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

## **WARNING: SUICIDAL THOUGHTS AND BEHAVIORS**

### **SUICIDALITY AND ANTIDEPRESSANT DRUGS**

**Antidepressants increased the risk of suicidal thoughts and behavior in children, adolescents, and young adults in short-term trials. These trials did not show an increase in the risk of suicidal thoughts and behavior with antidepressant use in subjects over age 24; there was a reduction in risk with antidepressant use in subjects aged 65 and older [see *Warnings and Precautions (5.1)*].**

**In patients of all ages who are started on antidepressant therapy, monitor closely for worsening, and for emergence of suicidal thoughts and behaviors. Advise families and caregivers of the need for close observation and communication with the prescriber [see *Warnings and Precautions (5.1)*].**

## **1 INDICATIONS AND USAGE**

Bupropion hydrochloride tablets USP are indicated for the treatment of major depressive disorder (MDD), as defined by the Diagnostic and Statistical Manual (DSM).

The efficacy of bupropion hydrochloride tablets USP in the treatment of a major depressive episode was established in two 4-week controlled inpatient trials and one 6-week controlled outpatient trial of adult subjects with MDD[see *Clinical Studies (14)*].

## **2 DOSAGE AND ADMINISTRATION**

### **2.1 General Instructions for Use**

To minimize the risk of seizure, increase the dose gradually [see *Warnings and Precautions (5.3)*]. Increases in dose should not exceed 100 mg per day in a 3-day period. Bupropion hydrochloride tablets should be swallowed whole and not crushed, divided, or chewed. Bupropion hydrochloride tablets may be taken with or without food.

The recommended starting dose is 200 mg per day, given as 100 mg twice daily. After 3 days of dosing, the dose may be increased to 300 mg per day, given as 100 mg 3 times daily, with at least 6 hours between successive doses. Dosing above 300 mg per day may be accomplished using the 75 mg or 100 mg tablets.

A maximum of 450 mg per day, given in divided doses of not more than 150 mg each, may be considered for patients who show no clinical improvement after several weeks of treatment at 300 mg per day. Administer the 100 mg tablet 4 times daily to not exceed the limit of 150 mg in a single dose.

It is generally agreed that acute episodes of depression require several months or longer of antidepressant drug treatment beyond the response in the acute episode. It is unknown whether the dose of bupropion hydrochloride tablets needed for maintenance treatment is identical to the dose that provided an initial response. Periodically reassess the need for maintenance treatment and the

appropriate dose for such treatment.

## **2.2 Dose Adjustment in Patients with Hepatic Impairment**

In patients with moderate to severe hepatic impairment (Child-Pugh score: 7 to 15), the maximum dose of bupropion hydrochloride tablet is 75 mg per day. In patients with mild hepatic impairment (Child-Pugh score: 5 to 6), consider reducing the dose and/or frequency of dosing [*see Use in Specific Populations (8.7) and Clinical Pharmacology (12.3)*].

## **2.3 Dose Adjustment in Patients with Renal Impairment**

Consider reducing the dose and/or frequency of bupropion hydrochloride tablets in patients with renal impairment (Glomerular Filtration Rate less than 90 mL per min) [*see Use in Specific Populations (8.6) and Clinical Pharmacology (12.3)*].

## **2.4 Switching a Patient to or from a Monoamine Oxidase Inhibitor (MAOI) Antidepressant**

At least 14 days should elapse between discontinuation of an MAOI intended to treat depression and initiation of therapy with bupropion hydrochloride tablets. Conversely, at least 14 days should be allowed after stopping bupropion hydrochloride tablets before starting an MAOI antidepressant [*see Contraindications (4) and Drug Interactions (7.6)*].

## **2.5 Use of Bupropion Hydrochloride Tablets with Reversible MAOIs Such as Linezolid or Methylene Blue**

Do not start bupropion hydrochloride tablets in a patient who is being treated with a reversible MAOI such as linezolid or intravenous methylene blue. Drug interactions can increase the risk of hypertensive reactions. In a patient who requires more urgent treatment of a psychiatric condition, non-pharmacological interventions, including hospitalization, should be considered [*see Contraindications (4), Drug Interactions (7.6)*].

In some cases, a patient already receiving therapy with bupropion hydrochloride tablets may require urgent treatment with linezolid or intravenous methylene blue. If acceptable alternatives to linezolid or intravenous methylene blue treatment are not available and the potential benefits of linezolid or intravenous methylene blue treatment are judged to outweigh the risks of hypertensive reactions in a particular patient, bupropion hydrochloride tablets should be stopped promptly, and linezolid or intravenous methylene blue can be administered. The patient should be monitored for 2 weeks or until 24 hours after the last dose of linezolid or intravenous methylene blue, whichever comes first. Therapy with bupropion hydrochloride tablets may be resumed 24 hours after the last dose of linezolid or intravenous methylene blue.

The risk of administering methylene blue by non-intravenous routes (such as oral tablets or by local injection) or in intravenous doses much lower than 1 mg per kg with bupropion hydrochloride tablet is unclear. The clinician should, nevertheless, be aware of the possibility of a drug interaction with such use [*see Contraindications (4), Drug Interactions (7.6)*].

## **3 DOSAGE FORMS AND STRENGTHS**

- 75 mg – light orange colored, round, biconvex film coated tablets debossed with ‘L295’ on one side and plain on other side.
- 100 mg – light yellow colored, round, biconvex film coated tablets debossed with ‘L296’ on one side and plain on other side.

## 4 CONTRAINDICATIONS

- Bupropion hydrochloride tablets are contraindicated in patients with a seizure disorder.
- Bupropion hydrochloride tablets are contraindicated in patients with a current or prior diagnosis of bulimia or anorexia nervosa as a higher incidence of seizures was observed in such patients treated with bupropion hydrochloride tablets [see *Warnings and Precautions (5.3)*].
- Bupropion hydrochloride tablets are contraindicated in patients undergoing abrupt discontinuation of alcohol, benzodiazepines, barbiturates, and antiepileptic drugs [see *Warnings and Precautions (5.3), Drug Interactions (7.3)*].
- The use of MAOIs (intended to treat psychiatric disorders) concomitantly with bupropion hydrochloride tablets or within 14 days of discontinuing treatment with bupropion hydrochloride tablets are contraindicated. There is an increased risk of hypertensive reactions when bupropion hydrochloride tablets are used concomitantly with MAOIs. The use of bupropion hydrochloride tablets within 14 days of discontinuing treatment with an MAOI is also contraindicated. Starting bupropion hydrochloride tablets in a patient treated with reversible MAOIs such as linezolid or intravenous methylene blue is contraindicated [see *Dosage and Administration (2.4, 2.5), Warnings and Precautions (5.4), Drug Interactions (7.6)*].
- Bupropion hydrochloride tablets are contraindicated in patients with known hypersensitivity to bupropion or other ingredients of bupropion hydrochloride tablets. Anaphylactoid/anaphylactic reactions and Stevens-Johnson syndrome have been reported [see *Warnings and Precautions (5.8)*].

## 5 WARNINGS AND PRECAUTIONS

### 5.1 Suicidal Thoughts and Behaviors in Children, Adolescents, and Young Adults

Patients with MDD, both adult and pediatric, may experience worsening of their depression and/or the emergence of suicidal ideation and behavior (suicidality) or unusual changes in behavior, whether or not they are taking antidepressant medications, and this risk may persist until significant remission occurs. Suicide is a known risk of depression and certain other psychiatric disorders, and these disorders themselves are the strongest predictors of suicide. There has been a long-standing concern that antidepressants may have a role in inducing worsening of depression and the emergence of suicidality in certain patients during the early phases of treatment.

Pooled analyses of short-term placebo-controlled trials of antidepressant drugs (selective serotonin reuptake inhibitors [SSRIs] and others) show that these drugs increase the risk of suicidal thinking and behavior (suicidality) in children, adolescents, and young adults (ages 18 to 24) with MDD and other psychiatric disorders. Short-term clinical trials did not show an increase in the risk of suicidality with antidepressants compared with placebo in adults beyond age 24; there was a reduction with antidepressants compared with placebo in adults aged 65 and older.

The pooled analyses of placebo-controlled trials in children and adolescents with MDD, obsessive compulsive disorder (OCD), or other psychiatric disorders included a total of 24 short-term trials of 9 antidepressant drugs in over 4,400 subjects. The pooled analyses of placebo-controlled trials in adults with MDD or other psychiatric disorders included a total of 295 short-term trials (median duration of 2 months) of 11 antidepressant drugs in over 77,000 subjects. There was considerable variation in risk of suicidality among drugs, but a tendency toward an increase in the younger subjects for almost all drugs studied. There were differences in absolute risk of suicidality across the different indications, with the highest incidence in MDD. The risk differences (drug vs. placebo), however, were relatively stable within age strata and across indications. These risk differences (drug-placebo difference in the number

of cases of suicidality per 1,000 subjects treated) are provided in Table 1.

**Table 1: Risk Differences in the Number of Suicidal Cases by Age Group in the Pooled Placebo-Controlled Trials of Antidepressants in Pediatric and Adult Subjects**

Age Range	Drug - Placebo Difference in Number of Cases of Suicidality per 1,000 Subjects Treated
Increases Compared with Placebo	
<18 years	14 additional cases
18 to 24 years	5 additional cases
Decreases Compared with Placebo	
25 to 64 years	1 fewer case
≥65 years	6 fewer cases

No suicides occurred in any of the pediatric trials. There were suicides in the adult trials, but the number was not sufficient to reach any conclusion about drug effect on suicide.

It is unknown whether the suicidality risk extends to longer-term use, i.e., beyond several months. However, there is substantial evidence from placebo-controlled maintenance trials in adults with depression that the use of antidepressants can delay the recurrence of depression.

**All patients being treated with antidepressants for any indication should be monitored appropriately and observed closely for clinical worsening, suicidality, and unusual changes in behavior, especially during the initial few months of a course of drug therapy, or at times of dose changes, either increases or decreases [see Boxed Warning].**

The following symptoms, anxiety, agitation, panic attacks, insomnia, irritability, hostility, aggressiveness, impulsivity, akathisia (psychomotor restlessness), hypomania, and mania, have been reported in adult and pediatric patients being treated with antidepressants for major depressive disorder as well as for other indications, both psychiatric and nonpsychiatric. Although a causal link between the emergence of such symptoms and either the worsening of depression and/or the emergence of suicidal impulses has not been established, there is concern that such symptoms may represent precursors to emerging suicidality.

Consideration should be given to changing the therapeutic regimen, including possibly discontinuing the medication, in patients whose depression is persistently worse, or who are experiencing emergent suicidality or symptoms that might be precursors to worsening depression or suicidality, especially if these symptoms are severe, abrupt in onset, or were not part of the patient's presenting symptoms.

**Families and caregivers of patients being treated with antidepressants for MDD or other indications, both psychiatric and nonpsychiatric, should be alerted about the need to monitor patients for the emergence of agitation, irritability, unusual changes in behavior, and the other symptoms described above, as well as the emergence of suicidality, and to report such symptoms immediately to healthcare providers. Such monitoring should include daily observation by families and caregivers. Prescriptions for bupropion hydrochloride tablets should be written for the smallest quantity of tablets consistent with good patient management, in order to reduce the risk of overdose.**

## 5.2 Neuropsychiatric Adverse Events and Suicide Risk in Smoking Cessation Treatment

Bupropion hydrochloride tablet is not approved for smoking cessation treatment; however, it contains the same active ingredient as the smoking cessation medication ZYBAN. Serious neuropsychiatric

adverse events have been reported in patients taking bupropion for smoking cessation. These postmarketing reports have included changes in mood (including depression and mania), psychosis, hallucinations, paranoia, delusions, homicidal ideation, aggression, hostility, agitation, anxiety, and panic, as well as suicidal ideation, suicide attempt, and completed suicide [see *Adverse Reactions (6.2)*]. Some patients who stopped smoking may have been experiencing symptoms of nicotine withdrawal, including depressed mood. Depression, rarely including suicidal ideation, has been reported in smokers undergoing a smoking cessation attempt without medication. However, some of these adverse events occurred in patients taking bupropion who continued to smoke.

Neuropsychiatric adverse events occurred in patients without and with pre-existing psychiatric disease; some patients experienced worsening of their psychiatric illnesses. Observe patients for the occurrence of neuropsychiatric adverse events. Advise patients and caregivers that the patient should stop taking bupropion hydrochloride tablet and contact a healthcare provider immediately if agitation, depressed mood, or changes in behavior or thinking that are not typical for the patient are observed, or if the patient develops suicidal ideation or suicidal behavior. In many postmarketing cases, resolution of symptoms after discontinuation of bupropion was reported. However, the symptoms persisted in some cases; therefore, ongoing monitoring and supportive care should be provided until symptoms resolve.

### 5.3 Seizure

Bupropion hydrochloride tablets can cause seizure. The risk of seizure is dose-related. The dose should not exceed 450 mg per day. Increase the dose gradually. Discontinue bupropion hydrochloride tablets and do not restart treatment if the patient experiences a seizure.

The risk of seizures is also related to patient factors, clinical situations, and concomitant medications that lower the seizure threshold. Consider these risks before initiating treatment with bupropion hydrochloride tablets. Bupropion hydrochloride tablets are contraindicated in patients with a seizure disorder, current or prior diagnosis of anorexia nervosa or bulimia, or undergoing abrupt discontinuation of alcohol, benzodiazepines, barbiturates, and antiepileptic drugs [see *Contraindications (4), Drug Interactions (7.3)*]. The following conditions can also increase the risk of seizure: severe head injury; arteriovenous malformation; CNS tumor or CNS infection; severe stroke; concomitant use of other medications that lower the seizure threshold (e.g., other bupropion products, antipsychotics, tricyclic antidepressants, theophylline, and systemic corticosteroids); metabolic disorders (e.g., hypoglycemia, hyponatremia, severe hepatic impairment, and hypoxia); use of illicit drugs (e.g., cocaine); or abuse or misuse of prescription drugs such as CNS stimulants. Additional predisposing conditions include diabetes mellitus treated with oral hypoglycemic drugs or insulin; use of anorectic drugs; and excessive use of alcohol, benzodiazepines, sedative/hypnotics, or opiates.

Incidence of Seizure with Bupropion Use: Bupropion is associated with seizures in approximately 0.4% (4/1,000) of patients treated at doses up to 450 mg per day. The estimated seizure incidence for bupropion hydrochloride tablets increases almost 10-fold between 450 mg and 600 mg per day.

The risk of seizure can be reduced if the dose of bupropion hydrochloride tablets does not exceed 450 mg per day, given as 150 mg 3 times daily, and the titration rate is gradual.

### 5.4 Hypertension

Treatment with bupropion hydrochloride tablets can result in elevated blood pressure and hypertension. Assess blood pressure before initiating treatment with bupropion hydrochloride tablets, and monitor periodically during treatment. The risk of hypertension is increased if bupropion hydrochloride tablets are used concomitantly with MAOIs or other drugs that increase dopaminergic or noradrenergic activity [see *Contraindications (4)*].

Data from a comparative trial of the sustained-release formulation of bupropion HCl, nicotine transdermal system (NTS), the combination of sustained-release bupropion plus NTS, and placebo as an aid to smoking cessation suggest a higher incidence of treatment-emergent hypertension in patients



treated with the combination of sustained-release bupropion and NTS. In this trial, 6.1% of subjects treated with the combination of sustained-release bupropion and NTS had treatment-emergent hypertension compared with 2.5%, 1.6%, and 3.1% of subjects treated with sustained-release bupropion, NTS, and placebo, respectively. The majority of these subjects had evidence of pre-existing hypertension. Three subjects (1.2%) treated with the combination of sustained-release bupropion and NTS and 1 subject (0.4%) treated with NTS had study medication discontinued due to hypertension compared with none of the subjects treated with sustained-release bupropion or placebo. Monitoring of blood pressure is recommended in patients who receive the combination of bupropion and nicotine replacement.

In a clinical trial of bupropion immediate-release in MDD subjects with stable congestive heart failure (N=36), bupropion was associated with an exacerbation of pre-existing hypertension in 2 subjects, leading to discontinuation of bupropion treatment. There are no controlled trials assessing the safety of bupropion in patients with a recent history of myocardial infarction or unstable cardiac disease.

### **5.5 Activation of Mania/Hypomania**

Antidepressant treatment can precipitate a manic, mixed, or hypomanic manic episode. The risk appears to be increased in patients with bipolar disorder or who have risk factors for bipolar disorder. Prior to initiating bupropion hydrochloride tablets, screen patients for a history of bipolar disorder and the presence of risk factors for bipolar disorder (e.g., family history of bipolar disorder, suicide, or depression). Bupropion hydrochloride tablets are not approved for use in treating bipolar depression.

### **5.6 Psychosis and Other Neuropsychiatric Reactions**

Depressed patients treated with bupropion hydrochloride tablets have had a variety of neuropsychiatric signs and symptoms, including delusions, hallucinations, psychosis, concentration disturbance, paranoia, and confusion. Some of these patients had a diagnosis of bipolar disorder. In some cases, these symptoms abated upon dose reduction and/or withdrawal of treatment. Instruct patients to contact a healthcare professional if such reactions occur.

### **5.7 Angle-Closure Glaucoma**

The pupillary dilation that occurs following use of many antidepressant drugs including bupropion hydrochloride tablets may trigger an angle-closure attack in a patient with anatomically narrow angles who does not have a patent iridectomy.

### **5.8 Hypersensitivity Reactions**

Anaphylactoid/anaphylactic reactions have occurred during clinical trials with bupropion. Reactions have been characterized by pruritus, urticaria, angioedema, and dyspnea requiring medical treatment. In addition, there have been rare, spontaneous postmarketing reports of erythema multiforme, Stevens-Johnson syndrome, and anaphylactic shock associated with bupropion. Instruct patients to discontinue bupropion hydrochloride tablets and consult a healthcare provider if they develop an allergic or anaphylactoid/anaphylactic reaction (e.g., skin rash, pruritus, hives, chest pain, edema, and shortness of breath) during treatment.

There are reports of arthralgia, myalgia, fever with rash and other serum sickness-like symptoms suggestive of delayed hypersensitivity.

## 6 ADVERSE REACTIONS

The following adverse reactions are discussed in greater detail in other sections of the labeling:

- Suicidal thoughts and behaviors in adolescents and young adults [see *Boxed Warning, Warnings and Precautions (5.1)*]
- Neuropsychiatric symptoms and suicide risk in smoking cessation treatment [see *Warnings and Precautions (5.2)*]
- Seizure [see *Warnings and Precautions (5.3)*]
- Hypertension [see *Warnings and Precautions (5.4)*]
- Activation of mania or hypomania [see *Warnings and Precautions (5.5)*]
- Psychosis and other neuropsychiatric reactions [see *Warnings and Precautions (5.6)*]
- Angle-closure glaucoma [see *Warnings and Precautions (5.7)*]
- Hypersensitivity reactions [see *Warnings and Precautions (5.8)*]

### 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared with rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

#### Adverse Reactions Leading to Discontinuation of Treatment

Adverse reactions were sufficiently troublesome to cause discontinuation of treatment with bupropion hydrochloride tablets in approximately 10% of the 2,400 subjects and healthy volunteers who participated in clinical trials during the product's initial development. The more common events causing discontinuation include neuropsychiatric disturbances (3%), primarily agitation and abnormalities in mental status; gastrointestinal disturbances (2.1%), primarily nausea and vomiting; neurological disturbances (1.7%), primarily seizures, headaches, and sleep disturbances; and dermatologic problems (1.4%), primarily rashes. It is important to note, however, that many of these events occurred at doses that exceed the recommended daily dose.

Commonly Observed Adverse Reactions: Adverse reactions commonly encountered in subjects treated with bupropion hydrochloride tablets are agitation, dry mouth, insomnia, headache/migraine, nausea/vomiting, constipation, tremor, dizziness, excessive sweating, blurred vision, tachycardia, confusion, rash, hostility, cardiac arrhythmia, and auditory disturbance.

Table 2 summarizes the adverse reactions that occurred in placebo-controlled trials at an incidence of at least 1% of subjects receiving bupropion hydrochloride tablets and more frequently in these subjects than in the placebo group.

**Table 2. Adverse Reactions Reported by at Least 1% of Subjects and at a Greater Frequency than Placebo in Controlled Clinical Trials**

<b>A dverse Reac tion</b>	<b>Bupropion Hydrochloride Tablets ( n =323) %</b>	<b>P lace bo ( n =185) %</b>
Cardiovascular		
Cardiac arrhythmias	5.3	4.3
Dizziness	22.3	16.2
Hypertension	4.3	1.6

Hypotension	2.5	2.2
Palpitations	3.7	2.2
Syncope	1.2	0.5
Tachycardia	10.8	8.6
Dermatologic		
Pruritus	2.2	0
Rash	8	6.5
Gastrointestinal		
Appetite increase	3.7	2.2
Constipation	26	17.3
Dyspepsia	3.1	2.2
Nausea/vomiting	22.9	18.9
Genitourinary		
Impotence	3.4	3.1
Menstrual complaints	4.7	1.1
Urinary frequency	2.5	2.2
Musculoskeletal		
Arthritis	3.1	2.7
Neurological		
Akathisia	1.5	1.1
Cutaneous temperature disturbance	1.9	1.6
Dry mouth	27.6	18.4
Excessive sweating	22.3	14.6
Headache/migraine	25.7	22.2
Impaired sleep quality	4	1.6
Insomnia	18.6	15.7
Sedation	19.8	19.5
Sensory disturbance	4	3.2
Tremor	21.1	7.6
Neuropsychiatric		
Agitation	31.9	22.2
Anxiety	3.1	1.1
Confusion	8.4	4.9
Decreased libido	3.1	1.6
Delusions	1.2	1.1
Euphoria	1.2	0.5
Hostility	5.6	3.8
Nonspecific		
Fever/chills	1.2	0.5
Special Senses		
Auditory disturbance	5.3	3.2
Blurred vision	14.6	10.3
Gustatory disturbance	3.1	1.1

#### Other Adverse Reactions Observed during the Clinical Development of Bupropion Hydrochloride Tablets

The conditions and duration of exposure to bupropion hydrochloride tablets varied greatly, and a substantial proportion of the experience was gained in open and uncontrolled clinical settings. During this experience, numerous adverse events were reported; however, without appropriate controls, it is

impossible to determine with certainty which events were or were not caused by bupropion hydrochloride tablets. The following enumeration is organized by organ system and describes events in terms of their relative frequency of reporting in the database.

The following definitions of frequency are used: Frequent adverse reactions are defined as those occurring in at least 1/100 subjects. Infrequent adverse reactions are those occurring in 1/100 to 1/1,000 subjects, while rare events are those occurring in less than 1/1,000 subjects. *Cardiovascular*: Frequent was edema; infrequent were chest pain, electrocardiogram (ECG) abnormalities (premature beats and nonspecific ST-T changes), and shortness of breath/dyspnea; rare were flushing, and myocardial infarction.

*Dermatologic*: Infrequent was alopecia.

*Endocrine*: Infrequent was gynecomastia; rare was glycosuria.

*Gastrointestinal*: Infrequent were dysphagia, thirst disturbance, and liver damage/jaundice; rare was intestinal perforation.

*Genitourinary*: Frequent was nocturia; infrequent were vaginal irritation, testicular swelling, urinary tract infection, painful erection, and retarded ejaculation; rare were enuresis, and urinary incontinence.

*Neurological*: Frequent were ataxia/incoordination, seizure, myoclonus, dyskinesia, and dystonia; infrequent were mydriasis, vertigo, and dysarthria; rare were electroencephalogram (EEG) abnormality, and impaired attention.

*Neuropsychiatric*: Frequent were mania/hypomania, increased libido, hallucinations, decrease in sexual function, and depression; infrequent were memory impairment, depersonalization, psychosis, dysphoria, mood instability, paranoia, and formal thought disorder ; rare was suicidal ideation.

*Oral Complaints*: Frequent was stomatitis; infrequent were toothache, bruxism, gum irritation, and oral edema.

*Respiratory*: Infrequent were bronchitis and shortness of breath/dyspnea; rare was pulmonary embolism.

*Special Senses*: Infrequent was visual disturbance; rare was diplopia.

*Nonspecific*: Frequent were flu-like symptoms; infrequent was nonspecific pain; rare was overdose.

#### Altered Appetite and Weight

A weight loss of greater than 5 lbs occurred in 28% of subjects receiving bupropion hydrochloride tablets. This incidence is approximately double that seen in comparable subjects treated with tricyclics or placebo. Furthermore, while 35% of subjects receiving tricyclic antidepressants gained weight, only 9.4% of subjects treated with bupropion hydrochloride tablets did. Consequently, if weight loss is a major presenting sign of a patient's depressive illness, the anorectic and/or weight-reducing potential of bupropion hydrochloride tablets should be considered.

## **6.2 Postmarketing Experience**

The following adverse reactions have been identified during post-approval use of bupropion hydrochloride tablets and are not described elsewhere in the label. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

#### Body (General)

Arthralgia, myalgia, and fever with rash and other symptoms suggestive of delayed hypersensitivity. These symptoms may resemble serum sickness [see *Warnings and Precautions (5.8)*].

### Cardiovascular

Hypertension (in some cases severe), orthostatic hypotension, third degree heart block.

### Endocrine

Hyponatremia, syndrome of inappropriate antidiuretic hormone secretion, hyperglycemia, hypoglycemia.

### Gastrointestinal

Esophagitis, hepatitis.

### Hemic and Lymphatic

Ecchymosis, leukocytosis, leukopenia, thrombocytopenia. Altered PT and/or INR, infrequently associated with hemorrhagic or thrombotic complications, were observed when bupropion was coadministered with warfarin.

### Musculoskeletal

Muscle rigidity/fever/rhabdomyolysis, muscle weakness.

### Nervous System

Aggression, coma, completed suicide, delirium, dream abnormalities, paranoid ideation, paresthesia, parkinsonism, restlessness, suicide attempt, unmasking of tardive dyskinesia.

### Skin and Appendages

Stevens-Johnson syndrome, angioedema, exfoliative dermatitis, urticaria.

### Special Senses

Tinnitus, increased intraocular pressure.

## **7 DRUG INTERACTIONS**

### **7.1 Potential for Other Drugs to Affect Bupropion Hydrochloride Tablets**

Bupropion is primarily metabolized to hydroxybupropion by CYP2B6. Therefore, the potential exists for drug interactions between bupropion hydrochloride tablets and drugs that are inhibitors or inducers of CYP2B6.

#### Inhibitors of CYP2B6

*Ticlopidine and Clopidogrel:* Concomitant treatment with these drugs can increase bupropion exposure but decrease hydroxybupropion exposure. Based on clinical response, dosage adjustment of bupropion hydrochloride tablets may be necessary when coadministered with CYP2B6 inhibitors (e.g., ticlopidine or clopidogrel) [see *Clinical Pharmacology (12.3)*].

#### Inducers of CYP2B6

*Ritonavir, Lopinavir, and Efavirenz:* Concomitant treatment with these drugs can decrease bupropion and hydroxybupropion exposure. Dosage increase of bupropion hydrochloride tablets may be necessary when coadministered with ritonavir, lopinavir, or efavirenz [see *Clinical Pharmacology (12.3)*] but should not exceed the maximum recommended dose.

*Carbamazepine, Phenobarbital, Phenytoin*: While not systematically studied, these drugs may induce the metabolism of bupropion and may decrease bupropion exposure [see *Clinical Pharmacology (12.3)*]. If bupropion is used concomitantly with a CYP inducer, it may be necessary to increase the dose of bupropion, but the maximum recommended dose should not be exceeded.

## **7.2 Potential for Bupropion Hydrochloride Tablets to Affect Other Drugs**

### Drugs Metabolized by CYP2D6

Bupropion and its metabolites (erythrohydrobupropion, threo hydrobupropion, hydroxybupropion) are CYP2D6 inhibitors. Therefore, coadministration of bupropion hydrochloride tablets with drugs that are metabolized by CYP2D6 can increase the exposures of drugs that are substrates of CYP2D6. Such drugs include certain antidepressants (e.g., venlafaxine, nortriptyline, imipramine, desipramine, paroxetine, fluoxetine, and sertraline), antipsychotics (e.g. haloperidol, risperidone, thioridazine), beta-blockers (e.g., metoprolol), and Type 1C antiarrhythmics (e.g., propafenone and flecainide). When used concomitantly with bupropion hydrochloride tablets, it may be necessary to decrease the dose of these CYP2D6 substrates, particularly for drugs with a narrow therapeutic index.

Drugs that require metabolic activation by CYP2D6 to be effective (e.g., tamoxifen) theoretically could have reduced efficacy when administered concomitantly with inhibitors of CYP2D6 such as bupropion. Patients treated concomitantly with bupropion hydrochloride tablets and such drugs may require increased doses of the drug [see *Clinical Pharmacology (12.3)*].

### Digoxin

Coadministration of bupropion hydrochloride tablets with digoxin may decrease plasma digoxin levels. Monitor plasma digoxin levels in patients treated concomitantly with bupropion hydrochloride tablets and digoxin [see *Clinical Pharmacology (12.3)*].

## **7.3 Drugs that Lower Seizure Threshold**

Use extreme caution when coadministering bupropion hydrochloride tablets with other drugs that lower the seizure threshold (e.g., other bupropion products, antipsychotics, antidepressants, theophylline, or systemic corticosteroids). Use low initial doses and increase the dose gradually [see *Contraindications (4), Warnings and Precautions (5.3)*].

## **7.4 Dopaminergic Drugs (Levodopa and Amantadine)**

Bupropion, levodopa, and amantadine have dopamine agonist effects. CNS toxicity has been reported when bupropion was coadministered with levodopa or amantadine. Adverse reactions have included restlessness, agitation, tremor, ataxia, gait disturbance, vertigo, and dizziness. It is presumed that the toxicity results from cumulative dopamine agonist effects. Use caution when administering bupropion hydrochloride tablets concomitantly with these drugs.

## **7.5 Use with Alcohol**

In postmarketing experience, there have been rare reports of adverse neuropsychiatric events or reduced alcohol tolerance in patients who were drinking alcohol during treatment with bupropion hydrochloride tablets. The consumption of alcohol during treatment with bupropion hydrochloride

tablets should be minimized or avoided.

## 7.6 MAO Inhibitors

Bupropion inhibits the reuptake of dopamine and norepinephrine. Concomitant use of MAOIs and bupropion is contraindicated because there is an increased risk of hypertensive reactions if bupropion is used concomitantly with MAOIs. Studies in animals demonstrate that the acute toxicity of bupropion is enhanced by the MAO inhibitor phenelzine. At least 14 days should elapse between discontinuation of an MAOI intended to treat depression and initiation of treatment with bupropion hydrochloride tablets. Conversely, at least 14 days should be allowed after stopping bupropion hydrochloride tablets before starting an MAOI antidepressant [see *Dosage and Administration (2.4, 2.5) and Contraindications (4)*].

## 7.7 Drug-Laboratory Test Interactions

False-positive urine immunoassay screening tests for amphetamines have been reported in patients taking bupropion. This is due to lack of specificity of some screening tests. False-positive test results may result even following discontinuation of bupropion therapy. Confirmatory tests, such as gas chromatography/mass spectrometry, will distinguish bupropion from amphetamines.

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

Pregnancy Category C

#### Risk Summary

Data from epidemiological studies of pregnant women exposed to bupropion in the first trimester indicate no increased risk of congenital malformations overall. All pregnancies, regardless of drug exposure, have a background rate of 2% to 4% for major malformations, and 15% to 20% for pregnancy loss. No clear evidence of teratogenic activity was found in reproductive developmental studies conducted in rats and rabbits; however, in rabbits, slightly increased incidences of fetal malformations and skeletal variations were observed at doses approximately equal to the maximum recommended human dose (MRHD) and greater and decreased fetal weights were seen at doses twice the MRHD and greater. Bupropion hydrochloride tablets should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

#### Clinical Considerations

Consider the risks of untreated depression when discontinuing or changing treatment with antidepressant medications during pregnancy and postpartum.

#### Human Data

Data from the international bupropion Pregnancy Registry (675 first-trimester exposures) and a retrospective cohort study using the United Healthcare database (1,213 first trimester exposures) did not show an increased risk for malformations overall.

No increased risk for cardiovascular malformations overall has been observed after bupropion exposure during the first trimester. The prospectively observed rate of cardiovascular malformations in pregnancies with exposure to bupropion in the first trimester from the international Pregnancy Registry was 1.3% (9 cardiovascular malformations/675 first-trimester maternal bupropion exposures), which is similar to the background rate of cardiovascular malformations (approximately 1%). Data from the United Healthcare database and a case-controlled study (6,853 infants with cardiovascular

malformations and 5,763 with non-cardiovascular malformations) from the National Birth Defects Prevention Study (NBDPS) did not show an increased risk for cardiovascular malformations overall after bupropion exposure during the first trimester.

Study findings on bupropion exposure during the first trimester and risk for left ventricular outflow tract obstruction (LVOTO) are inconsistent and do not allow conclusions regarding a possible association. The United Healthcare database lacked sufficient power to evaluate this association; the NBDPS found increased risk for LVOTO (n = 10; adjusted OR = 2.6; 95% CI: 1.2, 5.7), and the Slone Epidemiology case control study did not find increased risk for LVOTO.

Study findings on bupropion exposure during the first trimester and risk for ventricular septal defect (VSD) are inconsistent and do not allow conclusions regarding a possible association. The Slone Epidemiology Study found an increased risk for VSD following first trimester maternal bupropion exposure (n = 17; adjusted OR = 2.5; 95% CI: 1.3, 5) but did not find increased risk for any other cardiovascular malformations studied (including LVOTO as above). The NBDPS and United Healthcare database study did not find an association between first trimester maternal bupropion exposure and VSD.

For the findings of LVOTO and VSD, the studies were limited by the small number of exposed cases, inconsistent findings among studies, and the potential for chance findings from multiple comparisons in case control studies.

#### Animal Data

In studies conducted in rats and rabbits, bupropion was administered orally during the period of organogenesis at doses of up to 450 mg and 150 mg per kg per day, respectively (approximately 11 and 7 times the MRHD, respectively, on a mg per m<sup>2</sup> basis). No clear evidence of teratogenic activity was found in either species; however, in rabbits, slightly increased incidences of fetal malformations and skeletal variations were observed at the lowest dose tested (25 mg per kg per day, approximately equal to the MRHD on a mg per m<sup>2</sup> basis) and greater. Decreased fetal weights were observed at 50 mg per kg and greater.

When rats were administered bupropion at oral doses of up to 300 mg per kg per day (approximately 7 times the MRHD on a mg per m<sup>2</sup> basis) prior to mating and throughout pregnancy and lactation, there were no apparent adverse effects on offspring development.

### **8.3 Nursing Mothers**

Bupropion and its metabolites are present in human milk. In a lactation study of 10 women, levels of orally dosed bupropion and its active metabolites were measured in expressed milk. The average daily infant exposure (assuming 150 mL per kg daily consumption) to bupropion and its active metabolites was 2% of the maternal weight-adjusted dose. Exercise caution when bupropion hydrochloride tablets are administered to a nursing woman.

### **8.4 Pediatric Use**

Safety and effectiveness in the pediatric population have not been established [*see Boxed Warning, Warnings and Precautions (5.1)*].

### **8.5 Geriatric Use**

Of the approximately 6,000 subjects who participated in clinical trials with bupropion hydrochloride sustained-release tablets (depression and smoking cessation trials), 275 were ≥65 years and 47 were aged ≥75 years. In addition, several hundred subjects aged ≥65 years participated in clinical trials using



the immediate-release formulation of bupropion (depression trials). No overall differences in safety or effectiveness were observed between these subjects and younger subjects. Reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

Bupropion is extensively metabolized in the liver to active metabolites, which are further metabolized and excreted by the kidneys. The risk of adverse reactions may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, it may be necessary to consider this factor in dose selection; it may be useful to monitor renal function [see *Dosage and Administration (2.3)*, *Use in Specific Populations (8.6)*, *Clinical Pharmacology (12.3)*].

## **8.6 Renal Impairment**

Consider a reduced dose and/or dosing frequency of bupropion hydrochloride tablets in patients with renal impairment (Glomerular Filtration Rate: less than 90 mL per min). Bupropion and its metabolites are cleared renally and may accumulate in such patients to a greater extent than usual. Monitor closely for adverse reactions that could indicate high bupropion or metabolite exposures [see *Dosage and Administration (2.3)*, *Clinical Pharmacology (12.3)*].

## **8.7 Hepatic Impairment**

In patients with moderate to severe hepatic impairment (Child-Pugh score: 7 to 15), the maximum dose of bupropion hydrochloride tablet is 75 mg daily. In patients with mild hepatic impairment (Child-Pugh score: 5 to 6), consider reducing the dose and/or frequency of dosing [see *Dosage and Administration (2.2)*, *Clinical Pharmacology (12.3)*].

# **9 DRUG ABUSE AND DEPENDENCE**

## **9.1 Controlled Substance**

Bupropion is not a controlled substance.

## **9.2 Abuse**

### Humans

Controlled clinical trials conducted in normal volunteers, in subjects with a history of multiple drug abuse, and in depressed subjects showed some increase in motor activity and agitation/excitement, often typical of central stimulant activity.

In a population of individuals experienced with drugs of abuse, a single oral dose of 400 mg of bupropion produced mild amphetamine-like activity as compared with placebo on the Morphine-Benzedrine Subscale of the Addiction Research Center Inventories (ARCI) and a score greater than placebo but less than 15 mg of the Schedule II stimulant dextroamphetamine on the Liking Scale of the ARCI. These scales measure general feelings of euphoria and drug liking which are often associated with abuse potential.

Findings in clinical trials, however, are not known to reliably predict the abuse potential of drugs. Nonetheless, evidence from single-dose trials does suggest that the recommended daily dosage of bupropion when administered orally in divided doses is not likely to be significantly reinforcing to amphetamine or CNS stimulant abusers. However, higher doses (which could not be tested because of the risk of seizure) might be modestly attractive to those who abuse CNS stimulant drugs. Bupropion hydrochloride tablets are intended for oral use only. The inhalation of crushed tablets or injection of dissolved bupropion has been reported. Seizures and/or cases of death have been reported when bupropion has been administered intranasally or by parenteral injection.

### Animals

Studies in rodents and primates demonstrated that bupropion exhibits some pharmacologic actions common to psychostimulants. In rodents, it has been shown to increase locomotor activity, elicit a mild stereotyped behavior response, and increase rates of responding in several schedule-controlled behavior paradigms. In primate models assessing the positive-reinforcing effects of psychoactive drugs, bupropion was self-administered intravenously. In rats, bupropion produced amphetamine-like and cocaine-like discriminative stimulus effects in drug discrimination paradigms used to characterize the subjective effects of psychoactive drugs.

## **10 OVERDOSAGE**

### **10.1 Human Overdose Experience**

Overdoses of up to 30 grams or more of bupropion have been reported. Seizure was reported in approximately one-third of all cases. Other serious reactions reported with overdoses of bupropion alone included hallucinations, loss of consciousness, sinus tachycardia, and ECG changes such as conduction disturbances (including QRS prolongation) or arrhythmias. Fever, muscle rigidity, rhabdomyolysis, hypotension, stupor, coma, and respiratory failure have been reported mainly when bupropion was part of multiple drug overdoses.

Although most patients recovered without sequelae, deaths associated with overdoses of bupropion alone have been reported in patients ingesting large doses of the drug. Multiple uncontrolled seizures, bradycardia, cardiac failure, and cardiac arrest prior to death were reported in these patients.

### **10.2 Overdosage Management**


Consult a Certified Poison Control Center for up-to-date guidance and advice. Telephone numbers for certified poison control centers are listed in the Physician's Desk Reference (PDR). Call 1-800-222-1222 or refer to [www.poison.org](http://www.poison.org).

There are no known antidotes for bupropion. In case of an overdose, provide supportive care, including close medical supervision and monitoring. Consider the possibility of multiple drug overdose. Ensure an adequate airway, oxygenation, and ventilation. Monitor cardiac rhythm and vital signs. Induction of emesis is not recommended.

## **11 DESCRIPTION**

Bupropion hydrochloride, an antidepressant of the aminoketone class, is chemically unrelated to

tricyclic, tetracyclic, selective serotonin re-uptake inhibitor, or other known antidepressant agents. Its structure closely resembles that of diethylpropion; it is related to phenylethylamines. It is designated as ( $\pm$ )-1-(3-chlorophenyl)-2-[(1,1-dimethylethyl)amino]-1-propanone hydrochloride. The molecular weight is 276.2. The molecular formula is  $C_{13}H_{18}ClNO \cdot HCl$ . Bupropion hydrochloride powder is white, crystalline, and highly soluble in water. It has a bitter taste and produces the sensation of local anesthesia on the oral mucosa. The structural formula is:



Bupropion hydrochloride tablets USP is supplied for oral administration as 75 mg (light orange) and 100 mg (light yellow) film-coated tablets. Each tablet contains the labeled amount of bupropion hydrochloride USP and the inactive ingredients: 75 mg - microcrystalline cellulose, hypromellose, low-substituted hydroxypropyl cellulose, magnesium stearate, titanium dioxide, ethyl cellulose, triacetin, iron oxide yellow, iron oxide red, FD&C Red No. 40, 100 mg - microcrystalline cellulose, hypromellose, low-substituted hydroxypropyl cellulose, magnesium stearate, titanium dioxide, ethyl cellulose, triacetin, iron oxide yellow.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

The exact mechanism of the antidepressant action of bupropion is not known, but is presumed to be related to noradrenergic and/or dopaminergic mechanisms. Bupropion is a relatively weak inhibitor of the neuronal reuptake of norepinephrine and dopamine, and does not inhibit the reuptake of serotonin. Bupropion does not inhibit monoamine oxidase.

### 12.3 Pharmacokinetics

Bupropion is a racemic mixture. The pharmacological activity and pharmacokinetics of the individual enantiomers have not been studied. The mean elimination half-life ( $\pm$ SD) of bupropion after chronic dosing is 21 ( $\pm$ 9) hours, and steady-state plasma concentrations of bupropion are reached within 8 days.

#### Absorption

The absolute bioavailability of bupropion hydrochloride tablets in humans has not been determined because an intravenous formulation for human use is not available. However, it appears likely that only a small proportion of any orally administered dose reaches the systemic circulation intact. In rat and dog studies, the bioavailability of bupropion ranged from 5% to 20%.

In humans, following oral administration of bupropion hydrochloride tablets, peak plasma bupropion concentrations are usually achieved within 2 hours. Plasma bupropion concentrations are dose-proportional following single doses of 100 mg to 250 mg; however, it is not known if the proportionality between dose and plasma level is maintained in chronic use.

### Distribution

In vitro tests show that bupropion is 84% bound to human plasma proteins at concentrations up to 200 mcg per mL. The extent of protein binding of the hydroxybupropion metabolite is similar to that for bupropion, whereas the extent of protein binding of the threohydrobupropion metabolite is about half that seen with bupropion.

### Metabolism

Bupropion is extensively metabolized in humans. Three metabolites are active: hydroxybupropion, which is formed via hydroxylation of the *tert*-butyl group of bupropion, and the amino-alcohol isomers threohydrobupropion and erythrohydrobupropion, which are formed via reduction of the carbonyl group. In vitro findings suggest that CYP2B6 is the principal isoenzyme involved in the formation of hydroxybupropion, while cytochrome P450 enzymes are not involved in the formation of threohydrobupropion. Oxidation of the bupropion side chain results in the formation of a glycine conjugate of meta-chlorobenzoic acid, which is then excreted as the major urinary metabolite. The potency and toxicity of the metabolites relative to bupropion have not been fully characterized. However, it has been demonstrated in an antidepressant screening test in mice that hydroxybupropion is one-half as potent as bupropion, while threohydrobupropion and erythrohydrobupropion are 5-fold less potent than bupropion. This may be of clinical importance because the plasma concentrations of the metabolites are as high as or higher than those of bupropion.

Following a single dose in humans, peak plasma concentrations of hydroxybupropion occur approximately 3 hours after administration of bupropion hydrochloride tablets and are approximately 10 times the peak level of the parent drug at steady state. The elimination half-life of hydroxybupropion is approximately 20 ( $\pm 5$ ) hours, and its AUC at steady state is about 17 times that of bupropion. The times to peak concentrations for the erythrohydrobupropion and threohydrobupropion metabolites are similar to that of the hydroxybupropion metabolite. However, their elimination half-lives are longer, 33 ( $\pm 10$ ) and 37 ( $\pm 13$ ) hours, respectively, and steady-state AUCs are 1.5 and 7 times that of bupropion, respectively.

Bupropion and its metabolites exhibit linear kinetics following chronic administration of 300 mg to 450 mg per day.

### Elimination

Following oral administration of 200 mg of  $^{14}\text{C}$ -bupropion in humans, 87% and 10% of the radioactive dose were recovered in the urine and feces, respectively. Only 0.5% of the oral dose was excreted as unchanged bupropion.

### Population Subgroups

Factors or conditions altering metabolic capacity (e.g., liver disease, congestive heart failure [CHF], age, concomitant medications, etc.) or elimination may be expected to influence the degree and extent of accumulation of the active metabolites of bupropion. The elimination of the major metabolites of bupropion may be affected by reduced renal or hepatic function because they are moderately polar compounds and are likely to undergo further metabolism or conjugation in the liver prior to urinary excretion.

**Renal Impairment:** There is limited information on the pharmacokinetics of bupropion in patients with renal impairment. An inter-trial comparison between normal subjects and subjects with end-stage renal failure demonstrated that the parent drug  $C_{\text{max}}$  and AUC values were comparable in the 2 groups, whereas the hydroxybupropion and threohydrobupropion metabolites had a 2.3- and 2.8-fold increase, respectively, in AUC for subjects with end-stage renal failure. A second trial, comparing normal subjects and subjects with moderate-to-severe renal impairment (GFR  $30.9 \pm 10.8$  mL per min) showed that after a single 150 mg dose of sustained-release bupropion, exposure to bupropion was approximately 2-fold higher in subjects with impaired renal function, while levels of the

hydroxybupropion and threo/erythrohydrobupropion (combined) metabolites were similar in the 2 groups. Bupropion is extensively metabolized in the liver to active metabolites, which are further metabolized and subsequently excreted by the kidneys. The elimination of the major metabolites of bupropion may be reduced by impaired renal function. Bupropion hydrochloride tablets should be used with caution in patients with renal impairment and a reduced frequency and/or dose should be considered [see Use in Specific Populations (8.6)].

*Hepatic Impairment:* The effect of hepatic impairment on the pharmacokinetics of bupropion was characterized in 2 single-dose trials, one in subjects with alcoholic liver disease and one in subjects with mild-to-severe cirrhosis. The first trial demonstrated that the half-life of hydroxybupropion was significantly longer in 8 subjects with alcoholic liver disease than in 8 healthy volunteers ( $32 \pm 14$  hours versus  $21 \pm 5$  hours, respectively). Although not statistically significant, the AUCs for bupropion and hydroxybupropion were more variable and tended to be greater (by 53% to 57%) in volunteers with alcoholic liver disease. The differences in half-life for bupropion and the other metabolites in the 2 groups were minimal.

The second trial demonstrated no statistically significant differences in the pharmacokinetics of bupropion and its active metabolites in 9 subjects with mild-to-moderate hepatic cirrhosis compared with 8 healthy volunteers. However, more variability was observed in some of the pharmacokinetic parameters for bupropion ( $AUC$ ,  $C_{max}$ , and  $T_{max}$ ) and its active metabolites ( $t_{1/2}$ ) in subjects with mild-to-moderate hepatic cirrhosis. In subjects with severe hepatic cirrhosis, significant alterations in the pharmacokinetics of bupropion and its metabolites were seen (Table 3).

**Table 3. Pharmacokinetics of Bupropion and Metabolites in Patients with Severe Hepatic Cirrhosis: Ratio Relative to Healthy Matched Controls**

	$C_{max}$	AUC	$t_{1/2}$	$T_{max}^a$
Bupropion	1.69	3.12	1.43	0.5 h
Hydroxybupropion	0.31	1.28	3.88	19 h
Threo/erythrohydrobupropion amino alcohol	0.69	2.48	1.96	20 h

<sup>a</sup> = Difference

*Left Ventricular Dysfunction:* During a chronic dosing trial with bupropion in 14 depressed subjects with left ventricular dysfunction (history of CHF or an enlarged heart on x-ray), there was no apparent effect on the pharmacokinetics of bupropion or its metabolites, compared with healthy volunteers.

*Age:* The effects of age on the pharmacokinetics of bupropion and its metabolites have not been fully characterized, but an exploration of steady-state bupropion concentrations from several depression efficacy trials involving subjects dosed in a range of 300 mg to 750 mg per day, on a 3-times-daily schedule, revealed no relationship between age (18 to 83 years) and plasma concentration of bupropion. A single-dose pharmacokinetic trial demonstrated that the disposition of bupropion and its metabolites in elderly subjects was similar to that of younger subjects. These data suggest there is no prominent effect of age on bupropion concentration; however, another single- and multiple-dose pharmacokinetics trial suggested that the elderly are at increased risk for accumulation of bupropion and its metabolites [see Use in Specific Populations (8.5)].

*Gender:* Pooled analysis of bupropion pharmacokinetic data from 90 healthy male and 90 healthy female volunteers revealed no sex-related differences in the peak plasma concentrations of bupropion. The mean systemic exposure (AUC) was approximately 13% higher in male volunteers compared with female volunteers. The clinical significance of this finding is unknown.

**Smokers:** The effects of cigarette smoking on the pharmacokinetics of bupropion were studied in 34 healthy male and female volunteers; 17 were chronic cigarette smokers and 17 were nonsmokers. Following oral administration of a single 150 mg dose of bupropion, there were no statistically significant differences in  $C_{max}$ , half-life,  $T_{max}$ , AUC, or clearance of bupropion or its active metabolites between smokers and nonsmokers.

### Drug Interactions

**Potential for Other Drugs to Affect Bupropion Hydrochloride Tablets:** In vitro studies indicate that bupropion is primarily metabolized to hydroxybupropion by CYP2B6. Therefore, the potential exists for drug interactions between bupropion hydrochloride tablets and drugs that are inhibitors or inducers of CYP2B6. In addition, in vitro studies suggest that paroxetine, sertraline, norfluoxetine, fluvoxamine, and nelfinavir inhibit the hydroxylation of bupropion.

**Inhibitors of CYP2B6: Ticlopidine, Clopidogrel:** In a trial in healthy male volunteers, clopidogrel 75 mg once daily or ticlopidine 250 mg twice daily increased exposures ( $C_{max}$  and AUC) of bupropion by 40% and 60% for clopidogrel, and by 38% and 85% for ticlopidine, respectively. The exposures ( $C_{max}$  and AUC) of hydroxybupropion were decreased 50% and 52%, respectively, by clopidogrel, and 78% and 84%, respectively, by ticlopidine. This effect is thought to be due to the inhibition of the CYP2B6-catalyzed bupropion hydroxylation.

**Prasugrel:** Prasugrel is a weak inhibitor of CYP2B6. In healthy subjects, prasugrel increased bupropion  $C_{max}$  and AUC values by 14% and 18%, respectively, and decreased  $C_{max}$  and AUC values of hydroxybupropion, an active metabolite of bupropion, by 32% and 24%, respectively.

**Cimetidine:** The threohydrobupropion metabolite of bupropion does not appear to be produced by cytochrome P450 enzymes. The effects of concomitant administration of cimetidine on the pharmacokinetics of bupropion and its active metabolites were studied in 24 healthy young male volunteers. Following oral administration of bupropion 300 mg with and without cimetidine 800 mg, the pharmacokinetics of bupropion and hydroxybupropion were unaffected. However, there were 16% and 32% increases in the AUC and  $C_{max}$ , respectively, of the combined moieties of threohydrobupropion and erythrohydrobupropion.

**Citalopram:** Citalopram did not affect the pharmacokinetics of bupropion and its three metabolites.

**Inducers of CYP2B6: Ritonavir and Lopinavir:** In a healthy volunteer trial, ritonavir 100 mg twice daily reduced the AUC and  $C_{max}$  of bupropion by 22% and 21%, respectively. The exposure of the hydroxybupropion metabolite was decreased by 23%, the threohydrobupropion decreased by 38%, and the erythrohydrobupropion decreased by 48%.

In a second healthy volunteer trial, ritonavir at a dose of 600 mg twice daily decreased the AUC and the  $C_{max}$  of bupropion by 66% and 62%, respectively. The exposure of the hydroxybupropion metabolite was decreased by 78%, the threohydrobupropion decreased by 50%, and the erythrohydrobupropion decreased by 68%.

In another healthy volunteer trial, lopinavir 400 mg/ritonavir 100 mg twice daily decreased bupropion AUC and  $C_{max}$  by 57%. The AUC and  $C_{max}$  of hydroxybupropion were decreased by 50% and 31%, respectively.

**Efavirenz:** In a trial in healthy volunteers, efavirenz 600 mg once daily for 2 weeks reduced the AUC and  $C_{max}$  of bupropion by approximately 55% and 34%, respectively. The AUC of hydroxybupropion was unchanged, whereas  $C_{max}$  of hydroxybupropion was increased by 50%.

**Carbamazepine, Phenobarbital, Phenytoin:** While not systematically studied, these drugs may induce the

metabolism of bupropion.

### Potential for Bupropion Hydrochloride Tablets to Affect Other Drugs

Animal data indicated that bupropion may be an inducer of drug-metabolizing enzymes in humans. In one trial, following chronic administration of bupropion 100 mg three times daily to 8 healthy male volunteers for 14 days, there was no evidence of induction of its own metabolism. Nevertheless, there may be potential for clinically important alterations of blood levels of co-administered drugs.

*Drugs Metabolized by CYP2D6:* In vitro, bupropion and its metabolites (erythrohydrobupropion, threo hydrobupropion, hydroxybupropion) are CYP2D6 inhibitors. In a clinical trial of 15 male subjects (ages 19 to 35 years) who were extensive metabolizers of CYP2D6, bupropion 300 mg per day followed by a single dose of 50 mg desipramine increased the  $C_{max}$ , AUC, and  $t_{1/2}$  of desipramine by an average of approximately 2-, 5-, and 2-fold, respectively. The effect was present for at least 7 days after the last dose of bupropion. Concomitant use of bupropion with other drugs metabolized by CYP2D6 has not been formally studied.

*Citalopram:* Although citalopram is not primarily metabolized by CYP2D6, in one trial bupropion increased the  $C_{max}$  and AUC of citalopram by 30% and 40%, respectively.

*Lamotrigine:* Multiple oral doses of bupropion had no statistically significant effects on the single-dose pharmacokinetics of lamotrigine in 12 healthy volunteers.

*Digoxin:* Literature data showed that digoxin exposure was decreased when a single oral dose of 0.5 mg digoxin was administered 24 hours after a single oral dose of extended-release 150 mg bupropion in healthy volunteers.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

Lifetime carcinogenicity studies were performed in rats and mice at bupropion doses up to 300 mg and 150 mg per kg per day, respectively. These doses are approximately 7 and 2 times the MRHD, respectively, on a mg per  $m^2$  basis. In the rat study there was an increase in nodular proliferative lesions of the liver at doses of 100 mg to 300 mg per kg per day (approximately 2 to 7 times the MRHD on a mg per  $m^2$  basis); lower doses were not tested. The question of whether or not such lesions may be precursors of neoplasms of the liver is currently unresolved. Similar liver lesions were not seen in the mouse study, and no increase in malignant tumors of the liver and other organs was seen in either study.

Bupropion produced a positive response (2 to 3 times control mutation rate) in 2 of 5 strains in the Ames bacterial mutagenicity assay. Bupropion produced an increase in chromosomal aberrations in 1 of 3 *in vivo* rat bone marrow cytogenetic studies.

A fertility study in rats at doses up to 300 mg per kg per day revealed no evidence of impaired fertility.

## **14 CLINICAL STUDIES**

The efficacy of bupropion hydrochloride tablets in the treatment of major depressive disorder was established in two 4-week, placebo-controlled trials in adult inpatients with MDD (Trials 1 and 2 in Table 4) and in one 6-week, placebo-controlled trial in adult outpatients with MDD (Trial 3 in Table 4). In the first trial, the dose range of bupropion hydrochloride tablets was 300 mg to 600 mg per day administered in 3 divided doses; 78% of subjects were treated with doses of 300 mg to 450 mg per day. The trial demonstrated the efficacy of bupropion hydrochloride tablets as measured by the Hamilton Depression Rating Scale (HDRS) total score, the HDRS depressed mood item (Item 1), and the Clinical

Global Impressions-severity score (CGI-S). The second trial included 2 doses of bupropion hydrochloride tablets (300 mg and 450 mg per day) and placebo. This trial demonstrated the effectiveness of bupropion hydrochloride tablets for only the 450 mg per day dose. The efficacy results were statistically significant for the HDRS total score and the CGI-S score, but not for HDRS Item 1. In the third trial, outpatients were treated with 300 mg per day of bupropion hydrochloride tablets. This trial demonstrated the efficacy of bupropion hydrochloride tablets as measured by the HDRS total score, the HDRS Item 1, the Montgomery-Asberg Depression Rating Scale (MADRS), the CGI-S score, and the CGI-Improvement Scale (CGI-I) score. Effectiveness of bupropion hydrochloride tablets in long-term use, that is, for more than 6 weeks, has not been systematically evaluated in controlled trials.

**Table 4. Efficacy of Bupropion Hydrochloride Tablets for the Treatment of Major Depressive Disorder**

Trial number	Treatment Group	Primary Efficacy Measure: HDRS		
		Mean Baseline Score (SD)	LS Mean Score at Endpoint Visit (SE)	Placebo-subtracted Difference <sup>a</sup> (95% CI)
Trial 1	Bupropion Hydrochloride Tablets 300 to 600 mg/day <sup>b</sup> (n = 48)	28.5 (5.1)	14.9 (1.3)	-4.7 (-8.8, -0.6)
	Placebo (n = 27)	29.3 (7)	19.6 (1.6)	--
		Mean Baseline Score (SD)	LS Mean Change from Baseline (SE)	Placebo-subtracted Difference <sup>a</sup> (95% CI)
Trial 2	Bupropion Hydrochloride Tablets 300 mg/day (n = 36)	32.4 (5.9)	-15.5 (1.7)	-4.1
	Bupropion Hydrochloride Tablets 450 mg/day <sup>b</sup> (n = 34)	34.8 (4.6)	-17.4 (1.7)	-5.9 (-10.5, -1.4)
	Placebo (n = 39)	32.9 (5.4)	-11.5 (1.6)	--
Trial 3	Bupropion Hydrochloride Tablets 300 mg/day <sup>b</sup> (n = 110)	26.5 (4.3)	-12 (NA)	-3.9 (-5.7, -1)
	Placebo (n = 106)	27 (3.5)	-8.7 (NA)	--

n: sample size; SD: standard deviation; SE: standard error; LS Mean: least-squares mean; CI: unadjusted confidence interval included for doses that were demonstrated to be effective; NA: not available.

<sup>a</sup> Difference (drug minus placebo) in least-squares estimates with respect to the primary efficacy parameter. For Trial 1, it refers to the mean score at the endpoint visit; for Trials 2 and 3, it refers to the



mean change from baseline to the endpoint visit.

<sup>b</sup> Doses that are demonstrated to be statistically significantly superior to placebo.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

Bupropion hydrochloride tablets USP, 75 mg are light orange colored, round, biconvex film coated tablets debossed with 'L295' on one side and plain on other side.

Bottles of 100 NDC # 46708-118-31

Bottles of 500 NDC # 46708-118-71

Bupropion hydrochloride tablets USP, 100 mg are light yellow colored, round, biconvex film coated tablets debossed with 'L296' on one side and plain on other side.

Bottles of 100 NDC # 46708-127-31

Bottles of 500 NDC # 46708-127-71

**Store at 15° to 25°C (59° to 77°F). Protect from light and moisture.**

## 17 PATIENT COUNSELING INFORMATION

*Advise the patient to read the FDA-approved patient labeling (Medication Guide).*

Inform patients, their families, and their caregivers about the benefits and risks associated with treatment with bupropion hydrochloride tablets and counsel them in its appropriate use.

A patient Medication Guide about "Antidepressant Medicines, Depression and Other Serious Mental Illnesses, and Suicidal Thoughts or Actions," "Quitting Smoking, Quit-Smoking Medications, Changes in Thinking and Behavior, Depression, and Suicidal Thoughts or Actions," and "What Other Important Information Should I Know About Bupropion Hydrochloride Tablets?" is available for bupropion hydrochloride tablets. Instruct patients, their families, and their caregivers to read the Medication Guide and assist them in understanding its contents. Patients should be given the opportunity to discuss the contents of the Medication Guide and to obtain answers to any questions they may have. The complete text of the Medication Guide is reprinted at the end of this document.

Advise patients regarding the following issues and to alert their prescriber if these occur while taking bupropion hydrochloride tablets.

### Suicidal Thoughts and Behaviors

Instruct patients, their families, and/or their caregivers to be alert to the emergence of anxiety, agitation, panic attacks, insomnia, irritability, hostility, aggressiveness, impulsivity, akathisia (psychomotor restlessness), hypomania, mania, other unusual changes in behavior, worsening of depression, and suicidal ideation, especially early during antidepressant treatment and when the dose is adjusted up or down. Advise families and caregivers of patients to observe for the emergence of such symptoms on a day-to-day basis, since changes may be abrupt. Such symptoms should be reported to the patient's prescriber or healthcare professional, especially if they are severe, abrupt in onset, or were not part of the patient's presenting symptoms. Symptoms such as these may be associated with an increased risk for suicidal thinking and behavior and indicate a need for very close monitoring and possibly changes in the medication.

### Neuropsychiatric Adverse Events and Suicide Risk in Smoking Cessation Treatment

Although bupropion hydrochloride tablet is not indicated for smoking cessation treatment, it contains the same active ingredient as ZYBAN which is approved for this use. Inform patients that some patients have experienced changes in mood (including depression and mania), psychosis, hallucinations,

paranoia, delusions, homicidal ideation, aggression, hostility, agitation, anxiety, and panic, as well as suicidal ideation and suicide when attempting to quit smoking while taking bupropion. Instruct patients to discontinue bupropion and contact a healthcare professional if they experience such symptoms [*see Warnings and Precautions (5.2), Adverse Reactions (6.2)*].

### Severe Allergic Reactions

Educate patients on the symptoms of hypersensitivity and to discontinue bupropion hydrochloride tablets if they have a severe allergic reaction.

### Seizure

Instruct patients to discontinue and not restart bupropion hydrochloride tablets if they experience a seizure while on treatment. Advise patients that the excessive use or abrupt discontinuation of alcohol, benzodiazepines, antiepileptic drugs, or sedatives/hypnotics can increase the risk of seizure. Advise patients to minimize or avoid use of alcohol.

### Angle-Closure Glaucoma

Patients should be advised that taking bupropion hydrochloride tablets can cause mild pupillary dilation, which in susceptible individuals, can lead to an episode of angle-closure glaucoma. Pre-existing glaucoma is almost always open-angle glaucoma because angle-closure glaucoma, when diagnosed, can be treated definitively with iridectomy. Open-angle glaucoma is not a risk factor for angle-closure glaucoma. Patients may wish to be examined to determine whether they are susceptible to angle closure, and have a prophylactic procedure (e.g., iridectomy), if they are susceptible [*see Warnings and Precautions (5.7)*].

### Bupropion-Containing Products

Educate patients that bupropion hydrochloride tablets contains the same active ingredient (bupropion hydrochloride) found in ZYBAN, which is used as an aid to smoking cessation treatment, and that bupropion hydrochloride tablets should not be used in combination with ZYBAN or any other medications that contain bupropion (such as WELLBUTRIN SR, the sustained-release formulation and WELLBUTRIN XL or FORFIVO XL, the extended-release formulations, and APLENZIN, the extended-release formulation of bupropion hydrobromide). In addition, there are a number of generic bupropion HCl products for the immediate-, sustained-, and extended-release formulations.

### Potential for Cognitive and Motor Impairment

Advise patients that any CNS-active drug like bupropion hydrochloride tablets may impair their ability to perform tasks requiring judgment or motor and cognitive skills. Advise patients that until they are reasonably certain that bupropion hydrochloride tablets do not adversely affect their performance, they should refrain from driving an automobile or operating complex, hazardous machinery. Bupropion hydrochloride tablets may lead to decreased alcohol tolerance.

### Concomitant Medications

Counsel patients to notify their healthcare provider if they are taking or plan to take any prescription or over-the-counter drugs because bupropion hydrochloride tablets and other drugs may affect each others' metabolisms.

### Pregnancy

Advise patients to notify their healthcare provider if they become pregnant or intend to become pregnant during therapy.

### Precautions for Nursing Mothers

Advise patients that bupropion hydrochloride tablets are present in human milk in small amounts.

### Storage Information

Instruct patients to store bupropion hydrochloride tablets at room temperature, between 15° to 25°C (59° to 77°F) and keep the tablets dry and out of the light.

### Administration Information

Instruct patients to take bupropion hydrochloride tablets in equally divided doses 3 or 4 times a day, with doses separated by at least 6 hours to minimize the risk of seizure. Instruct patients if they miss a

dose, not to take an extra tablet to make up for the missed dose and to take the next tablet at the regular time because of the dose-related risk of seizure. Instruct patients that bupropion hydrochloride tablets should be swallowed whole and not crushed, divided, or chewed. Bupropion hydrochloride tablets can be taken with or without food.

Trademarks are owned by or licensed to the GSK group of companies. The other brands listed are trademarks owned by or licensed to their respective owners and are not owned by or licensed to the GSK group of companies. The makers of these brands are not affiliated with and do not endorse the GSK group of companies or its products.

Manufactured by:

Alembic Pharmaceuticals Limited (Formulation Division),  
Village Panelav, P. O. Tajpura, Near Baska,  
Taluka-Halol, Panchmahal 389350, Gujarat, India.

Revised: 06/2018

## Medication Guide

### Bupropion Hydrochloride (bue-PROE-pee-on HYE-droe-KLOR-ide) Tablets USP

**IMPORTANT: Be sure to read the three sections of this Medication Guide. The first section is about the risk of suicidal thoughts and actions with antidepressant medicines; the second section is about the risk of changes in thinking and behavior, depression and suicidal thoughts or actions with medicines used to quit smoking; and the third section is entitled “What Other Important Information Should I Know About Bupropion Hydrochloride Tablets?”**

#### **Antidepressant Medicines, Depression and Other Serious Mental Illnesses, and Suicidal Thoughts or Actions**

This section of the Medication Guide is only about the risk of suicidal thoughts and actions with antidepressant medicines.

**What is the most important information I should know about antidepressant medicines, depression and other serious mental illnesses, and suicidal thoughts or actions?**

**1. Antidepressant medicines may increase the risk of suicidal thoughts or actions in some children, teenagers, or young adults within the first few months of treatment.**

**2. Depression or other serious mental illnesses are the most important causes of suicidal thoughts and actions. Some people may have a particularly high risk of having suicidal thoughts or actions. These include people who have (or have a family history of) bipolar illness (also called manic-depressive illness) or suicidal thoughts or actions.**

**3. How can I watch for and try to prevent suicidal thoughts and actions in myself or a family member?**

- Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or feelings. This is very important when an antidepressant medicine is started or when the dose is changed.
- Call your healthcare provider right away to report new or sudden changes in mood, behavior, thoughts, or feelings.
- Keep all follow-up visits with your healthcare provider as scheduled. Call the healthcare provider between visits as needed, especially if you have concerns about symptoms.

**Call your healthcare provider right away if you or your family member has any of the following symptoms, especially if they are new, worse, or worry you:**

• thoughts about suicide or dying	• trouble sleeping (insomnia)
-----------------------------------	-------------------------------

•attempts to commit suicide	•new or worse irritability
•new or worse depression	•acting aggressive, being angry, or violent
•new or worse anxiety	•acting on dangerous impulses
•feeling very agitated or restless	•an extreme increase in activity and talking (mania)
•panic attacks	•other unusual changes in behavior or mood

### What else do I need to know about antidepressant medicines?

- Never stop an antidepressant medicine without first talking to a healthcare provider.** Stopping an antidepressant medicine suddenly can cause other symptoms.
- Antidepressants are medicines used to treat depression and other illnesses. It is important to discuss all the risks of treating depression and also the risks of not treating it. Patients and their families or other caregivers should discuss all treatment choices with the healthcare provider, not just the use of antidepressants.
- Antidepressant medicines have other side effects.** Talk to the healthcare provider about the side effects of the medicine prescribed for you or your family member.
- Antidepressant medicines can interact with other medicines.** Know all of the medicines that you or your family member takes. Keep a list of all medicines to show the healthcare provider. Do not start new medicines without first checking with your healthcare provider.

It is not known if bupropion hydrochloride tablets are safe and effective in children under the age of 18.

### Quitting Smoking, Quit-Smoking Medications, Changes in Thinking and Behavior, Depression, and Suicidal Thoughts or Actions

This section of the Medication Guide is only about the risk of changes in thinking and behavior, depression and suicidal thoughts or actions with drugs used to quit smoking. Although bupropion hydrochloride tablets are not a treatment for quitting smoking, it contains the same active ingredient (bupropion hydrochloride) as ZYBAN which is used to help patients quit smoking.

#### **Talk to your healthcare provider or your family member's healthcare provider about:**

- all risks and benefits of quit-smoking medicines.
- all treatment choices for quitting smoking.

When you try to quit smoking, with or without bupropion, you may have symptoms that may be due to nicotine withdrawal, including:

•urge to smoke	•frustration	•restlessness
•depressed mood	•anger	• decreased heart rate
•trouble sleeping	• feeling anxious	•increased appetite
•irritability	•difficulty concentrating	•weight gain

Some people have even experienced suicidal thoughts when trying to quit smoking without medication.

Sometimes quitting smoking can lead to worsening of mental health problems that you already have, such as depression.

Some people have had serious side effects while taking bupropion to help them quit smoking, including: **New or worse mental health problems, such as changes in behavior or thinking, aggression, hostility, agitation, depression, or suicidal thoughts or actions.** Some people had these symptoms when they began taking bupropion, and others developed them after several weeks of treatment, or after stopping bupropion. These symptoms happened more often in people who had a history of mental health problems before taking bupropion than in people without a history of mental health problems.

**Stop taking bupropion hydrochloride tablets and call your healthcare provider right away if you, your family, or caregiver notice any of these symptoms.** Work with your healthcare provider to decide whether you should continue to take bupropion hydrochloride tablets. In many people, these symptoms went away after stopping bupropion hydrochloride tablets, but in some people symptoms continued after stopping bupropion hydrochloride tablets. It is important for you to follow-up with your healthcare provider until your symptoms go away.

**Before taking bupropion hydrochloride tablets,** tell your healthcare provider if you have ever had depression or other mental health problem. You should also tell your healthcare provider about any symptoms you had during other times you tried to quit smoking, with or without bupropion.

### **What Other Important Information Should I Know About Bupropion Hydrochloride Tablets?**

**•Seizures: There is a chance of having a seizure (convulsion, fit) with bupropion hydrochloride tablets, especially in people:**

- o with certain medical problems.
- o who take certain medicines.

The chance of having seizures increases with higher doses of bupropion hydrochloride tablets. For more information, see the sections “Who should not take bupropion hydrochloride tablets?” and “What should I tell my healthcare provider before taking bupropion hydrochloride tablets?” Tell your healthcare provider about all of your medical conditions and all the medicines you take. **Do not take any other medicines while you are taking bupropion hydrochloride tablets unless your healthcare provider has said it is okay to take them.**

**If you have a seizure while taking bupropion hydrochloride tablets, stop taking the tablets and call your healthcare provider right away.** Do not take bupropion hydrochloride tablets again if you have a seizure.

**•High blood pressure (hypertension). Some people get high blood pressure that can be severe, while taking bupropion hydrochloride tablets.** The chance of high blood pressure may be higher if you also use nicotine replacement therapy (such as a nicotine patch) to help you stop smoking (see the section of this Medication Guide called “How should I take bupropion hydrochloride tablets?”).

**•Manic episodes.** Some people may have periods of mania while taking bupropion hydrochloride tablets, including:

- o Greatly increased energy
- o Severe trouble sleeping
- o Racing thoughts
- o Reckless behavior
- o Unusually grand ideas
- o Excessive happiness or irritability
- o Talking more or faster than usual

If you have any of the above symptoms of mania, call your healthcare provider.

**•Unusual thoughts or behaviors.** Some patients have unusual thoughts or behaviors while taking bupropion hydrochloride tablets, including delusions (believe you are someone else), hallucinations (seeing or hearing things that are not there), paranoia (feeling that people are against you), or feeling confused. If this happens to you, call your healthcare provider.

**•Visual problems.**

- eye pain
- changes in vision
- swelling or redness in or around the eye

Only some people are at risk for these problems. You may want to undergo an eye examination to see if you are at risk and receive preventative treatment if you are.

**•Severe allergic reactions. Some people can have severe allergic reactions to bupropion hydrochloride tablets. Stop taking bupropion hydrochloride tablets and call your healthcare provider right away** if you get a rash, itching, hives, fever, swollen lymph glands, painful sores in the mouth or around the eyes, swelling of the lips or tongue, chest pain, or have trouble breathing. These could be signs of a serious allergic reaction.

### **What is bupropion hydrochloride tablet?**

Bupropion hydrochloride tablet is a prescription medicine used to treat adults with a certain type of

depression called major depressive disorder.

### **Who should not take bupropion hydrochloride tablets?**

#### **Do not take bupropion hydrochloride tablets if you**

- have or had a seizure disorder or epilepsy.
- have or had an eating disorder such as anorexia nervosa or bulimia.
- **are taking any other medicines that contain bupropion, including ZYBAN (used to help people stop smoking) WELLBUTRIN SR, WELLBUTRIN XL, APLENZIN, FORFIVO XL.** Bupropion is the same active ingredient that is in bupropion hydrochloride tablets.
- drink a lot of alcohol and abruptly stop drinking, or use medicines called sedatives (these make you sleepy), or benzodiazepines, or anti-seizure medicines, and you stop using them all of a sudden.
- take a monoamine oxidase inhibitor (MAOI). Ask your healthcare provider or pharmacist if you are not sure if you take an MAOI, including the antibiotic linezolid.
  - o do not take an MAOI within 2 weeks of stopping bupropion hydrochloride tablets unless directed to do so by your healthcare provider.
  - o do not start bupropion hydrochloride tablets if you stopped taking an MAOI in the last 2 weeks unless directed to do so by your healthcare provider.
- are allergic to the active ingredient in bupropion hydrochloride tablets, bupropion, or to any of the inactive ingredients. See the end of this Medication Guide for a complete list of ingredients in bupropion hydrochloride tablets.

### **What should I tell my healthcare provider before taking bupropion hydrochloride tablets?**

Tell your healthcare provider if you have ever had depression, suicidal thoughts or actions, or other mental health problems. See “Antidepressant Medicines, Depression and Other Serious Mental Illnesses, and Suicidal Thoughts or Actions.”

#### **• Tell your healthcare provider about your other medical conditions including if you:**

- o have liver problems, especially cirrhosis of the liver.
- o have kidney problems.
- o have, or have had, an eating disorder, such as anorexia nervosa or bulimia.
- o have had a head injury.
- o have had a seizure (convulsion, fit).
- o have a tumor in your nervous system (brain or spine).
- o have had a heart attack, heart problems, or high blood pressure.
- o are a diabetic taking insulin or other medicines to control your blood sugar.
- o drink alcohol.
- o abuse prescription medicines or street drugs.
- o are pregnant or plan to become pregnant.
- o are breastfeeding. Bupropion hydrochloride passes into your milk in small amounts.

**Tell your healthcare provider about all the medicines you take**, including prescription, over-the-counter medicines, vitamins, and herbal supplements. Many medicines increase your chances of having seizures or other serious side effects if you take them while you are taking bupropion hydrochloride tablets.

### **How should I take bupropion hydrochloride tablets?**

- Take bupropion hydrochloride tablets exactly as prescribed by your healthcare provider. Do not change your dose or stop taking bupropion hydrochloride tablets without talking with your healthcare provider first.
- **Swallow bupropion hydrochloride tablets whole. Do not chew, cut, or crush bupropion hydrochloride tablets.**
- Take bupropion hydrochloride tablets at the same time each day.
- Take your doses of bupropion hydrochloride tablets at least 6 hours apart.
- You may take bupropion hydrochloride tablets with or without food.
- If you miss a dose, do not take an extra dose to make up for the dose you missed. Wait and take your

next dose at the regular time. **This is very important.** Too much bupropion hydrochloride tablets can increase your chance of having a seizure.

- If you take too much bupropion hydrochloride tablets, or overdose, call your local emergency room or poison control center right away.

- Do not take any other medicines while taking bupropion hydrochloride tablets unless your healthcare provider has told you it is okay.**

- If you are taking bupropion hydrochloride tablets for the treatment of major depressive disorder, it may take several weeks for you to feel that bupropion hydrochloride tablets are working. Once you feel better, it is important to keep taking bupropion hydrochloride tablets exactly as directed by your healthcare provider. Call your healthcare provider if you do not feel bupropion hydrochloride tablets are working for you.

### **What should I avoid while taking bupropion hydrochloride tablets?**

- Limit or avoid using alcohol during treatment with bupropion hydrochloride tablets. If you usually drink a lot of alcohol, talk with your healthcare provider before suddenly stopping. If you suddenly stop drinking alcohol, you may increase your risk of having seizures.

- Do not drive a car or use heavy machinery until you know how bupropion hydrochloride tablets affect you. Bupropion hydrochloride tablets can affect your ability to do these things safely.

### **What are the possible side effects of bupropion hydrochloride tablets?**

Bupropion hydrochloride tablets can cause serious side effects. See the sections at the beginning of this Medication Guide for information about serious side effects of bupropion hydrochloride tablets. The most common side effects of bupropion hydrochloride tablets include:

•nervousness	•heavy sweating
•dry mouth	•shakiness (tremor)
•constipation	•trouble sleeping
•headache	•blurred vision
•nausea or vomiting	•fast heartbeat
•dizziness	

If you have nausea, take your medicine with food. If you have trouble sleeping, do not take your medicine too close to bedtime.

Tell your healthcare provider right away about any side effects that bother you.

These are not all the possible side effects of bupropion hydrochloride tablets. For more information, ask your healthcare provider or pharmacist.

Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### **How should I store bupropion hydrochloride tablets?**

- Store bupropion hydrochloride tablets at room temperature between 15° to 25°C (59° to 77°F).
- Keep bupropion hydrochloride tablets dry and out of the light.

**Keep bupropion hydrochloride tablets and all medicines out of the reach of children.**

### **General information about the safe and effective use of bupropion hydrochloride tablets**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use bupropion hydrochloride tablets for a condition for which it was not prescribed. Do not give bupropion hydrochloride tablets to other people, even if they have the same symptoms you have. It may harm them.

If you take a urine drug screening test, bupropion hydrochloride tablets, may make the test result positive for amphetamines. If you tell the person giving you the drug screening test that you are taking

bupropion hydrochloride tablets, they can do a more specific drug screening test that should not have this problem.

This Medication Guide summarizes important information about bupropion hydrochloride tablets. If you would like more information, talk with your healthcare provider. You can ask your healthcare provider or pharmacist for information about bupropion hydrochloride tablets that is written for healthcare professionals.

**What are the ingredients in bupropion hydrochloride tablets?**

Active ingredient: bupropion hydrochloride

Inactive ingredients: 75 mg (light orange) and 100 mg (light yellow) film-coated tablets. Each tablet contains the labeled amount of bupropion hydrochloride and the inactive ingredients: 75 mg - microcrystalline cellulose, hypromellose, low-substituted hydroxypropyl cellulose, magnesium stearate, titanium dioxide, ethyl cellulose, triacetin, iron oxide yellow, iron oxide red, FD&C Red No. 40, 100 mg - microcrystalline cellulose, hypromellose, low-substituted hydroxypropyl cellulose, magnesium stearate, titanium dioxide, ethyl cellulose, triacetin, iron oxide yellow.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Trademarks are owned by or licensed to the GSK group of companies. The other brands listed are trademarks owned by or licensed to their respective owners and are not owned by or licensed to the GSK group of companies. The makers of these brands are not affiliated with and do not endorse the GSK group of companies or its products.

Manufactured by:

Alembic Pharmaceuticals Limited (Formulation Division),  
Village Panelav, P. O. Tajpura, Near Baska,  
Taluka-Halol, Panchmahal 389350, Gujarat, India.

Revised: 06/2018

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL - 75 mg**

Bupropion Hydrochloride Tablets USP 75 mg (100 Tablets in one Bottle)

Each film-coated tablet contains 75 mg bupropion HCl USP.

46708-118-31



**Each film-coated tablet contains:**  
bupropion HCl USP.....75 mg

**Usual Dosage:** See prescribing information for dosage information.

Store at 15° to 25°C (59° to 77°F). Protect from light and moisture.

Zyban® is a registered trademark of GlaxoSmithKline.

Manufactured by:  
Alembic Pharmaceuticals Limited (Formulation Division),  
Village Panelav, P. O. Tajpura, Near Baska, Taluka-Halol, Panchmahal 389350, Gujarat, India.


**NDC 46708-118-31**  
**Three Times Daily (After Initial Titration)**  
**buPROPion Hydrochloride Tablets, USP**  
**75 mg**

**WARNING:** Do not use in combination with Zyban® or any other medicines that contain bupropion HCl

**PHARMACIST: DISPENSE WITH MEDICATION GUIDE PROVIDED SEPARATELY.**

**100 Tablets Rx only**

**Alembic**

**N 3**  **46708-118-31**  
Mfg. Lic. No. G/959

LOT:   
EXP:   
11/2017

**PACKAGE LABEL PRINCIPAL DISPLAY PANEL - 100 mg**

Bupropion Hydrochloride Tablets, USP 100 mg (100 Tablets in one Bottle)  
Each film-coated tablet contains: bupropion HCl USP .....100 mg .  
46708-127-31

**Each film-coated tablet contains:**  
bupropion HCl USP.....100 mg

**Usual Dosage:** See prescribing information for dosage information.

Store at 15° to 25°C (59° to 77°F). Protect from light and moisture.

Zyban® is a registered trademark of GlaxoSmithKline.

Manufactured by:  
Alembic Pharmaceuticals Limited (Formulation Division),  
Village Panelav, P. O. Tajpura, Near Baska, Taluka-Halol, Panchmahal 389350, Gujarat, India.


**NDC 46708-127-31**  
**Three Times Daily (After Initial Titration)**  
**buPROPion Hydrochloride Tablets, USP**  
**100 mg**

**WARNING:** Do not use in combination with Zyban® or any other medicines that contain bupropion HCl

**PHARMACIST: Dispense with medication guide provided separately.**

**100 Tablets Rx only**

**Alembic**

**N 3**  **46708-127-31**  
Mfg. Lic. No. G/959

LOT:   
EXP:   
11/2017

**BUPROPION HYDROCHLORIDE**

bupropion hydrochloride tablet, film coated

**Product Information**

**Product Type**

HUMAN PRESCRIPTION DRUG

**Item Code (Source)**

NDC:46708-118

Route of Administration ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BUPROPION HYDROCHLORIDE (UNII: ZG7E5POY8O) (BUPROPION - UNII:01ZG3TPX31)	BUPROPION HYDROCHLORIDE	75 mg

### Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
LOW-SUBSTITUTED HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 2165RE0K14)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
ETHYLCELLULOSE, UNSPECIFIED (UNII: 7Z8S9VYZ4B)	
TRIACETIN (UNII: XHX3C3X673)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	

### Product Characteristics

Color	ORANGE (LIGHT ORANGE)	Score	no score
Shape	ROUND (ROUND)	Size	11mm
Flavor		Imprint Code	L295
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46708-118-31	100 in 1 BOTTLE; Type 0: Not a Combination Product	06/18/2018	
2	NDC:46708-118-71	500 in 1 BOTTLE; Type 0: Not a Combination Product	06/18/2018	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA203013	06/18/2018	

## BUPROPION HYDROCHLORIDE

bupropion hydrochloride tablet, film coated

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:46708-127
Route of Administration	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>BUPROPION HYDROCHLORIDE</b> (UNII: ZG7E5POY8O) (BUPROPION - UNII:01ZG3TPX31)	BUPROPION HYDROCHLORIDE	100 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>LOW-SUBSTITUTED HYDROXYPROPYL CELLULOSE, UNSPECIFIED</b> (UNII: 2165RE0K14)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>ETHYLCELLULOSE, UNSPECIFIED</b> (UNII: 7Z8S9VYZ4B)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	

**Product Characteristics**

<b>Color</b>	YELLOW (LIGHT YELLOW)	<b>Score</b>	no score
<b>Shape</b>	ROUND (ROUND)	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	L296
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46708-127-31	100 in 1 BOTTLE; Type 0: Not a Combination Product	06/18/2018	
2	NDC:46708-127-71	500 in 1 BOTTLE; Type 0: Not a Combination Product	06/18/2018	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA203013	06/18/2018	

**Labeler** - Alembic Pharmaceuticals Limited (650574663)**Establishment**

Name	Address	ID/FEI	Business Operations
Alembic Pharmaceuticals Limited		650574671	MANUFACTURE(46708-118, 46708-127)