

**COLD AND FLU DAYTIME- acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled  
Safeway, Inc.**

-----

**Drug Facts**

**Active ingredients (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 10 mg

Phenylephrine HCL 5 mg

**Purpose**

**Pain reliever/fever reducer**

Cough suppressant

Nasal decongestant

**Uses**

- temporarily relieves common cold and flu symptoms:
  - minor aches & pains
  - headache
  - sore throat
  - nasal congestion
  - fever
  - cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- diabetes
- heart disease
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

**When using this product**

**do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- nervousness, dizziness or sleeplessness occur
- fever gets worse or last more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause serious liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults and for children even if you do not notice any signs or symptoms

**Directions**

- **do not take more than directed (see overdose warning)**
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 4 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- **when using other Daytime or Nighttime products, carefully read each label to insure correct dosing**

**Other information**

- store between 15– 30°C (59-86°F)
- avoid excessive heat

**Inactive ingredients**

butylated hydroxyanisole, butylated hydroxytoluene, FD&C Yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

**Questions or comments?**

Call **1-888-723-3929** Monday- Friday 7AM- 6PM PST

**Principal Display Panel**

Compare to Vicks® DayQuil® Cold & Flu LiquiCaps® active ingredients†

**Non-Drowsy Daytime****Cold & Flu Relief**

**ACETAMINOPHEN 325mg** - Pain Reliever/Fever Reducer

**DEXTROMETHORPHAN HBr 10 mg** - Cough Suppressant

**PHENYLEPHRINE HCL 5mg** - Nasal Decongestant

- Non-drowsy
- Alcohol-free
- Antihistamine-free

**SOFTGELS\*\***

(\*\*Liquid-filled capsules)

†This product is not manufactured or distributed by The Procter & Gamble Company, Vicks®, DayQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

DISTRIBUTED BY: BETTER LIVING BRANDS LLC

P.O. BOX 99, PLEASANTON, CA 94566-0009

[www.betterlivingbrandsLLC.com](http://www.betterlivingbrandsLLC.com)

**Product Label**

<p><b>Questions or comments?</b> Call 1-877-753-3935 Monday-Friday 9AM-5PM EST</p>	<p><b>When using this product, do not exceed recommended dosage.</b> thinning drug warfarin.</p> <p><b>Ask a doctor or pharmacist before use if you are taking the blood</b></p> <p>■ cough that occurs with too much phlegm (mucus) ■ emphysema ■ persistent or chronic cough such as occurs with smoking, asthma, or ■ trouble urinating due to an enlarged prostate gland</p> <p>■ heart disease ■ thyroid disease ■ high blood pressure</p> <p><b>Ask a doctor before use if you have</b> ■ liver disease ■ diabetes</p> <p>pharmacist before taking this product.</p> <p>do not know if your prescription drug contains an MAOI, ask a doctor or (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) acetylaminophen, ask a doctor or pharmacist.</p> <p>■ with any other drug containing acetylaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetylaminophen, ask a doctor or pharmacist.</p> <p><b>Do not use</b> vomiting, consult a doctor promptly.</p> <p>days, is accompanied or followed by fever, headache, rash, nausea, or ■ 3 or more alcoholic drinks every day while using this product</p> <p><b>Sore throat warning:</b> If sore throat is severe, persists for more than 2 If a skin reaction occurs, stop use and seek medical help right away. may include: ■ skin reddening ■ blisters ■ rash</p> <p><b>Allergy alert:</b> Acetylaminophen may cause severe skin reactions. Symptoms ■ with other drugs containing acetylaminophen ■ more than 4,000 mg of acetylaminophen in 24 hours may occur if you take: <b>Liver warning:</b> This product contains acetylaminophen. Severe liver damage</p>
<p><b>Inactive ingredients</b> butylated hydroxyanisole, butylated hydroxytoluene, FD&amp;C yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink</p> <p><b>Other information</b> ■ store between 15-30°C (59-86°F) ■ avoid excessive heat</p> <p><b>Directions</b> ■ do not take more than directed (see Overdose warning) ■ adults and children 12 years and over: take 2 softgels with water every 4 hours ■ do not take more than 4 doses in 24 hours ■ swallow whole; do not crush, chew, or dissolve ■ children under 12 years: do not use ■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing</p>	<p><b>Warnings</b> Liver warning: This product contains acetylaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetylaminophen in 24 hours ■ with other drugs containing acetylaminophen ■ 3 or more alcoholic drinks every day while using this product</p> <p><b>Allergy alert:</b> Acetylaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash</p> <p><b>Sore throat warning:</b> If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p><b>Do not use</b> ■ with any other drug containing acetylaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetylaminophen, ask a doctor or pharmacist.</p> <p>■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p><b>Ask a doctor before use if you have</b> ■ liver disease ■ diabetes</p> <p>■ heart disease ■ thyroid disease ■ high blood pressure</p> <p>■ trouble urinating due to an enlarged prostate gland</p> <p>■ persistent or chronic cough such as occurs with smoking, asthma, or ■ emphysema</p> <p>■ cough that occurs with too much phlegm (mucus)</p> <p><b>Ask a doctor or pharmacist before use if you are taking the blood</b> thinning drug warfarin.</p> <p><b>When using this product, do not exceed recommended dosage.</b></p>
<p><b>Drug Facts (continued)</b></p> <p><b>Stop use and ask a doctor if</b> ■ pain, cough, or nasal congestion gets worse or lasts more than 7 days ■ nervousness, dizziness, or sleeplessness occur ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.</p> <p><b>If pregnant or breast-feeding, ask a health professional before use.</b> <b>Keep out of reach of children.</b> Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Warnings</b> Liver warning: This product contains acetylaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetylaminophen in 24 hours ■ with other drugs containing acetylaminophen ■ 3 or more alcoholic drinks every day while using this product</p> <p><b>Allergy alert:</b> Acetylaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash</p> <p><b>Sore throat warning:</b> If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p><b>Do not use</b> ■ with any other drug containing acetylaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetylaminophen, ask a doctor or pharmacist.</p> <p>■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p><b>Ask a doctor before use if you have</b> ■ liver disease ■ diabetes</p> <p>■ heart disease ■ thyroid disease ■ high blood pressure</p> <p>■ trouble urinating due to an enlarged prostate gland</p> <p>■ persistent or chronic cough such as occurs with smoking, asthma, or ■ emphysema</p> <p>■ cough that occurs with too much phlegm (mucus)</p> <p><b>Ask a doctor or pharmacist before use if you are taking the blood</b> thinning drug warfarin.</p> <p><b>When using this product, do not exceed recommended dosage.</b></p>

<p><b>Drug Facts (continued)</b></p> <p><b>Uses</b> ■ temporarily relieves common cold and flu symptoms: ■ minor aches and pains ■ headache ■ sore throat ■ fever ■ nasal congestion ■ cough due to minor throat and bronchial irritation</p>	<p><b>Drug Facts</b></p> <p><b>Active ingredients (in each softgel)</b> Acetaminophen 325 mg.....Pain reliever/fever reducer Dextromethorphan HBr 10 mg.....Cough suppressant Phenylephrine HCl 5 mg.....Nasal decongestant</p>
---	---



**Compare to**  
**Vicks® DayQuil®**  
**Cold & Flu LiquiCaps®**  
active ingredients†

NDC 21130-650-24

# Non-Drowsy Daytime Cold & Flu Relief

**ACETAMINOPHEN** 325 mg - Pain Reliever/Fever Reducer  
**DEXTROMETHORPHAN HBr** 10 mg - Cough Suppressant  
**PHENYLEPHRINE HCl** 5 mg - Nasal Decongestant



- Non-drowsy
- Alcohol-free
- Antibiotamine-free

**24 SOFTGELS\*\***

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.  
KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

at No.:  
xp. Date:

Antihistamine-free

(\*\*Liquid-filled capsules)

DISTRIBUTED BY: BETTER LIVING BRANDS LLC  
P.O. BOX 99, PLEASANTON, CA 94566-0009  
1-888-723-3929  
www.betterlivingbrandsLLC.com

This product is not manufactured or distributed by The Procter & Gamble Company.  
Vicks®, DayQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.

OUR PROMISE

QUALITY & SATISFACTION

100% GUARANTEED

OR YOUR MONEY BACK

S2801

RD21343

3 2 1130 76334 5

Scan here for more information

PLD-H568B

FC008030

SIGNATURE CARE Non Drowsy Daytime Cold & Flu Relief

COLD AND FLU DAYTIME			
acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:21130-650
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)		DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)		PHENYLEPHRINE HYDROCHLORIDE	5 mg
Inactive Ingredients			
Ingredient Name			Strength
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)			
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)			
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
GELATIN (UNII: 2G86QN327L)			
GLYCERIN (UNII: PDC6A3C0OX)			
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)			
POVIDONE (UNII: FZ989GH94E)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
WATER (UNII: 059QF0KO0R)			
SORBITAN (UNII: 6O92ICV9RU)			
SORBITOL (UNII: 506T60A25R)			
MANNITOL (UNII: 3OWL53L36A)			
Product Characteristics			
Color	orange	Score	no score
Shape	CAPSULE	Size	20mm

Flavor		Imprint Code		P19
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-650-24	24 in 1 CARTON	06/30/2019	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M012	06/30/2019	

**Labeler** - Safeway, Inc. (009137209)