

**DG HEALTH TUSSIN CF ADULT COUGH AND COLD- dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution**  
**Dolgencorp, LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Dolgencorp, LLC Tussin CF Drug Facts**

**Active ingredients (in each 10 mL)**

Dextromethorphan HBr, USP 20 mg

Guaifenesin, USP 200 mg

Phenylephrine HCl, USP 10 mg

**Purposes**

Cough suppressant

Expectorant

Nasal decongestant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
- nasal congestion
- cough due to minor throat and bronchial irritation

**Warnings**

**Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- diabetes
- high blood pressure
- thyroid disease
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland

**When using this product**

do not use more than directed

**Stop use and ask a doctor if**

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	10 mL every 4 hours
children under 12 years	do not use

**Other information**

- each 10 mL contains: sodium 5 mg
- store at 20-25°C (68-77°F).  
Do not refrigerate.

**Inactive ingredients**

anhydrous citric acid, edetate disodium, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

**Questions or comments?**

1-888-309-9030

**Principal Display Panel**

Compare to active ingredients of Robitussin® Multi-Symptom Cold

For Ages 12 & Over

Adult

Tussin

Cough Suppressant (Dextromethorphan HBr)

Expectorant (Guaifenesin)

Nasal Decongestant (Phenylephrine HCl)

Multi-Symptom Cold

Relieves:

Cough

Nasal congestion

Mucus

Non-Drowsy

Peak Cold

4 FL OZ (118 mL)

**DG™** health

Compare  
to active  
ingredients of  
**Robitussin®**  
Multi-Symptom  
Cold\*

For Ages 12 & Over  
**Adult**

# Tussin

**Cough Suppressant**  
(Dextromethorphan HBr)  
**Expectorant** (Guaifenesin)  
**Nasal Decongestant**  
(Phenylephrine HCl)

**Multi-Symptom  
Cold**

Relieves:

- Cough
- Nasal congestion
- Mucus

Non Drowsy

Peak Cold



4 FL OZ (118 mL)

**DG™** health

Compare  
to active  
ingredients of  
**Robitussin®**  
Multi-Symptom  
Cold\*

For Ages 12 & Over  
**Adult**

# Tussin

**Cough Suppressant**  
(Dextromethorphan HBr)  
**Expectorant** (Guaifenesin)  
**Nasal Decongestant**  
(Phenylephrine HCl)

**Multi-Symptom  
Cold**

Relieves:

- Cough
- Nasal congestion
- Mucus

Non Drowsy


Peak Cold



4 FL OZ (118 mL)



For Ages 12 & Over  
**Tussin**  
 Adult  
 Cough Suppressant  
 (Dextromethorphan HBr)  
 Expectorant (Guaifenesin)  
 Nasal Decongestant  
 (Phenylephrine HCl)



## Drug Facts

### Active ingredients (in each 10 mL) Purposes

Dextromethorphan HBr,  
 USP 20 mg.....Cough suppressant  
 Guaifenesin, USP 200 mg.....Expectorant  
 Phenylephrine HCl,  
 USP 10 mg.....Nasal decongestant

### Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
  - nasal congestion
  - cough due to minor throat and bronchial irritation

### Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

When using this product do not use more than directed

### Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition. ▶

## Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.  
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	10 mL every 4 hours
children under 12 years	do not use

### Other information

- each 10 mL contains: sodium 5 mg
- store at 20-25°C (68-77°F).  
Do not refrigerate.

**Inactive ingredients** anhydrous citric acid, edetate disodium, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

### Questions or comments?

1-888-309-9030

\*This product is not manufactured or distributed by Pfizer, distributor of Robitussin® Multi-Symptom Cold.

DISTRIBUTED BY  
 DOLGENCORP, LLC  
 100 MISSION RIDGE  
 GOODLETTSVILLE,  
 TN 37072 USA


**100%**  
**Satisfaction**  
**Guaranteed!**  
 (888) 309-9030

**DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING**

**PARENTS:**

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)



**DG HEALTH TUSSIN CF ADULT COUGH AND COLD**  
 dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:559 10-516
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 10 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 10 mL

Inactive Ingredients	
Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

**Product Characteristics**

<b>Color</b>	RED (clear)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:559 10-516-26	1 in 1 CARTON	05/22/2010	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	05/22/2010	

**Labeler** - Dolgencorp, LLC (068331990)

Revised: 12/2019

Dolgencorp, LLC