

**VICKS DAYQUIL AND VICKS NYQUIL COLD AND FLU- acetaminophen,
phenylephrine hydrochloride, dextromethorphan hydrobromide, and
doxylamine succinate**
The Procter & Gamble Manufacturing Company

**Vicks[®] DayQuil[®] and Vicks[®] NyQuil[®] COLD & FLU
LiquiCaps[™] Convenience Pack**

DayQuil Cold & Flu

Drug Facts

Active ingredients (in each LiquiCap)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/Fever reducer
Cough suppressant
Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

Warnings

Liver warning:

This product contains acetaminophen.

Severe liver damage may occur if you take

- more than 8 LiquiCaps in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening

- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough get worse or last more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 8 LiquiCaps per 24 hrs

adults & children 12 yrs & over	2 LiquiCaps with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- store at no greater than 25°C

Inactive ingredients

FD&C Red No. 40, FD&C Yellow No. 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide

Questions?

1-800-362-1683

NyQuil Cold & Flu

Drug Facts

Active ingredients (in each LiquiCap)

Acetaminophen 325 mg
Dextromethorphan HBr 15 mg
Doxylamine succinate 6.25 mg

Purpose

Pain reliever/Fever reducer
Cough suppressant
Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever
- runny nose & sneezing

Warnings

Liver warning:

This product contains acetaminophen.

Severe liver damage may occur if you take

- more than 8 LiquiCaps in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleep

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur

- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, & tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 8 LiquiCaps per 24 hrs

adults & children 12 yrs & over	2 LiquiCaps with water every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- store at no greater than 25C

Inactive ingredients

D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide

Questions?

1-800-362-1683

TAMPER EVIDENT:This package is safety-sealed & child resistant. Use only if blisters are intact. If difficult to open, use scissors.

Made in Canada

DIST. BY PROCTER & GAMBLE,

CINCINNATI OH 45202

PRINCIPAL DISPLAY PANEL - 24 Capsule Convenience Pack

VICKS®

DayQuil™

COLD & FLU

Acetaminophen, Phenylephrine HCl, Dextromethorphan HBr

Headache, Fever, Sore Throat, Minor Aches & Pains

Nasal Congestion, Sinus Pressure

Cough

Non-Drowsy

16 DayQuil LiquiCaps

NyQuil™

COLD & FLU

Acetaminophen, Doxylamine Succinate, Dextromethorphan HBr

Headache, Fever, Sore Throat, Minor Aches & Pains

Sneezing, Runny Nose

Cough

Nighttime Relief

8 NyQuil LiquiCaps: 24 Total LiquiCaps



<p>Drug Facts NyQuil™ Cold & Flu</p> <p>Active ingredients (in each LiqueCap)</p> <p>Acetaminophen 325 mg Pain reliever/fever reducer Dextromethorphan HBr 15 mg Cough suppressant Doxylamine succinate 6.25 mg Antihistamine</p> <p>Uses temporarily relieves common cold/flu symptoms:</p> <ul style="list-style-type: none"> cough due to minor throat & bronchial irritation sore throat headache minor aches & pains fever runny nose & sneezing <p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take</p> <ul style="list-style-type: none"> more than 8 LiqueCaps in 24 hours, which is the maximum daily amount for this product with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product <p>Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening • blisters • rash</p>	<p>Drug Facts DayQuil™ Cold & Flu</p> <p>Active ingredients (in each LiqueCap)</p> <p>Acetaminophen 325 mg Pain reliever/fever reducer Dextromethorphan HBr 10 mg Cough suppressant Phenylephrine HCl 5 mg Nasal decongestant</p> <p>Uses temporarily relieves common cold/flu symptoms:</p> <ul style="list-style-type: none"> nasal congestion cough due to minor throat & bronchial irritation sore throat headache minor aches & pains fever <p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take</p> <ul style="list-style-type: none"> more than 8 LiqueCaps in 24 hours, which is the maximum daily amount for this product with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product <p>Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening • blisters • rash</p>	<p>Drug Facts (continued)</p> <p>If a skin reaction occurs, stop use and seek medical help right away</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use</p> <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> liver disease glaucoma cough that occurs with too much phlegm (mucus) a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema trouble urinating due to enlarged prostate gland <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> taking sedatives or tranquilizers taking the blood thinning drug warfarin <p>When using this product</p> <ul style="list-style-type: none"> do not use more than directed excitability may occur, especially in children marked drowsiness may occur avoid alcoholic drinks be careful when driving a motor vehicle or operating machinery alcohol, sedatives, and tranquilizers may increase drowsiness <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> pain or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. 	<p>Drug Facts (continued)</p> <p>If a skin reaction occurs, stop use and seek medical help right away</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use</p> <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> liver disease heart disease high blood pressure thyroid disease diabetes trouble urinating due to enlarged prostate gland cough that occurs with too much phlegm (mucus) persistent or chronic cough such as occurs with smoking, asthma, or emphysema <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> taking the blood thinning drug warfarin <p>When using this product, do not use more than directed.</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> you get nervous, dizzy or sleepless pain, nasal congestion, or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>
<p>Drug Facts (continued)</p> <p>Directions</p> <ul style="list-style-type: none"> take only as directed do not exceed 8 LiqueCaps per 24 hrs adults & children 12 yrs & over 2 LiqueCaps with water every 6 hrs children 4 to under 12 yrs ask a doctor children under 4 yrs do not use <p>Other information</p> <ul style="list-style-type: none"> store at no greater than 25°C <p>Inactive ingredients D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide</p> <p>Questions? 1-800-362-1683</p>	<p>Drug Facts (continued)</p> <p>Directions</p> <ul style="list-style-type: none"> take only as directed do not exceed 8 LiqueCaps per 24 hrs adults & children 12 yrs & over 2 LiqueCaps with water every 4 hrs children 4 to under 12 yrs ask a doctor children under 4 yrs do not use <p>Other information</p> <ul style="list-style-type: none"> store at no greater than 25°C <p>Inactive ingredients FD&C Red No. 40, FD&C Yellow No. 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide</p> <p>Questions? 1-800-362-1683</p>	<p>Drug Facts (continued)</p> <p>Directions</p> <ul style="list-style-type: none"> take only as directed do not exceed 8 LiqueCaps per 24 hrs adults & children 12 yrs & over 2 LiqueCaps with water every 4 hrs children 4 to under 12 yrs ask a doctor children under 4 yrs do not use <p>Other information</p> <ul style="list-style-type: none"> store at no greater than 25°C <p>Inactive ingredients FD&C Red No. 40, FD&C Yellow No. 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide</p> <p>Questions? 1-800-362-1683</p>	<p>Drug Facts (continued)</p> <p>Directions</p> <ul style="list-style-type: none"> take only as directed do not exceed 8 LiqueCaps per 24 hrs adults & children 12 yrs & over 2 LiqueCaps with water every 4 hrs children 4 to under 12 yrs ask a doctor children under 4 yrs do not use <p>Other information</p> <ul style="list-style-type: none"> store at no greater than 25°C <p>Inactive ingredients FD&C Red No. 40, FD&C Yellow No. 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide</p> <p>Questions? 1-800-362-1683</p>

VICKS DAYQUIL AND VICKS NYQUIL COLD AND FLU

acetaminophen, phenylephrine hydrochloride, dextromethorphan hydrobromide, and doxylamine succinate kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69423-996
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423-996-12	1 in 1 PACKAGE; Type 1: Convenience Kit of Co-Package	01/01/2020	

2	NDC:69423-996-24	1 in 1 PACKAGE; Type 1: Convenience Kit of Co-Package	01/01/2020	
3	NDC:69423-996-60	1 in 1 PACKAGE; Type 1: Convenience Kit of Co-Package	01/01/2021	
4	NDC:69423-996-72	1 in 1 PACKAGE; Type 1: Convenience Kit of Co-Package	01/01/2020	
5	NDC:69423-996-48	1 in 1 PACKAGE; Type 1: Convenience Kit of Co-Package	01/01/2020	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BLISTER PACK	8
Part 2	1 BLISTER PACK	4

Part 1 of 2

VICKS DAYQUIL COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride capsule, liquid filled

Product Information

Item Code (Source)	NDC:69423-994
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POVIDONE (UNII: FZ989GH94E)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
SORBITOL (UNII: 506T60A25R)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	orange	Score	no score
Shape	OVAL	Size	21mm
Flavor		Imprint Code	DQuil
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/01/2020	

Part 2 of 2

VICKS NYQUIL COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate capsule, liquid filled

Product Information

Item Code (Source)	NDC:69423-995
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
SORBITOL (UNII: 506T60A25R)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
GELATIN (UNII: 2G86QN327L)	

FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
POVIDONE (UNII: FZ989GH94E)	
WATER (UNII: 059QF0KO0R)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
GLYCERIN (UNII: PDC6A3C0OX)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	

Product Characteristics

Color	green	Score	no score
Shape	OVAL	Size	21mm
Flavor		Imprint Code	NQUIL
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		4 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/01/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/01/2020	

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Establishment

Name	Address	ID/FEI	Business Operations
Catalent Ontario Limited		243944050	manufacture(69423-995, 69423-994)

Revised: 10/2023

The Procter & Gamble Manufacturing Company