

ASPIRIN- aspirin tablet, coated
WR Group, Inc.

Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

***nonsteroidal anti-inflammatory drug**

Purpose

Pain reliever

Uses

- for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**
- ask your doctor about other uses for safety coated 81 mg aspirin

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or last more than 3 days
- redness or swelling is present
- ringing in the ears or loss of hearing occurs
- new symptoms occur

These could be signs of a serious condition

If pregnant or breast-feeding,

ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use

Other information

- store between 15-30°C (59-86°F)

Inactive ingredients

D&C yellow #10, FD&C yellow #6, methacrylic acid copolymer, microcrystalline cellulose, pregelatinized starch (corn), silicon dioxide, sodium bicarbonate, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triethyl citrate

Principal Display Panel

Compare to the active ingredient in Bayer® Low Dose Aspirin®†

Aspirin

LOW DOSE

Enteric Safety Coated††

Pain reliever (**NSAID**)

Tablets

≠Talk to your doctor or other healthcare provider before using this product for your heart.

†This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Bayer® Low Dose Aspirin.

≠Coating helps protect against stomach upset.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

MANUFACTURED EXCLUSIVELY FOR:

KOLBE & SCHMITT

9160 E. BAHIA DR.

SCOTTSDALE, AZ 85260 USA

Product Label

NDC 69607-0867-6

Kolbe+Schmitt.

Aspirin 81mg

LOW DOSE
Enteric Safety Coated^{††}
Aspirin Regimen[†]

**Pain Reliever (NSAID)
60 Tablets**

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Drug Facts	Active ingredient (in each tablet) Aspirin 81 mg (NSAID)* *nonsteroidal anti-inflammatory drug	Purpose Pain reliever
Uses	<ul style="list-style-type: none"> for the temporary relief of minor aches and pains or as recommended by your doctor. <p>Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.</p> <ul style="list-style-type: none"> ask your doctor about other uses for safety coated 81 mg aspirin 	
Warnings	<p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p>	
Drug Facts (continued under label)		
<p>^{††}Coating helps protect against stomach upset</p> <p>[†]Talk to your doctor or other healthcare provider before using this product for your heart. Aspirin is not right for everyone.</p> <p>MANUFACTURED EXCLUSIVELY FOR: KOLBE & SCHMITT • 9160 E. BAHIA DR. SCOTTSDALE, AZ 85260 USA</p> <p>#14696 • L19</p> <p>PLD-0567A Lot No.: Exp. Date: LB007280</p>		

PEEL HERE →

Drug Facts (continued)
<p>Allergy alert: Aspirin may cause a severe allergic reaction, which may include:</p> <ul style="list-style-type: none"> hives facial swelling shock asthma (wheezing) <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:</p> <ul style="list-style-type: none"> are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.</p>
Directions
<ul style="list-style-type: none"> drink a full glass of water with each dose adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor children under 12 years: consult a doctor
Other information
<ul style="list-style-type: none"> store between 15-30°C (59-86°F)

KOLBE & SCHMITT Low Dose Aspirin

ASPIRIN			
aspirin tablet, coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69607-0867
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg
Inactive Ingredients			
	Ingredient Name		Strength
	D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)		
	FD&C YELLOW NO. 6 (UNII: H77VEI93A8)		
	METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13)		
	CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)		

STARCH, CORN (UNII: O8232NY3SJ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	E
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69607-0867-6	1 in 1 BOX	01/31/2020	
1		60 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/31/2020	

Labeler - WR Group, Inc. (089173699)

Revised: 4/2024

WR Group, Inc.