

**COLD SORE FEVER BLISTER TREATMENT- docosanol cream  
Meijer Distribution Inc.**

-----  
**Docosanol Cream, 10%**

***Active ingredient***

Docosanol 10%

***Purpose***

Cold sore/fever blister treatment

***Uses***

- treats cold sores/fever blisters on the face or lips
- shortens healing time and duration of symptoms:
  - tingling, pain, burning, and/or itching

**Warnings**

**Allergy Alert:** This product may cause a severe allergic reaction. Symptoms may include:

- hives
- facial swelling
- wheezing/difficulty breathing
- shock
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

**For external use only**

**Do not use**

- if you are allergic to any ingredient in this product

**When using this product**

- apply only to the affected areas
- do not use in or near the eyes
- avoid applying directly inside your mouth
- do not share this product with anyone. This may spread the infection.

**Stop use and ask a doctor if**

- your cold sore gets worse or the cold sore is not healed within 10 days

## **Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### ***Directions***

- **adults and children 12 years or over:**
  - wash hands before and after applying cream
  - apply to affected area on the face or lips at the first sign of cold sore/fever blister (tingle).
  - early treatment ensures the best results
  - rub in gently but completely
  - use 5 times a day until healed
- **children under 12 years:** ask a doctor

### ***Other information***

- store at 20° to 25°C (68° to 77°F)
- do not freeze

### ***Inactive ingredients***

benzyl alcohol, light mineral oil, propylene glycol, purified water, sucrose distearate, sucrose stearate

### **Questions or comments?**

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

### **Principal Display Panel**

**COMPARE TO ABREVA® ACTIVE INGREDIENT\*\***

Docosanol Cream, 10%

COLD SORE/FEVER BLISTER TREATMENT

Only FDA-approved medicine to shorten healing time\*

Cold sore treatment

Portable convenient pump

NET WT 2 g (0.07 oz)

\*Docosanol Cream, 10% contains the only OTC medicine approved by the FDA to shorten healing time and duration of symptoms.

\*\*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Abreva®.

**RETAIN THIS INSERT FOR FULL PRODUCT USES, DIRECTIONS AND WARNINGS.  
 DIST. BY MEIJER DISTRIBUTION, INC.  
 GRAND RAPIDS, IN 49544**

**Package Label**



**Thank you for purchasing Docosanol Cream, 10%.** We hope that since Docosanol Cream has been demonstrated to shorten the healing time of cold sores, it helps you look and feel better faster. Please follow insert directions for proper use and refer to the information below.

**You are not alone**

An estimated one-in-five Americans is affected by cold sores each year. Most people who experience them get two to three cold sore outbreaks each year. Just before and during an outbreak, the cold sore infection can be easily passed on to others. It is especially important to be aware of early warning signs that a cold sore is coming, and to be sure to avoid close physical contact when you have one.

**Getting to the Root of the Problem**

A blister, or cluster of blisters around your mouth, is caused by the cold sore virus. After the cold sore has healed, the virus stays dormant in your body but can be reactivated by everyday triggers like stress, sunlight, extreme hot or cold weather, colds and flu, lack of sleep, or menstruation.

**The Sooner, the Better**

A typical cold sore outbreak can last as long as seven to ten days. The sooner you begin using Docosanol Cream, 10%, the more effective the medicine will be. So for best results, use Docosanol Cream, 10% at the very start of the cold sore - when you feel the first sign of the tingle, redness, bump or itch.

Docosanol Cream, 10% is different from other over-the-counter cold sore products. Only Docosanol Cream, 10% contains 10% docosanol, which helps to shorten the healing time of cold sores and the duration of symptoms including tingling, pain, itching and burning.

**How to use your Docosanol Cream, 10% Pump**

- For adults and Children 12 years and over
- Docosanol Cream, 10% should be used five times a day until the cold sore is healed. Apply it directly to the cold sore at the first sign of a tingle, redness, bump or itch for best results.
- If you wish, you can apply cosmetics such as lipstick over Docosanol Cream, 10%. However, use a separate applicator, like a cotton swab, to apply cosmetics over an unhealed cold sore to avoid spreading the infection. For best results, remove any cosmetics prior to applying Docosanol Cream, 10%.

**Press Pump completely about 6-8 times to get first dose.**



**STEP 1:** Wash your hands before and after treating the cold sore.

**STEP 2:** Remove cap and press pump firmly. When first used, expect to press pump completely about 6-8 times before obtaining first dose.

**STEP 3:** Rub in gently but completely 5 times a day until healed for maximum of 10 days.

**STEP 4:** Place cap back on pump after each use.

**For more information:** Call us at 1-877-753-3935

7 08820 29480 6  
 PCD-4724 PFD 490820 F000694

www.meijer.com  
 DIST. BY MEIJER DISTRIBUTION, INC.  
 GRAND RAPIDS, MI 49544

RETAIN THIS INSERT FOR FULL PRODUCT USES, DIRECTIONS AND WARNINGS.

© Docosanol Cream, 10% contains the only OTC medicine approved by the FDA to shorten healing time and duration of symptoms.

\*\*\*This product is not manufactured or distributed by GSK Consumer Healthcare, owner of the registered trademark Abreva®.

QUESTIONS OR COMMENTS?  
 Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

**Stop use and ask a doctor if**  
 ■ your cold sore gets worse or the cold sore is not healed within 10 days  
 ■ you do not use or near the eyes  
 ■ you do not share this product with anyone. This may spread the infection.  
 ■ you do not use in or near the eyes  
 ■ you avoid applying directly inside your mouth

**When using this product**  
 ■ apply only to the affected areas  
 ■ if you are allergic to any ingredient in this product

**Do not use**  
 ■ if you are allergic to any ingredient in this product

**For external use only**  
 If an allergic reaction occurs, stop use and seek medical help right away.  
 Symptoms may include:  
 ■ hives ■ rash ■ itching, pain, burning, and/or itching  
 ■ facial swelling ■ severe allergic reaction

**Warnings**  
 ■ Allergy alert: This product may cause a severe allergic reaction.  
 ■ Avoid eye contact. If eye contact occurs, wash eyes with water. If irritation persists, seek medical help.  
 ■ Avoid contact with others. This product may be contagious.

**Uses**  
 ■ treats cold sores/fever blisters on the face or lips  
 ■ shortens healing time and duration of symptoms:  
 ■ tingling, pain, burning, and/or itching

**Directions**  
 ■ adults and children 12 years or over:  
 ■ apply to affected area on the face or lips at the first sign of cold sore/fever blister (tingle)  
 ■ wash hands before and after applying cream  
 ■ apply to affected area on the face or lips at the first sign of cold sore/fever blister (tingle)  
 ■ use 5 times a day until healed  
 ■ do not freeze  
 ■ store at 20°-25°C (68°-77°F)

**Other information**  
 ■ do not freeze  
 ■ store at 20°-25°C (68°-77°F)

**Inactive ingredients**  
 benzyl alcohol, light mineral oil, propylene glycol, purified water, sucrose disaccharide, sucrose stearate

**Keep out of reach of children.**  
 If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Drug Facts (continued)**

**MEIJER Cold Sore Fever Blister Treatment**

**COLD SORE FEVER BLISTER TREATMENT**

docosanol cream

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:79481-0065
<b>Route of Administration</b>	TOPICAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>DOCOSANOL</b> (UNII: 9G1OE216XY) (DOCOSANOL - UNII:9G1OE216XY)	DOCOSANOL	100 mg in 1 g

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	
<b>LIGHT MINERAL OIL</b> (UNII: N6K5787QVP)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SUCROSE DISTEARATE</b> (UNII: 33X4X4B90S)	
<b>SUCROSE STEARATE</b> (UNII: 274KW0050M)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:79481-0065-7	2 g in 1 BOTTLE, PUMP; Type 0: Not a Combination Product	07/07/2023	

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA212385	07/07/2023	

**Labeler** - Meijer Distribution Inc. (006959555)