

# **FIRDAPSE- amifampridine phosphate tablet**

## **Catalyst Pharmaceuticals, Inc.**

-----

### **HIGHLIGHTS OF PRESCRIBING INFORMATION**

**These highlights do not include all the information needed to use FIRDAPSE® safely and effectively. See full prescribing information for FIRDAPSE®.**

**FIRDAPSE® (amifampridine) tablets, for oral use**  
**Initial U.S. Approval: 2018**

### **INDICATIONS AND USAGE**

FIRDAPSE is a potassium channel blocker indicated for the treatment of Lambert-Eaton myasthenic syndrome (LEMS) in adults. (1)

### **DOSAGE AND ADMINISTRATION**

- The recommended starting dosage is 15 mg to 30 mg daily taken orally in divided doses (3 to 4 times daily). (2.1)
- Starting dosage is 15 mg daily for patients with renal impairment, hepatic impairment, and in known N-acetyltransferase 2 (NAT2) poor metabolizers (2.2, 2.3, 2.4)
- Dosage can be increased by 5 mg daily every 3 to 4 days. (2.1)
- Dosage is not to exceed a maximum of 80 mg daily. (2.1)
- The maximum single dose is 20 mg. (2.1)

### **DOSAGE FORMS AND STRENGTHS**

- Tablets: 10 mg, functionally scored. (3)

### **CONTRAINDICATIONS**

FIRDAPSE is contraindicated in patients with:

- A history of seizures (4)
- Hypersensitivity to amifampridine or another aminopyridine (4)

### **WARNINGS AND PRECAUTIONS**

- Seizures: FIRDAPSE can cause seizures. Consider discontinuation or dose-reduction of FIRDAPSE in patients who have a seizure while on treatment. (5.1)
- Hypersensitivity reactions: If a hypersensitivity reaction such as anaphylaxis occurs, FIRDAPSE should be discontinued and appropriate therapy initiated. (5.2)

### **ADVERSE REACTIONS**

The most common (> 10%) adverse reactions are: paresthesia, upper respiratory tract infection, abdominal pain, nausea, diarrhea, headache, elevated liver enzymes, back pain, hypertension, and muscle spasms. (6)

**To report SUSPECTED ADVERSE REACTIONS, contact Catalyst Pharmaceuticals at 1-844-347-3277 (1-844-FIRDAPSE) or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

### **DRUG INTERACTIONS**

- Drugs that lower seizure threshold: The concomitant use of FIRDAPSE and drugs that lower seizure threshold may lead to an increased risk of seizures. (7.1)
- Drugs with cholinergic effects: The concomitant use of FIRDAPSE and drugs with cholinergic effects (e.g., direct or indirect cholinesterase inhibitors) may increase the cholinergic effects of FIRDAPSE and of those drugs and increase the risk of adverse reactions. (7.2)

### **USE IN SPECIFIC POPULATIONS**

- Pregnancy: Based on animal data, may cause fetal harm (8.1)

**See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.**

**Revised: 3/2021**

---

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

### **1 INDICATIONS AND USAGE**

### **2 DOSAGE AND ADMINISTRATION**

- 2.1 Dosage Information
- 2.2 Patients with Renal Impairment
- 2.3 Patients with Hepatic Impairment
- 2.4 Known N-acetyltransferase 2 (NAT2) Poor Metabolizers
- 2.5 Administration Instructions

### **3 DOSAGE FORMS AND STRENGTHS**

### **4 CONTRAINDICATIONS**

### **5 WARNINGS AND PRECAUTIONS**

- 5.1 Seizures
- 5.2 Hypersensitivity

### **6 ADVERSE REACTIONS**

- 6.1 Clinical Trials Experience

### **7 DRUG INTERACTIONS**

- 7.1 Drugs that Lower Seizure Threshold
- 7.2 Drugs with Cholinergic Effects

### **8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment
- 8.8 NAT2 Poor Metabolizers

### **10 OVERDOSAGE**

### **11 DESCRIPTION**

### **12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics
- 12.5 Pharmacogenomics

### **13 NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

### **14 CLINICAL STUDIES**

### **16 HOW SUPPLIED/ STORAGE AND HANDLING**

- 16.1 How Supplied
- 16.2 Storage and Handling

### **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

## 1 INDICATIONS AND USAGE

FIRDAPSE<sup>®</sup> is indicated for the treatment of Lambert-Eaton myasthenic syndrome (LEMS) in adults.

## 2 DOSAGE AND ADMINISTRATION

### 2.1 Dosage Information

- The recommended starting dosage of FIRDAPSE is 15 mg to 30 mg daily, taken orally in divided doses (3 to 4 times daily).
- The dosage can be increased by 5 mg daily every 3 or 4 days.
- The maximum recommended total daily dosage is 80 mg.
- The maximum single dose is 20 mg.
- If a dose is missed, patients should not take double or extra doses.

### 2.2 Patients with Renal Impairment

The recommended starting dosage of FIRDAPSE in patients with renal impairment (creatinine clearance 15 to 90 mL/min) is 15 mg daily, taken orally in 3 divided doses. No dosage recommendation for FIRDAPSE can be made for patients with end-stage renal disease [see *Use in Specific Populations (8.6)* and *Clinical Pharmacology (12.3, 12.5)*].

### 2.3 Patients with Hepatic Impairment

The recommended starting dosage of FIRDAPSE in patients with any degree of hepatic impairment is 15 mg daily, taken orally in 3 divided doses [see *Use in Specific Populations (8.7)* and *Clinical Pharmacology (12.3, 12.5)*].

### 2.4 Known N-acetyltransferase 2 (NAT2) Poor Metabolizers

The recommended starting dosage of FIRDAPSE in known N-acetyltransferase 2 (NAT2) poor metabolizers is 15 mg daily, taken orally in 3 divided doses [see *Use in Specific Populations (8.8)* and *Clinical Pharmacology (12.3, 12.5)*].

### 2.5 Administration Instructions

FIRDAPSE can be taken without regard to food.

## 3 DOSAGE FORMS AND STRENGTHS

FIRDAPSE tablets contain 10 mg amifampridine and are white to off-white, round, and functionally scored. Each tablet is debossed on the non-scored side with "CATALYST" and on the scored side with "211" above the score and "10" below the score.

## 4 CONTRAINDICATIONS

FIRDAPSE is contraindicated in patients with:

- A history of seizures [see *Warnings and Precautions (5.1)*]
- Hypersensitivity to amifampridine phosphate or another aminopyridine [see *Warnings and Precautions (5.2)*]

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Seizures**

FIRDAPSE can cause seizures. Seizures have been observed in patients without a history of seizures taking FIRDAPSE at the recommended doses, at various times after initiation of treatment, at an incidence of approximately 2%. Many of the patients were taking medications or had comorbid medical conditions that may have lowered the seizure threshold [see *Drug Interactions (7.1)*]. Seizures may be dose-dependent. Consider discontinuation or dose-reduction of FIRDAPSE in patients who have a seizure while on treatment. FIRDAPSE is contraindicated in patients with a history of seizures.

### **5.2 Hypersensitivity**

In clinical trials, hypersensitivity reactions and anaphylaxis associated with FIRDAPSE administration have not been reported. Anaphylaxis has been reported in patients taking another aminopyridine; therefore, it may occur with FIRDAPSE. If anaphylaxis occurs, administration of FIRDAPSE should be discontinued and appropriate therapy initiated.

## **6 ADVERSE REACTIONS**

The following serious adverse reactions are described elsewhere in the labeling:

- Seizures [see *Warnings and Precautions (5.1)*]
- Hypersensitivity [see *Warnings and Precautions (5.2)*]

### **6.1 Clinical Trials Experience**

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

In controlled and uncontrolled trials (Study 1 and 2) in patients with LEMS, 63 patients were treated with FIRDAPSE, including 40 patients treated for more than 6 months, and 39 patients treated for more than 12 months. In an expanded access program, 139 patients with LEMS were treated with FIRDAPSE, including 102 patients treated for more than 6 months, 77 patients treated for more than 12 months, and 53 patients treated for more than 18 months.

Study 1 was a double-blind, placebo-controlled, randomized discontinuation study in adults with LEMS. Following an initial open-label run-in phase (up to 90 days), patients were randomized to either continue FIRDAPSE treatment or transition to placebo, for a 14-day double-blind phase. Following final assessments, patients were allowed to resume FIRDAPSE treatment for up to 2 years (open-label long-term safety phase of the study).

During the open-label run-in phase of Study 1, 53 patients received FIRDAPSE for an average of 81 days at a mean daily dosage of 50.5 mg/day. The mean patient age was 52.1 years and 66% were female. There were 42 patients who had no prior exposure to FIRDAPSE at the initiation of this study. Table 1 shows adverse reactions with an incidence of 5% or greater occurring in the 42 LEMS patients newly initiated on treatment with FIRDAPSE during the run-in phase of the study.

**Table 1. Adverse Reactions in  $\geq 5\%$  of LEMS Patients Newly Treated with FIRDAPSE in Study 1**

Adverse Reaction	FIRDAPSE N=42 %
Paresthesia*	62
Upper respiratory tract infection	33
Abdominal pain	14
Nausea	14
Diarrhea	14
Headache	14
Elevated liver enzymes†	14
Back pain	14
Hypertension	12
Muscle spasms	12
Dizziness	10
Asthenia	10
Muscular weakness	10
Pain in extremity	10
Cataract	10
Constipation	7
Bronchitis	7
Fall	7
Lymphadenopathy	7

\* Includes paresthesia, oral paresthesia, oral hypoesthesia

† Includes elevated alanine aminotransferase (ALT), aspartate aminotransferase (AST), lactate dehydrogenase (LDH), and gamma-glutamyl transferase (GGT)

### Other Adverse Reactions

In the overall population treated in Study 1 (n=53), including the double-blind phase and the 2-year open-label long-term safety phase, additional adverse reactions occurring in at least 5% of the patients included: dyspnea, urinary tract infection, gastroesophageal reflux, insomnia, peripheral edema, pyrexia, viral infection, blood creatine phosphokinase increase, depression, erythema, hypercholesterolemia, and influenza. These patients received a mean daily dosage of 66 mg of FIRDAPSE.

## **7 DRUG INTERACTIONS**

### **7.1 Drugs that Lower Seizure Threshold**

The concomitant use of FIRDAPSE and drugs that lower seizure threshold may lead to an increased risk of seizures [see *Warnings and Precautions (5.1)*]. The decision to administer FIRDAPSE concomitantly with drugs that lower the seizure threshold should be carefully considered in light of the severity of the associated risks.

## 7.2 Drugs with Cholinergic Effects

The concomitant use of FIRDAPSE and drugs with cholinergic effects (e.g., direct or indirect cholinesterase inhibitors) may increase the cholinergic effects of FIRDAPSE and of those drugs and increase the risk of adverse reactions.

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Pregnancy Exposure Registry

There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to FIRDAPSE during pregnancy. Physicians are encouraged to enroll pregnant patients, or pregnant women may register themselves in the registry by calling 855-212-5856 (toll-free), using the Fax number 877-867-1874 (toll-free), by contacting the Pregnancy Coordinating Center at [firdapsepregnancyregistry@ubc.com](mailto:firdapsepregnancyregistry@ubc.com), or by visiting the study website [www.firdapsepregnancystudy.com](http://www.firdapsepregnancystudy.com).

#### Risk Summary

There are no data on the developmental risk associated with the use of FIRDAPSE in pregnant women. In animal studies, administration of amifampridine phosphate to rats during pregnancy and lactation resulted in developmental toxicity (increase in stillbirths and pup deaths, reduced pup weight, and delayed sexual development) at doses associated with maternal plasma drug levels lower than therapeutic drug levels (see Animal Data). In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively. The background risk of major birth defects and miscarriage for the indicated population is unknown.

#### Data

##### *Animal Data*

Oral administration of amifampridine phosphate (0, 7.5, 22.5, or 75 mg/kg/day) to female rats prior to and during mating and continuing throughout organogenesis produced no adverse effects on embryofetal development. Plasma amifampridine exposure (AUC) at the highest dose tested is approximately 7 times that in humans at the maximum recommended human dose (MRHD) of 80 mg amifampridine/day. Oral administration of amifampridine phosphate (0, 9, 30, or 57 mg/kg/day) to pregnant rabbits throughout organogenesis produced no adverse effects on embryofetal development. The highest dose tested is approximately 7 times the MRHD (80 mg/day amifampridine) on a body surface area (mg/m<sup>2</sup>) basis.

Oral administration of amifampridine phosphate (0, 7.5, 22.5, or 75 mg/kg/day) to female rats throughout pregnancy and lactation resulted in an increase in stillbirths and pup deaths, reduced pup weight, and delayed sexual development in female pups at the mid and high doses tested. The no-effect dose (7.5 mg/kg/day amifampridine phosphate) for adverse developmental effects is associated with a plasma amifampridine exposure (AUC) less than that in humans at the MRHD.

### 8.2 Lactation

## Risk Summary

There are no data on the presence of FIRDAPSE in human milk, the effects on the breastfed infant, or the effects on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for FIRDAPSE and any potential adverse effects on the breastfed infant from FIRDAPSE or from the underlying maternal condition.

In lactating rat, amifampridine was excreted in milk and reached levels similar to those in maternal plasma.

### **8.4 Pediatric Use**

Safety and effectiveness in pediatric patients have not been established.

### **8.5 Geriatric Use**

Clinical studies of FIRDAPSE did not include sufficient numbers of subjects aged 65 and over (19 of 63 patients in Studies 1 and 2) to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy [*see Dosage and Administration (2.2, 2.3) and Drug Interactions (7.2, 7.3)*].

### **8.6 Renal Impairment**

Renal clearance is an elimination pathway for amifampridine and the inactive metabolite, 3-N-acetyl amifampridine, and exposure of amifampridine is higher in subjects with renal impairment [*see Clinical Pharmacology (12.3)*]. Therefore, in patients with renal impairment, FIRDAPSE should be initiated at the lowest recommended starting dosage (15 mg/day), and patients should be closely monitored for adverse reactions [*see Dosage and Administration (2.2)*]. Consider dosage modification or discontinuation of FIRDAPSE for patients with renal impairment as needed based on clinical effect and tolerability. The safety, efficacy, and pharmacokinetics of amifampridine have not been studied in patients with end-stage renal disease (CL<sub>cr</sub> <15 mL/min or patients requiring dialysis). No dosage recommendation for FIRDAPSE can be made for patients with end-stage renal disease.

### **8.7 Hepatic Impairment**

The effects of FIRDAPSE have not been studied in patients with hepatic impairment. FIRDAPSE is extensively metabolized by N-acetyltransferase 2 (NAT2) and hepatic impairment may cause an increase in exposure. Therefore, initiate FIRDAPSE in patients with any degree of hepatic impairment at the lowest recommended starting dosage (15 mg/day) and monitor for adverse reactions [*see Dosage and Administration (2.3)*]. Consider dosage modification or discontinuation of FIRDAPSE for patients with hepatic impairment as needed based on clinical effect and tolerability.

### **8.8 NAT2 Poor Metabolizers**

Exposure of FIRDAPSE is increased in patients who are N-acetyltransferase 2 (NAT2)

poor metabolizers [see *Clinical Pharmacology (12.5)*]. Therefore, initiate FIRDAPSE in patients who are known NAT2 poor metabolizers at the lowest recommended starting dosage (15 mg/day) and monitor for adverse reactions [see *Dosage and Administration (2.4)*]. Consider dosage modification of FIRDAPSE for patients who are known NAT2 poor metabolizers as needed based on clinical effect and tolerability.

## 10 OVERDOSAGE


Overdose with FIRDAPSE was not reported during clinical studies.

In a case report, a 65-year-old patient with LEMS inadvertently received a total daily amifampridine dose of 360 mg/day (more than 4 times the maximum recommended total daily dose) and was hospitalized for general weakness, paresthesia, nausea, vomiting, and palpitations. The patient developed convulsions and paroxysmal supraventricular tachycardia, and four days after admission, experienced cardiac arrest. The patient was resuscitated and ultimately recovered following withdrawal of amifampridine.

Patients with suspected overdose with FIRDAPSE should be monitored for signs or symptoms of exaggerated FIRDAPSE adverse reactions or effects, and appropriate symptomatic treatment instituted immediately.

## 11 DESCRIPTION

The active ingredient of FIRDAPSE is amifampridine phosphate, which is a voltage-gated potassium channel blocker. Amifampridine phosphate is described chemically as 3,4-diaminopyridine phosphate with a molecular weight of 207.1 and a molecular formula of  $C_5H_7N_3 \cdot H_3PO_4$ . The structural formula is:



Amifampridine phosphate is a white, crystalline powder that is freely soluble in water, and slightly soluble in solvents ethanol, methanol and acetic acid. A 1% aqueous solution of amifampridine phosphate has a pH of 4.4 at ambient conditions.

Each FIRDAPSE tablet contains 10 mg amifampridine (equivalent to 18.98 mg amifampridine phosphate). The tablet formulation includes the following inactive ingredients: calcium stearate, colloidal silicon dioxide, and microcrystalline cellulose.

FIRDAPSE tablets are intended for oral administration only.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action



The mechanism by which amifampridine exerts its therapeutic effect in LEMS patients has not been fully elucidated. Amifampridine is a broad spectrum potassium channel blocker.

## 12.2 Pharmacodynamics

The effect of FIRDAPSE on QTc interval prolongation was studied in a double blind, randomized, placebo and positive controlled study in 52 healthy individuals who are slow acetylators. At an exposure 2-fold the expected maximum therapeutic exposure of amifampridine, FIRDAPSE did not prolong QTc to any clinically relevant extent.

## 12.3 Pharmacokinetics

The pharmacokinetics of amifampridine are similar between healthy individuals and LEMS patients. Following single and multiple doses, AUC,  $C_{max}$  and  $C_{min}$  were highly variable between individuals. FIRDAPSE exposure increased proportionally with dose across the range of 20 mg to 80 mg single oral doses.

### Absorption

Amifampridine peak plasma concentration is reached 20 minutes to 1 hour after administration. Food does not have a clinically significant effect on the exposure of amifampridine.

### Elimination

Amifampridine is eliminated primarily through metabolism to 3-N-acetyl-amifampridine and to a smaller extent through the kidneys. The terminal half-life ranges from 1.8 to 2.5 hours in healthy subjects.

### Metabolism

Amifampridine is extensively metabolized by N-acetyltransferase 2 (NAT2) to 3-N-acetyl-amifampridine, which is considered an inactive metabolite.

### Excretion

Following administration of FIRDAPSE to healthy subjects, 93% to 100% of the administered dose was eliminated in the urine as amifampridine or 3-N-acetyl amifampridine over 24 hours.

### Specific Populations

#### Patients with Renal Impairment

Pharmacokinetic data are available from a study of 24 otherwise healthy subjects with impaired renal function who received a single 10-mg dose of FIRDAPSE. The exposure of amifampridine (measured as AUC) was 2- to 3-fold higher in subjects with moderate (CLcr 30-59 mL/min) or severe (CLcr 15-29 mL/min) renal impairment than in subjects with normal renal function (CLcr greater than or equal to 90 mL/min). Compared with subjects with normal renal function, subjects with mild renal impairment (CLcr 60-89 mL/min) had a 36% increase in exposure. Therefore, FIRDAPSE should be initiated at the lowest recommended starting dosage (15 mg/day) in patients with renal impairment, and such patients should be closely monitored for adverse reactions [see *Dosage and Administration (2.2) and Use in Specific Populations (8.6)*].  $C_{max}$  was marginally affected by renal impairment.

## 12.5 Pharmacogenomics

Genetic variants in the N-acetyltransferase gene 2 (NAT2) affect the rate and extent of FIRDAPSE metabolism. Poor metabolizers, also referred to as "slow acetylators" (i.e., carriers of two reduced function alleles), have 3.5- to 4.5-fold higher  $C_{max}$ , and 5.6- to 9- fold higher AUC than normal metabolizers, also referred to as "fast/rapid acetylators" (i.e., carriers of two normal function alleles). Therefore, FIRDAPSE should be initiated at the lowest recommended starting dosage (15 mg/day) in known NAT2 poor metabolizers, and such patients should be closely monitored for adverse reactions [see *Dosage and Administration (2.4) and Use in Specific Populations (8.8)*]. In the general population, the NAT2 poor metabolizer phenotype prevalence is 40–60% in the White and African American populations, and in 10–30% in Asian ethnic populations (individuals of Japanese, Chinese, or Korean descent).

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

#### Carcinogenicity

In a 104-week carcinogenicity study, oral administration of amifampridine phosphate (0, 15, 48, or 105 mg/kg/day) resulted in an increase in uterine tumors (endometrial carcinoma and combined endometrial adenoma/endometrial carcinoma/squamous cell carcinoma) at the mid and high doses tested. The low dose, not associated with an increase in tumors, is similar to the maximum recommended human dose (80 mg/day amifampridine) on a body surface area (mg/m<sup>2</sup> basis).

#### Mutagenesis

Amifampridine phosphate was negative in the *in vitro* bacterial reverse mutation and *in vivo* rat micronucleus assays. Amifampridine phosphate was positive for clastogenicity in the *in vitro* mouse lymphoma *tk* assay in the absence of metabolic activation.

#### Impairment of Fertility

Oral administration of amifampridine phosphate (0, 7.5, 22.5, or 75 mg/kg/day) to male and female rats prior to and during mating, and continuing in females throughout organogenesis, produced no adverse effects on fertility. Plasma amifampridine exposure (AUC) at the highest dose tested is approximately 7 times that in humans at the maximum recommended human dose (MRHD) of 80 mg amifampridine/day.

## 14 CLINICAL STUDIES

The efficacy of FIRDAPSE for the treatment of LEMS was demonstrated in two randomized, double-blind, placebo-controlled discontinuation studies. A total of 64 adults (age 21 to 88 years) with LEMS were enrolled (Study 1 and Study 2). The studies enrolled patients with a confirmed diagnosis of LEMS based on either neurophysiology studies or a positive anti-P/Q type voltage-gated calcium channel antibody test. Patients were required to be on an adequate and stable dosage (30 to 80 mg daily) of amifampridine phosphate prior to entering the randomized discontinuation phases of both studies.

The two co-primary measures of efficacy in both studies were the change from baseline to the end of the discontinuation period in the Quantitative Myasthenia Gravis (QMG) score and in the Subject Global Impression (SGI) score.

The QMG is a 13-item physician-rated categorical scale assessing muscle weakness. Each item is assessed on a 4 -point scale, where a score of 0 represents no weakness, and a score of 3 represents severe weakness (total score 0-39). Higher scores represent greater impairment.

The SGI is a 7-point scale on which patients rated their global impression of the effects of the study treatment on their physical well-being. Lower scores on the SGI represent lower perceived benefit with the study treatment.

A key secondary efficacy endpoint was the clinical global impression improvement (CGI-I) score, a 7-point scale on which the treating physician rated the global impression of change in clinical symptoms. A higher CGI-I score indicates a perceived worsening of clinical symptoms.

### Study 1 (NCT01377922)

After an initial open-label run-in phase, 38 patients were randomized in a double-blind fashion to either continue treatment with FIRDAPSE (n=16) or to a downward titration to placebo (n=22) over 7 days. Following the downward titration period, patients remained on blinded FIRDAPSE or placebo for 7 more days. Efficacy was assessed at Day 14 of the double-blind period. Patients were allowed to use stable dosages of peripherally acting cholinesterase inhibitors or oral immunosuppressants. Twenty-six percent of patients randomized to FIRDAPSE were receiving cholinesterase inhibitors, versus 36% in the placebo group, and 28% of patients randomized to FIRDAPSE were receiving oral immunosuppressant therapies, versus 34% in the placebo group.

Patients had a median age of 54 years (range: 21 to 88 years), 61% were female, and 90% were White. Eighty-four percent of patients had a diagnosis of autoimmune LEMS, and 16% of patients had a diagnosis of paraneoplastic LEMS.

During the double-blind period (from Baseline to Day 14), the QMG scores tended to worsen in both treatment groups, but there was significantly greater worsening in the placebo group than in the FIRDAPSE group ( $p=0.045$ ). Similarly, the SGI score tended to worsen in both treatment groups during the double-blind period, but there was significantly greater worsening in the placebo group than in the FIRDAPSE group ( $p=0.003$ ), as summarized in Table 2. These results indicate that in Study 1, patients randomized to placebo had a significantly greater worsening of muscle weakness and of global impression of the effects of the study treatment on their physical well-being, compared to patients who continued FIRDAPSE in the double-blind period.

**Table 2. Change from Baseline to Day 14 in QMG Score and SGI Score in Study 1**

Assessment	FIRDAPSE (n=16)	Placebo (n=21)
<b>Primary Endpoints</b>		
<b>QMG Score*</b>		
Baseline (mean)	6.4	5.6
Change from Baseline (Least Square Mean)	0.4	2.2

FIRDAPSE-placebo Treatment Difference (Least Square Mean (95% CI))	-1.7 (-3.4, -0.0)	
p-value <sup>†</sup>	0.045	
<b>SGI Score<sup>‡</sup></b>		
Baseline (mean)	5.6	5.9
Change from Baseline (Least Square Mean))	-0.8	-2.6
FIRDAPSE-placebo Treatment Difference, (Least Square Mean (95% CI))	1.8 (0.7, 3.0)	
p-value <sup>†</sup>	0.003	

\* QMG Score range 0 (no impairment) to 39 (worst impairment)

† Pairwise contrast at Day 14 from mixed-effects model with repeated measures.

‡ SGI Score range 0 (least perceived benefit) to 7 (most perceived benefit)

The CGI-I score was significantly greater for patients randomized to placebo than for patients who continued treatment with FIRDAPSE, with a mean difference between FIRDAPSE and placebo of -1.1 ( $p=0.02$ ), indicating that clinicians perceived a greater worsening of clinical symptoms in patients who were randomized to placebo and discontinued from FIRDAPSE treatment, compared to patients who continued FIRDAPSE in the double-blind period.

### Study 2 (NCT02970162)

Patients on stable treatment with FIRDAPSE were randomized 1:1 in a double-blind fashion to either continue treatment with FIRDAPSE ( $n=13$ ) or change to placebo ( $n=13$ ) for 4 days. Efficacy was assessed at the end of the 4-day double-blind discontinuation period. Patients were allowed to use stable doses of peripherally acting cholinesterase inhibitors or corticosteroids. Sixty-one percent of patients randomized to FIRDAPSE were receiving cholinesterase inhibitors, versus 54% of patients randomized to placebo. Corticosteroid use was similar between FIRDAPSE and placebo (8%). Patients with recent use of immunomodulatory therapies (e.g., azathioprine, mycophenolate, cyclosporine), rituximab, intravenous immunoglobulin G, and plasmapheresis were excluded from the study. Patients had a median age of 55.5 years (range: 31 to 75 years), 62% were female, and 88% were White.

From Baseline to Day 4, there was significantly greater worsening in the QMG score in the placebo group than in the FIRDAPSE group ( $p=0.0004$ ), and also significantly greater worsening in the SGI score in the placebo group than in the FIRDAPSE group ( $p=0.0003$ ), as summarized in Table 3. These results indicate that in Study 2, patients randomized to placebo had a significantly greater worsening of muscle weakness and of global impression of the effects of the study treatment on their physical well-being, compared to patients who continued FIRDAPSE in the double-blind period.

**Table 3. Change from Baseline to Day 4 in QMG Scores and SGI Scores in Study 2**

<b>Assessment</b>	<b>FIRDAPSE (n=13)</b>	<b>Placebo (n=13)</b>
<b>QMG Scores*</b>		
Baseline, Mean	7.8	8.5
Change from Baseline, Least Square Mean <sup>†</sup>	0.00	6.54
FIRDAPSE-placebo Treatment Difference, Least Square Mean (95% CI)	-6.54 (-9.78, -3.29)	
p-value <sup>‡</sup>	0.0004	
<b>SGI Scores<sup>§</sup></b>		
Baseline, Mean	6.1	5.8
Change from Baseline, Least Square Mean <sup>†</sup>	-0.64	-3.59
FIRDAPSE-placebo Treatment Difference, Least Square Mean (95% CI)	2.95 (1.53, 4.38)	
p-value <sup>‡</sup>	0.0003	

\* QMG Score range 0 (no impairment) to 39 (worst impairment)

† Change from baseline for QMG total score was modeled as the response, with fixed effects terms for treatment and QMG at Baseline

‡ p-value based on the Wilcoxon Rank Sum Test for treatment differences.

§ SGI Score range 0 (least perceived benefit) to 7 (most perceived benefit)

The clinical global impression improvement (CGI-I) score was significantly greater for patients randomized to placebo than for patients who continued treatment with FIRDAPSE, with a mean difference between FIRDAPSE and placebo of -2.7 ( $p=0.002$ ), indicating that clinicians perceived a greater worsening of clinical symptoms in patients who were randomized to placebo and discontinued from FIRDAPSE treatment, compared to patients who continued FIRDAPSE in the double-blind period.

## **16 HOW SUPPLIED/ STORAGE AND HANDLING**

### **16.1 How Supplied**

FIRDAPSE 10 mg tablets are white to off white, round, and functionally scored. Each tablet is debossed on the non-scored side with "CATALYST" and on the scored side with "211" above the score and "10" below the score. Tablets can be divided in half at the score. FIRDAPSE is supplied as follows:

#### Child Resistant Blister Pack

- blister pack containing 10 tablets
- carton containing 12 blister packs (120 tablets total) NDC 69616-211-04  
NDC 69616-211-06

## Bottles

- 60 tablets
- 240 tablets

NDC 69616-211-08

NDC 69616-211-03

## **16.2 Storage and Handling**

Store FIRDAPSE tablets at 20°C to 25°C (68°F to 77°F) with excursions permitted from 15°C to 30°C (59°F to 86°F) [see USP controlled room temperature].

## **17 PATIENT COUNSELING INFORMATION**

Advise the patient and/or caregiver to read the FDA-approved patient labeling (Medication Guide).

### Risk of Seizures

Inform patients that FIRDAPSE can cause seizures, and to notify their healthcare provider if they experience a seizure [see *Warnings and Precautions (5.1)*].

### Hypersensitivity

Instruct patients to inform their healthcare provider if they have signs or symptoms of hypersensitivity, and to seek emergency help if symptoms of anaphylaxis occur [see *Warnings and Precautions (5.2)*].

### FIRDAPSE Dosing

Instruct patients to take FIRDAPSE exactly as prescribed. Patients should carefully follow the dose escalation schedule provided by their healthcare provider to safely achieve the therapeutic dosage [see *Dosage and Administration (2)*]. Inform patients that the tablets may be divided in half at the score, if needed. Instruct patients not to take a double dose to make up for a missed dose.

### Drug Interactions

Instruct patients to notify their healthcare provider prior to starting any new medication, including over-the-counter drugs [see *Drug Interactions (7)*].

### Pregnancy

Instruct patients that if they are pregnant or plan to become pregnant while taking FIRDAPSE they should inform their healthcare provider. Advise patients that there is a pregnancy registry that monitors pregnancy outcomes in women exposed to FIRDAPSE during pregnancy and encourage them to enroll if they become pregnant while taking FIRDAPSE [see *Use in Specific Populations (8.1)*].

### Storage

Advise patients to store FIRDAPSE at 68°F to 77°F (20°C to 25°C).

Distributed by Catalyst Pharmaceuticals, Inc., Coral Gables, FL 33134

FIRDAPSE is a trademark of Catalyst Pharmaceuticals, Inc.

Catalyst and the Catalyst logo are trademarks of Catalyst Pharmaceuticals, Inc.

**MEDICATION GUIDE**  
**FIRDAPSE® (FIR-dapse)**  
**(amifampridine)**  
**tablets, for oral use**

Read this Medication Guide before you start taking FIRDAPSE and each time you get a refill. There may be new information. This information does not take the place of talking with your doctor about your medical condition or your treatment.

**What is the most important information I should know about FIRDAPSE?**

FIRDAPSE can cause seizures.

- You could have a seizure even if you never had a seizure before.
- **Do not** take FIRDAPSE if you have ever had a seizure.

**Stop taking FIRDAPSE and call your doctor right away if you have a seizure while taking FIRDAPSE.**

**What is FIRDAPSE?**

FIRDAPSE is a prescription medicine used to treat Lambert-Eaton myasthenic syndrome (LEMS) in adults.

It is not known if FIRDAPSE is safe or effective in children.

**Do not take FIRDAPSE if you:**

- have ever had a seizure.
- are allergic to amifampridine phosphate, or another aminopyridine.

**Before you take FIRDAPSE, tell your doctor about all of your medical conditions.**

**including if you:**

- are taking another aminopyridine, such as compounded 3,4-diaminopyridine (3,4-DAP)
- have had a seizure
- have kidney problems
- liver problems
- are pregnant or plan to become pregnant. It is not known if FIRDAPSE will harm your unborn baby. You and your doctor will decide if you should take FIRDAPSE while you are pregnant.
- There is a registry for women who become pregnant during treatment with FIRDAPSE. The purpose of this registry is to collect information about your health and your baby's health. Contact the registry as soon as you learn that you are pregnant, or ask your healthcare provider to contact for you by calling 855-212-5856 (toll free), contacting the Fax number 877-867-1874 (toll free), emailing the Pregnancy Coordinating Center at [firdapsepregnancyregistry@ubc.com](mailto:firdapsepregnancyregistry@ubc.com), or visiting the study website [www.firdapsepregnancystudy.com](http://www.firdapsepregnancystudy.com)
- are breastfeeding or plan to breastfeed. It is not known if FIRDAPSE passes into your breast milk. Talk to your doctor about the best way to feed your baby while taking FIRDAPSE.

**Tell your doctor about all the medicines you take,** including prescription and over-the-counter medicines, vitamins and herbal supplements.

**How should I take FIRDAPSE?**

- Take FIRDAPSE exactly as your doctor tells you to take it. Do not change your dose of FIRDAPSE.
- Do not take more than 2 tablets of FIRDAPSE at one time or more than 8 tablets in a 24-hour period.
- FIRDAPSE can be taken with or without food.
- If you miss a dose of FIRDAPSE, skip that dose and take your next dose at your next scheduled dose time. Do not double your dose to make up the missed dose.
- Do not take FIRDAPSE together with other medicines known to increase the risk of seizures.
- If you take too much FIRDAPSE, call your doctor or go to the nearest hospital emergency room right away.

### **What are the possible side effects of FIRDAPSE?**

#### **FIRDAPSE may cause serious side effects, including:**

- **Seizures. See "What is the most important information I should know about FIRDAPSE?"**
- **Serious allergic reactions, such as anaphylaxis.** FIRDAPSE can cause serious allergic reactions. Stop taking FIRDAPSE and call your doctor right away or get emergency medical help if you have:
  - shortness of breath or trouble breathing
  - swelling of your throat or tongue
  - hives

#### **The most common side effects of FIRDAPSE include:**

- tingling around the mouth, tongue, face, fingers, toes, and other body parts
- upper respiratory infection
- stomach pain
- nausea
- diarrhea
- headache
- increased liver enzymes
- back pain
- high blood pressure
- muscle spasms

Tell your doctor if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of FIRDAPSE.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

#### **How should I store FIRDAPSE?**

- Store FIRDAPSE at 68°F to 77°F (20°C to 25°C).
- Safely throw away FIRDAPSE that is out of date or no longer needed.

#### **Keep FIRDAPSE and all medicines out of the reach of children.**

#### **General Information about the safe and effective use of FIRDAPSE**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use FIRDAPSE for a condition for which it was not prescribed. Do not give FIRDAPSE to other people, even if they have the same symptoms that you have. It may harm them.



If you would like more information, talk to your doctor or pharmacist. You can ask your pharmacist or doctor for information about FIRDAPSE that is written for health professionals.

**What are the ingredients in FIRDAPSE?**

**Active ingredient:** amifampridine

**Inactive ingredients:** calcium stearate, colloidal silicon dioxide, and microcrystalline cellulose.

Distributed by Catalyst Pharmaceuticals, Inc., Coral Gables, FL 33134

For more information, go to [www.YourCatalystPathways.com](http://www.YourCatalystPathways.com) or call 1-833-422-8259

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Revised 02/2021

**PRINCIPAL DISPLAY PANEL - 10 mg Tablet Blister Pack Carton**

Rx Only

NDC 69616-211-06

120 Tablets Total

Provided in 12 Child-Resistant Blister Cards

Of 10 Tablets Each

FIRDAPSE®


(amifampridine) Tablets

10 mg per Tablet

Manufactured for:

Catalyst Pharmaceuticals, Inc.

Coral Gables, FL 33134



120 Tablets Total  
Provided in 12 Child-Resistant Blister Cards  
Of 10 Tablets Each

**FIRDAPSE®**  
(amifampridine) Tablets  
10 mg per Tablet

NDC 69616-211-06

Rx Only

Rx Only

NDC 69616-211-06

120 Tablets Total  
Provided in 12 Child-Resistant Blister Cards  
Of 10 Tablets Each

**FIRDAPSE®**  
(amifampridine) Tablets  
10 mg per Tablet

Manufactured for:  
Catalyst Pharmaceuticals, Inc.  
Coral Gables, FL 33134

Rx Only

NDC 69616-211-06

**FIRDAPSE®**  
(amifampridine) Tablets  
10 mg per Tablet

120 Tablets Total  
Provided in 12 Child-Resistant Blister Cards  
Of 10 Tablets Each

Each tablet contains:  
 Amifampridine.....10 mg  
 (Equivalent to 18.98 mg of amifampridine phosphate)

**Usual Dosage:** See package insert for full prescribing information.

**Storage:** Store at 20°C to 25°C (68°F to 77°F), with excursions permitted from 15°C to 30°C (59°F to 86°F)

[See USP controlled room temperature].

Store in the original child resistant package.

For oral use only.

Caution: Federal law prohibits dispensing without a prescription.

Keep out of reach of children.



Rx Only

NDC 69616-211-06



120 Tablets Total  
 Provided in 12 Child-Resistant Blister Cards  
 Of 10 Tablets Each

GTIN XXXXXXXXXX  
 SN XXXXXXXXXXXXXXXX  
 EXP MMYYYY  
 LOT XXXXXXXXXX

# FIRDAPSE

amifampridine phosphate tablet

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:69616-211
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>amifampridine phosphate</b> (UNII: 8HF8FIN815) (amifampridine - UNII:RU4S6E2G0J)	amifampridine	10 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>microcrystalline cellulose</b> (UNII: OP1R32D61U)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>calcium stearate</b> (UNII: 776XM7047L)	

## Product Characteristics

<b>Color</b>	WHITE (white to off white)	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	CATALYST;211;10
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69616-211-06	12 in 1 CARTON	01/07/2019	
1	NDC:69616-211-04	10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:69616-211-08	60 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/07/2019	
3	NDC:69616-211-03	240 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/07/2019	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA208078	01/07/2019	

**Labeler** - Catalyst Pharmaceuticals, Inc. (053742248)

## Establishment

Name	Address	ID/FEI	Business Operations
Patheon API Manufacturing, Inc.		962538372	API MANUFACTURE(69616-211) , analysis(69616-211)

## Establishment

Name	Address	ID/FEI	Business Operations
Catalent Micron Technologies, Inc.		015966157	particle size reduction(69616-211) , ANALYSIS(69616-211)

## Establishment

Name	Address	ID/FEI	Business Operations
Pharmaceutics International, Inc.		878265586	MANUFACTURE(69616-211) , analysis(69616-211)

## Establishment

Name	Address	ID/FEI	Business Operations
PCI Pharma Services Canada, Inc.		204123822	PACK(69616-211)

## Establishment

Name	Address	ID/FEI	Business Operations
Quality Chemical Laboratories, LLC		071344167	ANALYSIS(69616-211)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Patheon Inc.		240769596	ANALYSIS(69616-211)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Patheon Pharmaceuticals Inc.		005286822	MANUFACTURE(69616-211) , PACK(69616-211)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Patheon Softgels Inc.		002193829	ANALYSIS(69616-211)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Robertson-Microlit Laboratories Inc.		177250289	ANALYSIS(69616-211)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
SGS Canada Inc.		203668041	analysis(69616-211)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
SGS North America Inc.		049859261	analysis(69616-211)

Revised: 7/2022

Catalyst Pharmaceuticals, Inc.