

SUDAFED PE HEAD CONGESTION PLUS MUCUS- acetaminophen, guaifenesin, and phenylephrine hydrochloride tablet, film coated
Johnson & Johnson Consumer Inc.

SUDAFED PE Head Congestion + Mucus

Drug Facts

<i>Active ingredients (in each tablet)</i>	<i>Purpose</i>
Acetaminophen 325 mg	Pain reliever/fever reducer
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these symptoms associated with hay fever or other respiratory allergies, and the common cold:
 - sinus congestion and pressure
 - headache
 - minor aches and pains
 - nasal congestion
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dose**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

-
- take 2 tablets every 4 hours

adults and children 12 years and over	▪ do not take more than 10 tablets in 24 hours
children under 12 years	ask a doctor

Other information

- each tablet contains: **sodium 3 mg**
- store between 20-25°C (68-77°F)
- **do not use if blister unit is torn or broken**

Inactive ingredients

carnauba wax, croscarmellose sodium, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, titanium dioxide, triacetin

Questions or comments?

call **1-888-217-2117** (toll-free) or **215-273-8755** (collect)

PRINCIPAL DISPLAY PANEL

PREVIOUSLY SUDAFED PE [®] PRESSURE+PAIN+MUCUS
NDC 50580-447-01

SUDAFED PE[®]

HEAD CONGESTION
+ MUCUS

Acetaminophen, Guaifenesin, Phenylephrine HCl,
Pain Reliever/Fever Reducer, Expectorant, Nasal Decongestant

actual
size

- SINUS PRESSURE
- HEADACHE
- CHEST CONGESTION

24 TABLETS

NON-DROWSY



Made in Italy
 Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
 McKel Consumer Health Care Division
 Fort Washington, PA 19034 USA
 © J&J 2018

Drug Facts (continued)

■ do not use if blister unit is torn or broken
 ■ each tablet contains: sodium 3 mg
 ■ store between 20-25°C (68-77°F)

Drug Facts (continued)

Inactive Ingredients
 carnauba wax, croscarmellose sodium, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrated starch, titanium dioxide, triacetin

Important: Read all product information before using. Keep this box for important information.

Drug Facts

Active Ingredients (in each tablet)
 Acetaminophen 325 mg
 Guaifenesin 200 mg
 Pseudoephedrine HCl 5 mg

Purpose
 Pain reliever/fever reducer
 Expectorant
 Nasal decongestant

Uses
 ■ temporarily relieves these symptoms associated with hay fever or other respiratory allergies, and the common cold: ■ sinus congestion and pressure ■ headache ■ minor aches and pains ■ nasal congestion ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive ■ temporarily reduces fever

Warnings
 Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
 Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ rash ■ skin redness ■ blisters
 If a skin reaction occurs, stop use and seek medical help right away.
 Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks

Directions
 ■ do not take more than directed (see overdose warning) ■ take 2 tablets every 4 hours ■ do not take more than 10 tablets in 24 hours
 ask a doctor
 children under 12 years
 and over

Other information
 ■ do not use if blister unit is torn or broken
 ■ store between 20-25°C (68-77°F)
 ■ each tablet contains: sodium 3 mg

Questions or comments? call 1-888-217-2117 (toll-free) or 1-215-273-8755 (collect)

Does Not Contain Pseudoephedrine

Open from other side

The makers of the SUDAFED® family of products are dedicated to helping you and your family reduce head congestion and sinus pressure.

PREVIOUSLY SUDAFED PE® PRESSURE+PAIN+MUCUS NDC 50580-447-01

SUDAFED PE®

HEAD CONGESTION + MUCUS

Acetaminophen, Guaifenesin, Phenylephrine HCl, Pain Reliever/Fever Reducer, Expectorant, Nasal Decongestant



- SINUS PRESSURE
- HEADACHE
- CHEST CONGESTION

24 TABLETS

NON-DROWSY



30 (42/49) / 120 (478)

SUDAFED PE HEAD CONGESTION PLUS MUCUS

acetaminophen, guaifenesin, and phenylephrine hydrochloride tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50580-447
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	SUPE;SU02
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-447-01	2 in 1 CARTON	06/17/2019	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/17/2019	

Labeler - Johnson & Johnson Consumer Inc. (878046358)

Revised: 1/2024

Johnson & Johnson Consumer Inc.