# FIRST AID ANTIBIOTIC AND PAIN RELIEF- neomycin, polymyxin b, pramoxine hcl cream

## **Meijer Distribution Inc**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

Meijer Distribution, Inc. First Aid Antibiotic + Pain Relief Cream Drug Facts

## Active ingredient (in each gram)

Neomycin sulfate equivalent to 3.5 mg of neomycin base Polymyxin B 10,000 units as polymyxin B sulfate Pramoxine HCl 10 mg

### **Purpose**

First aid antibiotic

External analgesic

#### Uses

first aid to help prevent infection and for temporary relief of pain or discomfort in minor:

- cuts
- scrapes
- burns

# **Warnings**

For external use only.

#### Do not use

- if you are allergic to any of the ingredients
- in the eyes
- over large areas of the body

# Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

# Stop use and ask a doctor if

- you need to use longer than 1 week
- condition persists or gets worse
- symptoms persist for more than 1 week, or clear up and occur again within a few days
- rash or other allergic reaction develops

## Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

#### **Directions**

- adults and children 2 years of age or older:
- clean the affected area
- apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage
- children under 2 years of age: ask a doctor

#### Other information

store at 20°-25°C (68°-77°F)

## **Inactive ingredients**

emulsifying wax, methylparaben, mineral oil, poloxamer 188, propylene glycol, purified water, white petrolatum

#### Questions or comments?

1-800-719-9260

# **Principal Display Panel**

MAXIMUM STRENGTH

first aid antibiotic + pain relief cream

Neomycin Sulfate | Polymyxin B Sulfate | Pramoxine HCl

+ PAIN RELIEF

Sting Free

Soothing Infection Protection

Compare to Neosporin® + Pain Relief

active ingredients

# MAXIMUM STRENGTH

first aid antibiotic + pain relief cream

Neomycin Sulfate | Polymyxin B Sulfate | Pramoxine HCl

+ PAIN RELIEF

NET WT 0.5 OZ (14 g)

First Aid Antibiotic/Pain Relieving Cream



**MAXIMUM STRENGTH** 

# first aid antibiotic + pain relief cream

Neomycin Sulfate | Polymyxin B Sulfate | Pramoxine HCl

#### + PAIN RELIEF

Sting Free Soothing Infection Protection

# Soothing Infection Protection for Cuts | Scrapes | Burns

# meijer

NDC 79481-0809-( Compare to Neosporin® + Pain Relief active ingredients®

# first aid antibiotic + pain relief cream

Neomycin Sulfate | Polymyxin B Sulfate | Pramoxine HCl

+ PAIN RELIEF

NET WT 0.5 OZ (14 g)


First Aid Antibiotic/Pain Relieving Cream

DIST. BY MEIJER DISTRIBUTION, INC. GRAND RAPIDS, MI 49544 www.meijer.com









\*This product is not affiliated with, manufactured by, or produced by the makers of Neosporin®.

#### Drug Facts

#### Active ingredients (in each gram)

# Purpose

**Uses** first aid to help prevent infection and for temporary relief of pain or decomfort in minor:

■ cuts ■ scrapes ■ burns

#### Warnings

For external use only.

#### Do not use

- if you are allergic to any of the ingredents
- in theeyes
- over large areasof the body

#### Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

#### Drug Facts (continued)

- Stop use and ask a doctor if
- you ne ed to use longer than 1 week
- condition persists or gets worse
- symptoms persist for more than 1 we ek, or clear up and oc cur a gain within a few days
- rash or other allergic reaction develops

Keep out of reach of children. If swallowed, get medical help or contact a Roison Control Center righta way (1-800-222-1222)

#### Directions

- adults and children 2 years of ageor older:
- clean the affected area
- apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage
- children under 2 years of age: ask a doctor

Other information store at 20°-25°C (68°-77°F)

Inactive ingredients emulsifying wax, mehylparaben, mineral di, poloxamer 188, propylene glycol, purified water, white petro latum

Questions or comments? 1-800-719-9260





# FIRST AID ANTIBIOTIC AND PAIN RELIEF

neomycin, polymyxin b, pramoxine hcl cream

<b>Product Information</b>	roduct Information		
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79481-0809
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety			
Ingredient Name	<b>Basis of Strength</b>	Strength	
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:116QD7X297)	NEOMYCIN	3.5 mg in 1 g	
PRAMOXINE HYDROCHLORIDE (UNII: 88AYB867L5) (PRAMOXINE - UNII:068X84E056)	PRAMOXINE HYDROCHLORIDE	10 mg in 1 g	
POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII: J2VZ 07J96K)	POLYMYXIN B	10000 [USP'U] in 1 g	

Inactive Ingredients			
Ingredient Name	Strength		
METHYLPARABEN (UNII: A2I8C7HI9T)			
MINERAL OIL (UNII: T5L8T28FGP)			
POLOXAMER 188 (UNII: LQA7B6G8JG)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
WATER (UNII: 059QF0KO0R)			
PETROLATUM (UNII: 4T6H12BN9U)			

Product Characteristics			
Color	WHITE	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79481- 0809-0	1 in 1 CARTON	04/20/2021	
1		14 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:79481- 0809-1	1 in 1 CARTON	06/06/2022	
2		28 g in 1 TUBE; Type 0: Not a Combination Product		

# **Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333B	04/20/2021	

# Labeler - Meijer Distribution Inc (006959555)

Revised: 6/2022 Meijer Distribution Inc