

PAIN RELIEVING GEL-PATCH- lidocaine patch
Shanghai Chuangshi Medical Technology (Group) Co., Ltd.

Zeey, Pain Relieving Gel-Patch, Lidocaine 4%, N 5, N 15, N 18

Active ingredient

Lidocaine 4% Purpose: Topical Anesthetic

Purpose

Topical Anesthetic

Uses

For temporary relief of pain

Warnings

For external use only

Ask a doctor if:

- localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
- Conditions worsen
- Symptoms persists for more than 7 days
- Symptoms clear up and occur again within a few days

When using this product

- Use only as directed
- Avoid contact with the eyes, mucous membranes or rashed
- Do not bandage tightly

Do not use

- More than 1 patch at a time
- On wounds or damaged skin
- With a heating pad
- If you are allergic to any ingredients of this product

Stop use if:

- localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
- Conditions worsen

- Symptoms persists for more than 7 days
- Symptoms clear up and occur again within a few days

Pregnancy or breast feeding

If pregnant or breastfeeding, ask a health professional before use

Keep out of reach of children

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Adult and Children 12 years of age and over:

- Clean and dry affected area
- Remove film from patch and apply to the skin (see illustration)
- Apply 1 patch at a time to affected area, not more than 3 to 4 times daily
- Remove patch from the skin after at most 8-hour application

Children under 12 years of age: Consult a doctor

Dosage forms & strengths

This is a strip, patch dosage form.

The active ingredient strength is 4% w/w.

One patch one time, no more than 8 hours one time.

Inactive ingredients

Glycerin, Sodium Polyacrylate, Dihydroxyaluminum Aminoacetate, Disodium Ethylenediaminetetraacetic Acid, Kaolin, Titanium Dioxide, Polysorbate 80, Propylene Glycol, Tartaric Acid, Polyacrylic Acid, Polyvinylpyrrolidone K90, Methylparaben, Propylparaben, Water

Questions or Comments?

1 646-223-0128 · Lu.FDAmanage@gmail.com

Other information

- Avoid storing product in direct sunlight
- Protect product from excessive moisture

Package label. Principal display panel

Zeeey[®]

NDC#73557-157-01

**FAST ACTING
PAIN RELIEF**

PAIN RELIEVING GEL-PATCH

**Desensitize
aggravated nerves
for temporary
relief of pain**

- Back
- Neck
- Shoulders
- Knees & Elbows

LIDOCAINE 4%

1 PATCH

1 PATCH-5.5" x 3.94" (14cm x 10cm)



PAIN RELIEVING GEL-PATCH

LIDOCAINE 4%

Drug Facts

Active Ingredient	Purpose
Lidocaine 4%	Topical anesthetic

Uses: For temporary relief of pain

Warnings

For external use only

Do not use: - More than 1 patch at a time - On wounds or damaged skin. - With a heating pad. - If you are allergic to any ingredients of this product.

When using this product:

- Use only as directed
- Avoid contact with the eyes, mucous membranes or rashes
- Do not bandage tightly

Stop use and ask a doctor if:

- Localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
- Conditions worsen
- Symptoms persist for more than 7 days
- Symptoms clear up and occur again within a few days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Adult and children 12 years of age and over:

- Clean and dry affected area
- Remove film from patch and apply to the skin (see illustration)
- Apply 1 patch at a time to affected area, not more than 3 to 4 times daily
- Remove patch from the skin after at most 8-hour application

Children under 12 years of age:

- Consult a doctor

Other Information

- Avoid storing product in direct sunlight
- Protect product from excessive moisture

Inactive Ingredients

Dihydroxyaluminum Aminoacetate, Edetate Disodium, Glycerin, Kaolin, Methylparaben, Polysorbate 80, Propylene Glycol, Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polyacrylate, Tartaric Acid, Titanium Dioxide, Water.

Questions or Comments?

1 646-225-0128 • Lu.FDKmanage@gmail.com



LOT



Produced by:

SHANGHAI CHUANGSHI MEDICAL TECHNOLOGY (GROUP) CO.,LTD
No.388, Zhang Jiangtang Rd, Qingpu District, Shanghai, 201716 China
Tel: 86-21-31166566
Fac: 86-21-31166568
E-mail: webmaster@csigroup.com.cn
www.csigroup.com.cn

Made in China



LIDOCAINE 4%

PAIN RELIEVING
GEL-PATCH

NDC#73557-157-05

Zeeey®

FAST ACTING
PAIN RELIEF

PAIN RELIEVING GEL-PATCH



Desensitize
aggravated nerves
for temporary
relief of pain

- Back
- Neck
- Shoulders
- Knees & Elbows

LIDOCAINE 4%

APPLY FOR
8
HOURS

5 PATCHES

CONTAINS FIVE 5.5" x 3.94" INDIVIDUALLY PACKAGED PATCHES

PAIN RELIEVING
GEL-PATCH

LIDOCAINE 4%

HOW TO APPLY



PAIN RELIEVING GEL-PATCH

LIDOCAINE 4%

Drug Facts

Active Ingredient Lidocaine 4% **Purpose** Topical anesthetic

Uses: For temporary relief of pain

Warnings

For external use only

Do not use: • More than 1 patch at a time • On wounds or damaged skin • With a heating pad • If you are allergic to any ingredients of this product

When using this product:

• Use only as directed
• Avoid contact with the eyes, mucous membranes or surfaces
• Do not bandage tightly

Stop use and ask a doctor if:

• Localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
• Conditions worsen
• Symptoms persist for more than 7 days
• Symptoms clear up and occur again within a few days

If pregnant or breast-feeding, ask a health professional before use

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Adult and children 12 years of age and over:

- Clean and dry affected area
- Remove film from patch and apply to the skin (see illustration)
- Apply 1 patch at a time to affected area, not more than 3 to 4 times daily

• Remove patch from the skin after at most 8-hour application
Children under 12 years of age: Consult a doctor

Other Information

- Avoid storing product in direct sunlight
- Protect product from excessive moisture

Inactive Ingredients

Styrene/ethylenehexanoate, Ethylate Dioxidum, Glycerin, Isobutyl Methacrylate, Polyacrylate 80, Propylene Glycol, Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polycrylate, Tartaric Acid, Titanium Dioxide, Water

Questions or Comments?

1-846-224-0728 • lidocaine@zeeey.com

Produced by:

SHANGHAI CHANGSHI MEDICAL TECHNOLOGY (GROUP) CO., LTD.
No. 88, Zhangkang Rd, Dongpu District, Shanghai, China, 201716
Tel: 86-21-31166566
Fax: 86-21-31166568
E-mail: webmaster@csigroup.com
www.csigroup.com.cn

Made in China



LOT



6 924239 705656



LIDOCAINE 4%

PAIN RELIEVING
GEL-PATCH

Zeey®

FAST ACTING
PAIN RELIEF

PAIN RELIEVING GEL-PATCH

Desensitize
aggravated nerves
for temporary
relief of pain

- Back
- Neck
- Shoulders
- Knees & Elbows

LIDOCAINE 4%

15 PATCHES

CONTAINS FIFTEEN 5.5" x 3.94" INDIVIDUALLY PACKAGED PATCHES

APPLY FOR
8 HOURS

PAIN RELIEVING
GEL-PATCH

LIDOCAINE 4%

HOW TO APPLY

1 Remove patch from outer package

2 Partially peel back protective film

3 Place on affected area

4 Remove remaining film while pressing patch to skin

PAIN RELIEVING GEL-PATCH

LIDOCAINE 4%

Drug Facts

Active Ingredient	Purpose
Lidocaine 4%	Topical anesthetic

Uses: For temporary relief of pain

Warnings

For external use only

Do not use: • More than 1 patch at a time • On wounds or damaged skin • With a heating pad • If you are allergic to any ingredients of this product

When using this product: • Use only as directed

• Avoid contact with the eyes, mucous membranes or rashes

• Do not bandage tightly

Stop use and ask a doctor if:

• Localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering

• Conditions worsen

• Symptoms persist for more than 7 days

• Symptoms clear up and occur again within a few days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If you loved, get medical help or contact a Poison Control Center right away.

Directions

Adult and children 12 years of age and over:

- Clean and dry affected area
- Remove film from patch and apply to the skin (see illustration)
- Apply 1 patch at a time to affected area, not more than 3 to 4 times daily
- Remove patch from the skin after at most 8-hour application

Children under 12 years of age: Consult a doctor

Other Information

- Avoid storing product in direct sunlight
- Protect product from excessive moisture

Inactive Ingredients

Dihydroxyaluminum Aminoacetate, Edetate Disodium, Glycerin, Karlin, Methylparaben, Polysorbate 80, Propylene Glycol, Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polycrylate, Tartaric Acid, Triammonium Diacetic Water.

Questions or Comments?

1-866-235-0128 • US-PS@zeey.com

Produced by:

SHANGHAI CHUANGSHI MEDICAL TECHNOLOGY (GROUP) CO., LTD
No. 388, Zhang Jiang Road, Qingshi District, Shanghai, China, 201716
Tel: 86-21-31166568
Fax: 86-21-31166568
E-mail: webmaster@csigroup.com.cn
www.csigroup.com.cn

Made in China





PAIN RELIEVING GEL-PATCH

lidocaine patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73557-157
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	0.04 g in 1 g

Inactive Ingredients

Ingredient Name	Strength
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
GLYCERIN (UNII: PDC6A3COOX)	
POVIDONE K90 (UNII: RDH86HJV5Z)	
METHYLPARABEN (UNII: A2I8C7HI9T)	

PROPYLPARABEN (UNII: Z8IX2SC1OH)
EDETATE DISODIUM (UNII: 7FLD91C86K)
KAOLIN (UNII: 24H4NWX5CO)
TARTARIC ACID (UNII: W4888I119H)
WATER (UNII: 059QF0KO0R)
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)
POLYACRYLIC ACID (250000 MW) (UNII: 9G2MAD7J6W)
SODIUM POLYACRYLATE (2500000 MW) (UNII: 05I15JNI2J)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
DIHYDROXYALUMINUM AMINOACETATE ANHYDROUS (UNII: 1K713C615K)

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73557-157-05	5 in 1 BOX	12/30/2022	
1	NDC:73557-157-01	12 g in 1 PATCH; Type 0: Not a Combination Product		
2	NDC:73557-157-15	15 in 1 BOX	01/09/2024	
2	NDC:73557-157-01	12 g in 1 PATCH; Type 0: Not a Combination Product		
3	NDC:73557-157-18	18 in 1 BOX	01/09/2024	
3	NDC:73557-157-01	12 g in 1 PATCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	12/30/2022	

Labeler - Shanghai Chuangshi Medical Technology (Group) Co., Ltd. (546872672)

Registrant - Shanghai Chuangshi Medical Technology (Group) Co., Ltd. (546872672)

Establishment

Name	Address	ID/FEI	Business Operations
Shanghai Chuangshi Medical Technology (Group) Co., Ltd.		546872672	manufacture(73557-157) , label(73557-157)

Revised: 4/2024

Shanghai Chuangshi Medical Technology (Group) Co., Ltd.