

XERMELO- telotristat ethyl tablet
Lexicon Pharmaceuticals, Inc.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use XERMELO safely and effectively. See full prescribing information for XERMELO.

XERMELO (telotristat ethyl) tablets, for oral use
Initial U.S. Approval: 2017

INDICATIONS AND USAGE

Xermelo is a tryptophan hydroxylase inhibitor indicated for the treatment of carcinoid syndrome diarrhea in combination with somatostatin analog (SSA) therapy in adults inadequately controlled by SSA therapy. (1)

DOSAGE AND ADMINISTRATION

- The recommended dosage of Xermelo in adult patients is 250 mg three times daily for patients whose diarrhea is inadequately controlled by a SSA therapy. (2)
- Take Xermelo with food. (2)
- When short-acting octreotide is used in combination with Xermelo, administer short-acting octreotide at least 30 minutes after administering Xermelo. (2, 7.1)
- Discontinue Xermelo if severe constipation develops. (2, 5.1)

DOSAGE FORMS AND STRENGTHS

Tablets: 250 mg telotristat ethyl (3)

CONTRAINDICATIONS

None. (4)

WARNINGS AND PRECAUTIONS

Constipation: Xermelo reduces bowel movement frequency; monitor patients for constipation and/or severe persistent or worsening abdominal pain. Discontinue Xermelo if severe constipation or abdominal pain develops. (5.1)

ADVERSE REACTIONS

Most common adverse reactions ($\geq 5\%$) are nausea, headache, increased GGT, depression, flatulence, decreased appetite, peripheral edema, and pyrexia. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Lexicon Pharmaceuticals, Inc. at 1-844-539-7427 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

DRUG INTERACTIONS

CYP3A4 Substrates (e.g., midazolam): Efficacy of concomitant drugs may be decreased; monitor for suboptimal efficacy and consider increasing the dose of the concomitant drug. (7.1)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 3/2017

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

5.1 Constipation

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

7 DRUG INTERACTIONS

- 7.1 CYP3A4 Substrates
- 7.2 Short-Acting Octreotide

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Xermelo is indicated for the treatment of carcinoid syndrome diarrhea in combination with somatostatin analog (SSA) therapy in adults inadequately controlled by SSA therapy.

2 DOSAGE AND ADMINISTRATION

The recommended dosage of Xermelo in adult patients is 250 mg three times daily for patients whose diarrhea is inadequately controlled by SSA therapy.

Administration

- Take Xermelo with food [see *Clinical Pharmacology (12.3)*, *Clinical Studies (14)*].
- When short-acting octreotide is used in combination with Xermelo, administer short-acting octreotide at least 30 minutes after administering Xermelo [see *Clinical Pharmacology (12.3)*, *Clinical Studies (14)*].
- If a dose is missed, take the next dose at the regular time. Do not take 2 doses at the same time to make up for a missed dose.
- Discontinue Xermelo if severe constipation develops [see *Warnings and Precautions (5.1)*].

3 DOSAGE FORMS AND STRENGTHS

Tablets: 250 mg telotristat ethyl; white to off-white, coated and oval with “T-E” debossed on one side and “250” debossed on the other side.

4 CONTRAINDICATIONS

None.

5 WARNINGS AND PRECAUTIONS

5.1 Constipation

Xermelo reduces bowel movement frequency. In a 12-week, placebo-controlled trial, in which patients had 4 or greater bowel movements per day, 2 out of 45 patients treated with a higher than recommended dosage of Xermelo reported constipation. In one patient the constipation was serious, resulting in hospitalization. During the 36-week extension period with higher than the recommended dosage, 10 of 115 patients reported constipation: one developed intestinal perforation and one developed obstruction. In another 12-week, placebo-controlled trial in which patients had less than 4 bowel movements per day, 4 out of 25 patients treated with the recommended dosage of Xermelo reported constipation. Given that patients with metastatic carcinoid tumors may have impaired integrity of the gastrointestinal tract wall, monitor for the development of constipation and/or severe, persistent, or worsening abdominal pain in patients taking Xermelo. Discontinue Xermelo if severe constipation or severe persistent or worsening abdominal pain develops [see *Dosage and Administration (2)*, *Adverse Reactions (6.1)*].

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Xermelo was studied in a double-blind, placebo-controlled clinical trial of 90 patients with metastatic neuroendocrine tumors and carcinoid syndrome diarrhea. Patients reported between 4 to 12 bowel movements daily despite the use of SSA therapy at a stable dose for at least 3 months [see *Clinical Studies (14)*]. Placebo or Xermelo 250 mg was administered three times daily for 12 weeks. Concomitant anti-diarrheal medications (e.g., loperamide) were used by 43% (36% and 51% in the placebo and Xermelo group, respectively), pancreatic enzyme replacement medications by 39% (36% and 42% in the placebo and Xermelo group, respectively), and opioid analgesics by 29% (24% and 33% in the placebo and Xermelo group, respectively) of patients during the 12-week double-blind period of the trial.

Table 1 below lists adverse reactions occurring at an incidence of at least 5% in the Xermelo group (N=45) and at an incidence greater than placebo (N=45) during the 12-week placebo-controlled period of the trial.

Table 1: Percent Common Adverse Reactions^a by Treatment Group at 12-Weeks in a Double-Blind Placebo-Controlled Clinical Trial of Patients with

Carcinoid Syndrome Diarrhea

Adverse Reaction	Xermelo 250 mg Three Times Daily, N=45 (%)	Placebo, N=45 (%)
Nausea	13	11
Headache	11	4
Increased gamma-glutamyl-transferase (GGT)	9	0
Depression ^b	9	7
Peripheral edema	7	2
Flatulence	7	2
Decreased appetite	7	4
Pyrexia	7	4

^a incidence of at least 5% in the Xermelo group and at an incidence greater than placebo

^b including depression, depressed mood and decreased interest

In another placebo-controlled clinical trial of patients with carcinoid syndrome diarrhea and less than 4 bowel movements per day, the following additional adverse reactions, not listed in Table 1, of abdominal pain (including upper and lower abdominal pain, abdominal distention and gastrointestinal pain) and constipation were reported in at least 5% of patients in the Xermelo treated group and at an incidence greater than placebo [see *Warnings and Precautions (5.1)*].

Less Common Adverse Reactions:

The following is a list of adverse reactions occurring in less than 5% of patients receiving Xermelo during the 12-week placebo-controlled period of the clinical trial:

Investigations: increased alkaline phosphatase, increased alanine aminotransferase, and increased aspartate aminotransferase.

Fecaloma was reported in one patient treated with Xermelo during the 36-week open-label extension period following the 12-week double-blind period of the trial.

7 DRUG INTERACTIONS

7.1 CYP3A4 Substrates

Concomitant use of Xermelo may decrease the efficacy of drugs that are CYP3A4 substrates (e.g., midazolam) by decreasing their systemic exposure. Monitor for suboptimal efficacy and consider increasing the dose for concomitant CYP3A4 substrates, if necessary [see *Clinical Pharmacology (12.3)*].

7.2 Short-Acting Octreotide

Concurrent administration of short-acting octreotide with Xermelo significantly decreased the systemic exposure of telotristat ethyl and telotristat, the active metabolite. If treatment with short-acting octreotide is needed in combination with

Xermelo, administer short-acting octreotide at least 30 minutes after administration of Xermelo [see *Clinical Pharmacology (12.3)*, *Clinical Studies (14)*].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There are no human data with Xermelo use in pregnant women to inform a drug-associated risk. In animal reproduction studies, no effects on embryo-fetal development were observed with the administration of oral telotristat ethyl to rats during organogenesis at doses up to 750 mg/kg/day (approximately 9 times the exposure at the RHD [recommended human dose]). Treatment of pregnant rabbits with oral telotristat ethyl during organogenesis produced maternal toxicity and post-implantation loss at doses of 250 mg/kg/day or higher (approximately 15 times the exposure at the RHD), and reduced fetal weight at 500 mg/kg/day (approximately 33 times the exposure at the RHD). In a pre-/postnatal development study, an increased incidence of mortality in rat offspring was observed during postnatal days 0 to 4 at the maternal oral dose of 500 mg/kg/day (approximately 5 times the exposure at the RHD), given during organogenesis through lactation [see *Data*].

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

Data

Animal data

An embryo-fetal development study performed in rats with oral telotristat ethyl at doses up to 750 mg/kg/day (approximately 9 times the AUC [area under the plasma concentration-time curve] for the active metabolite at the RHD) during organogenesis produced no harm to embryo-fetal development.

In pregnant rabbits treated orally with telotristat ethyl during organogenesis, an increased incidence of post-implantation loss at doses of 250 and 500 mg/kg/day (approximately 15 times the AUC for the active metabolite at RHD) and a decrease in fetal weight at 500 mg/kg/day (approximately 33 times the AUC for the active metabolite at the RHD) was observed. The adverse effects on embryo-fetal development were associated with maternal toxicity (impaired weight gain and/or mortality) at 250 and 500 mg/kg/day. No adverse effects on embryo-fetal development were observed at 125 mg/kg/day (approximately 5 times the AUC for the active metabolite at the RHD).

A pre-/postnatal development study was conducted in rats using oral administration of 100, 200, and 500 mg/kg/day telotristat ethyl during organogenesis through lactation. An increased incidence of pup mortality was observed during postnatal days 0 to 4 at the maternal dose of 500 mg/kg/day (approximately 5 times the AUC for the active metabolite at the RHD). No developmental abnormalities or effects on growth, learning and memory, or reproductive performance were observed through maturation of offspring at maternal doses of up to 500 mg/kg/day in surviving offspring.

8.2 Lactation

Risk Summary

There are no data on the presence of telotristat ethyl in human or animal milk, the effects on the breastfed infant, or the effects on milk production. The effects of local gastrointestinal and systemic exposure to telotristat ethyl on breastfed infants are unknown. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Xermelo and any potential adverse effects on the breastfed infant from Xermelo or from the underlying maternal condition.

Clinical Considerations

Monitor the breastfed infant for symptoms of constipation.

8.4 Pediatric Use

The safety and effectiveness of Xermelo in pediatric patients have not been established.

8.5 Geriatric Use


Of 45 patients in a clinical trial of Xermelo, 19 (42%) patients were 65 years of age and over. No overall differences in safety or effectiveness were observed between these patients and younger patients, and other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

11 DESCRIPTION

Xermelo (telotristat ethyl) tablets contain telotristat ethyl as telotristat etiprate, a tryptophan hydroxylase inhibitor. Telotristat etiprate is the hippurate salt of telotristat ethyl [(S)-ethyl 2-amino-3-(4-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)phenyl)propanoate], which undergoes hydrolysis to the active metabolite, (S)-2-amino-3-(4-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)phenyl)propanoic acid.

The molecular formula of telotristat etiprate is $C_{27}H_{26}ClF_3N_6O_3 \cdot C_9H_9NO_3$ and its molecular weight is 754.2. The molecular weight of the free base (telotristat ethyl) is 575.0.

Chemical Structure:



Telotristat etiprate is a white to off-white solid. The solubility is a function of pH at 25°C; at pH 1 (0.1N HCl), the solubility is greater than 71 mg/mL., at pH 3 phosphate buffer, the solubility is 0.30 mg/mL, at a pH of 5 to 9, the solubility is negligible. In organic solvents, telotristat etiprate is freely soluble in methanol, soluble in acetone, and sparingly soluble in ethanol.

Each Xermelo tablet contains 250 mg of telotristat ethyl (free base) which is equivalent to 328 mg telotristat etiprate. The inactive ingredients of Xermelo tablets include: colloidal silicon dioxide, croscarmellose sodium, hydroxypropyl cellulose, lactose anhydrous, macrogol/PEG, magnesium stearate, polyvinyl alcohol [part hydrolyzed], talc and titanium dioxide.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Telotristat, the active metabolite of telotristat ethyl, is an inhibitor of tryptophan hydroxylase, which mediates the rate limiting step in serotonin biosynthesis. The *in vitro* inhibitory potency of telotristat towards tryptophan hydroxylase is 29 times higher than that of telotristat ethyl. Serotonin plays a role in mediating secretion, motility, inflammation, and sensation of the gastrointestinal tract, and is over-produced in patients with carcinoid syndrome. Through inhibition of tryptophan hydroxylase, telotristat and telotristat ethyl reduce the production of peripheral serotonin, and the frequency of carcinoid syndrome diarrhea.

12.2 Pharmacodynamics

In healthy subjects, telotristat ethyl 500 mg three times daily (twice the recommended dosage) for 14 days decreased whole blood serotonin and 24-hour urinary 5-hydroxyindolacetic acid (u5-HIAA) from baseline. A decrease in 24-hour u5-HIAA was observed as early as after 5 days of treatment.

In patients with metastatic neuroendocrine tumors and carcinoid syndrome diarrhea, 24-hour u5-HIAA decreased from baseline following 6 and 12 weeks of treatment with Xermelo 250 mg three times a day, whereas placebo did not decrease u5-HIAA.

Cardiac Electrophysiology

At a dose 6 times the recommended dose of 250 mg, Xermelo does not prolong the QT interval to any clinically relevant extent.

12.3 Pharmacokinetics

Absorption

After a single oral dose of telotristat ethyl to healthy subjects, telotristat ethyl was absorbed and metabolized to its active metabolite, telotristat. Peak plasma concentrations of telotristat ethyl were achieved within 0.5 to 2 hours, and those of telotristat within 1 to 3 hours. Plasma concentrations thereafter declined in a biphasic manner. Following administration of a single 500 mg dose of telotristat ethyl (twice the recommended dosage) under fasted conditions in healthy subjects, the mean C_{max} and AUC_{0-inf} were 4.4 ng/mL and 6.23 ng•hr/mL, respectively for telotristat ethyl. The mean C_{max} and AUC_{0-inf} were 610 ng/mL and 2320 ng•hr/mL, respectively for telotristat. Peak plasma concentrations and AUC of telotristat ethyl and telotristat appeared to be dose proportional following administration of a single dose of telotristat ethyl in the range of 100 mg (0.4 times the lowest recommended dose to 1000 mg [4 times the highest recommended dose]) under fasted conditions.

Following multiple-dose administration of telotristat ethyl 500 mg three times daily, there was negligible accumulation at steady state for both telotristat ethyl and telotristat.

In patients with metastatic neuroendocrine tumors and carcinoid syndrome diarrhea treated with SSA therapy, the median T_{max} for telotristat ethyl and telotristat was approximately 1 and 2 hours, respectively. Following administration of 500 mg telotristat ethyl three times daily, with meals in patients, the mean C_{max} and AUC_{0-6hr} were approximately 7 ng/mL and 22 ng•hr/mL, respectively, for telotristat ethyl. The mean C_{max} and AUC_{0-6hr} were approximately 900 ng/mL and 3000 ng•hr/mL, respectively for telotristat. The pharmacokinetic parameters for both telotristat ethyl and telotristat were highly variable with about 55% coefficient of variation.

Food Effect

Administration of a single 500 mg dose of Xermelo (twice the recommended dose) with food resulted in higher exposure to both telotristat ethyl and telotristat. The systemic exposure to telotristat ethyl, was significantly increased following administration with a high-fat meal, with C_{max} , and AUC_{0-inf} being 112%, and 264% higher, respectively compared to the fasted state. Following administration of a single 500 mg dose of telotristat ethyl under the fed conditions in healthy subjects, the mean C_{max} and AUC_{0-inf} were 10.5 ng/mL and 21.6 ng•hr/mL, respectively for telotristat ethyl. The C_{max} and AUC_{0-inf} values for telotristat were also increased by 47% and 33%, respectively, with a high-fat meal compared to the fasted state. The mean C_{max} and AUC_{0-inf} were 908 ng/mL and 2980 ng•hr/mL, respectively for telotristat under the fed condition. [see *Dosage and Administration* (2)].

Distribution

Both telotristat ethyl and telotristat are greater than 99% bound to human plasma proteins.

In vitro data suggests that telotristat is a substrate of P-glycoprotein.

Elimination

Following a single 500 mg oral dose of telotristat ethyl in healthy subjects, the apparent half-life was approximately 0.6 hours for telotristat ethyl and 5 hours for telotristat. The apparent total clearance at steady state (CL/F_{SS}) following oral dosing with telotristat ethyl 500 mg three times daily for 14 days (twice the recommended dosage) in healthy subjects was 2.7 and 152 L/hr for telotristat ethyl and telotristat, respectively.

Metabolism

After oral administration, telotristat ethyl undergoes hydrolysis via carboxylesterases to telotristat, its active metabolite. Telotristat is further metabolized. Among the metabolites of telotristat, the systemic exposure to an acid metabolite of oxidative deaminated decarboxylated telotristat was about 35% of that of telotristat. *In vitro* data suggest that telotristat ethyl and telotristat are not substrates for CYP enzymes.

Excretion

Following a single 500 mg oral dose of ^{14}C -telotristat ethyl, 93.2% of the dose was recovered over 240 hours: 92.8% was recovered in the feces, with less than 0.4% being recovered in the urine.

Specific Populations

Age and Sex

Population pharmacokinetic analysis indicated that age (18 to 83 years) and sex do not affect the pharmacokinetics of telotristat.

Renal Impairment

Population pharmacokinetic analysis indicated that creatinine clearance (20 to 89 mL/min) does not affect the pharmacokinetics of telotristat. Xermelo was not studied in end-stage renal disease (ESRD) patients who require dialysis.

Hepatic Impairment

Population pharmacokinetic analysis indicated that mild hepatic impairment (defined as total bilirubin greater than 1 to 1.5 times the upper limit of normal [ULN] or AST greater than the ULN) does not affect the pharmacokinetics of telotristat. The effect of moderate or severe hepatic impairment (defined as total bilirubin greater than 1.5 times the ULN and any value for AST) is unknown.

Drug Interaction Studies

Effect of Telotristat Ethyl on Other Drugs

In vitro studies

The potential for telotristat ethyl and telotristat to induce CYP enzymes (1A2 and 2B6) or inhibit CYP enzymes (2B6, 2C8, and 2C9) has not been adequately studied *in vitro*.

In vitro telotristat ethyl inhibited P-glycoprotein (P-gp) and breast cancer resistance protein (BCRP), but telotristat did not inhibit P-gp and BCRP at the clinically relevant concentrations. However, *in vivo* drug interaction potential via inhibition of BCRP is low based on *in vitro* studies and *in vivo* findings.

Based on *in vitro* studies, *in vivo* drug interaction potential via inhibition of organic cation transporter 1 (OCT1), OCT2, organic anion transporter 1 (OAT1), OAT3, organic anion transporting polypeptide 1B1 (OATP1B1), OATP1B3, or bile salt export pump (BSEP)

transporters by telotristat ethyl and telotristat is low at the recommended dosage.

Midazolam (sensitive CYP3A4 substrate)

Following administration of multiple doses of telotristat ethyl, the systemic exposure to concomitant midazolam was significantly decreased. When 3 mg midazolam was co-administered orally after 5 day treatment with telotristat ethyl 500 mg three times daily (twice the recommended dosage), the mean C_{max} , and AUC_{0-inf} for midazolam were decreased by 25%, and 48%, respectively, compared to administration of midazolam alone. The mean C_{max} , and AUC_{0-inf} for the active metabolite, 1'-hydroxymidazolam, were also decreased by 34%, and 48%, respectively. The reduction in the systemic exposure to both midazolam and its active metabolite suggests that the glucuronidation of 1'-hydroxymidazolam may have been increased by telotristat ethyl [see *Drug Interactions (7.1)*].

Fexofenadine (sensitive P-gp substrate)

The C_{max} and AUC of fexofenadine increased by 16% when a single 180 mg dose of fexofenadine was co-administered orally with the final dose of telotristat ethyl 500 mg administered three times daily (twice the recommended dosage) for 5 days. Clinically meaningful interactions with P-gp substrates are unlikely.

Effect of Other Drugs on Telotristat Ethyl

Short-Acting Octreotide

The mean C_{max} and AUC_{0-last} of telotristat ethyl were decreased by 86% and 81%, respectively, following administration of a single 500 mg dose of Xermelo (twice the recommended dose), co-administered with short-acting octreotide 200 mcg injected subcutaneously in healthy subjects. The mean C_{max} and AUC_{0-last} of telotristat were decreased by 79% and 68%, respectively [see *Clinical Studies (14)*].

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

In a 26-week study in transgenic (Tg.rasH2) mice, telotristat ethyl was not tumorigenic at oral doses up to 300 mg/kg/day.

Telotristat ethyl was negative in the *in vitro* Ames test, the *in vitro* chromosomal aberration test using Chinese hamster ovary cells, and the *in vivo* rat micronucleus test.

Telotristat ethyl at oral doses up to 500 mg/kg/day (approximately 5 times the AUC for the active metabolite at the RHD) was found to have no effect on fertility and reproductive performance of male or female rats.

14 CLINICAL STUDIES

A 12-week double-blind, placebo-controlled, randomized, multicenter trial of Xermelo was conducted in adult patients with a well-differentiated metastatic neuroendocrine tumor and carcinoid syndrome diarrhea who were having between 4 to 12 daily bowel movements despite the use of SSA therapy at a stable dose for at least 3 months. Patients were randomized to placebo or treatment with Xermelo 250 mg three times

daily.

Study medication was administered within 15 minutes before or within 1 hour after a meal or snack [see *Dosage and Administration (2)*]. All patients were required to stay on their baseline SSA regimen and were allowed to use rescue medication (short-acting octreotide) and antidiarrheals (e.g., loperamide) for symptomatic relief. A total of 90 patients were evaluated for efficacy. The mean age of the population was 63 years of age (range 37 to 83 years), 50% were male, and 90% were White.

The primary efficacy endpoint was the change from baseline in the number of daily bowel movements averaged over the 12-week treatment period. The analysis results can be found in Table 2 below. The average was based on the number of days with valid, non-missing data. When a patient had more than 6 weeks of missing data, the change from baseline was considered equal to zero. A week of missing data was defined as a patient missing at least 4 days of diary data in that week.


Table 2: Change from Baseline in Bowel Movements/Day Averaged Over 12 Weeks in Adult Patients with Carcinoid Syndrome Diarrhea

Parameter		Xermelo 250 mg three times daily	Placebo
Bowel Movements/Day At Baseline^a	Number of Patients	45	45
	Baseline Mean (SD)	6.1 (2.1)	5.2 (1.4)
	Median (Min, Max)	5.5 (3.5, 13.0)	5.1 (3.5, 9.0)
Change From Baseline In Bowel Movements/Day Averaged Over 12 Weeks	Change Averaged over 12 Weeks: Mean (SD)	-1.4 (1.4)	-0.6 (0.8)
	Median (Min, Max)	-1.3 (-6.1, 1.6)	-0.6 (-2.7, 0.8)
	Estimate of Treatment Difference (97.5% CL) ^b	-0.8 ^c (-1.3, -0.3)	---
CL=confidence limit; SD=standard deviation.			
^a Baseline Bowel Movements/Day was assessed over the 3-4 week screening/run-in period.			
^b Statistical tests used a blocked 2-sample Wilcoxon Rank Sum statistic (van Elteren test) stratified by the u5-HIAA stratification at randomization. CLs were based on the Hodges-Lehmann estimator of the median paired difference.			
^c p<0.001			

In the 12-week study, a difference in average weekly reductions in bowel movement frequency between Xermelo and placebo was observed as early as 1 to 3 weeks, and persisted for the remaining 9 weeks of the study.

To aid in the interpretation of the bowel movement reduction results, the proportion of patients reporting any particular level of reduction in overall average bowel movement frequency is depicted in Figure 1 below. For example, 33% of patients randomized to Xermelo and 4% of patients randomized to placebo experienced a reduction in overall average bowel movements from baseline of at least 2 bowel movements per day.

Figure 1: Cumulative Proportion of Patients with Carcinoid Syndrome Diarrhea Reporting Change in Overall Average Bowel Movement Frequency



Other symptoms of carcinoid syndrome (abdominal pain or flushing) did not show improvement in the comparison of Xermelo to placebo.

The average number of daily short-acting octreotide injections used for rescue therapy over the 12-week double-blind treatment period was 0.3 and 0.7 in the Xermelo and placebo groups, respectively. In the subgroup of patients who received short-acting octreotide injections, observed reductions in the number of bowel movements per day and treatment differences were generally consistent with the reductions and differences observed in patients who did not receive rescue therapy, and were similar to the overall data presented in Table 2 above [see *Dosage and Administration (2)*, *Drug Interactions (7.2)*].

A third randomized treatment arm of Xermelo 500 mg three times daily did not demonstrate additional treatment benefit on the primary endpoint and had a greater incidence of adverse reactions than Xermelo 250 mg three times daily. Therefore, Xermelo 500 mg three times daily is not recommended [see *Dosage and Administration (2)*].

16 HOW SUPPLIED/STORAGE AND HANDLING

How Supplied

250 mg tablet: white to off-white coated oval tablet with “T-E” debossed on one side and “250” debossed on the other side.

Xermelo is dispensed in a monthly case for a total of 28 days of therapy. Each monthly case contains four weekly boxes. Each weekly box contains seven daily dose packs (day pack).

- NDC 70183-125-84: Monthly case of 84 tablets. Each child resistant daily dose pack (day pack) contains three 250 mg tablets.

Storage

Store at 25°C (77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].

17 PATIENT COUNSELING INFORMATION

Advise patients:

- If they experience severe constipation or severe persistent or worsening abdominal pain, to discontinue Xermelo and contact their healthcare provider [see *Warnings and Precautions (5.1)*].
- To take Xermelo with food [see *Clinical Pharmacology (12.3)*, *Clinical Studies (14)*].
- When short-acting octreotide is used in combination with Xermelo, administer short-acting octreotide at least 30 minutes after administering Xermelo [see *Clinical Pharmacology (12.3)*, *Clinical Studies (14)*].
- If a dose is missed, take the next dose at the regular time. Do not take 2 doses at the same time to make up for a missed dose.

Distributed by:

Lexicon Pharmaceuticals, Inc.
8800 Technology Forest Place
The Woodlands, TX 77381

Principal Display Panel - ShellPak Label

NDC 70183-125-03

Rx Only

To Open:

1. Press and hold button below
2. Pull out blister from the right

XERMELO™

(telotristat ethyl) tablets

250 mg
per tablet

This daily dose pack contains

3 TABLETS

**For oral use only. As with all medications,
keep out of reach of children.**

To Open:
1. Press and hold button below
2. Pull out blister from the right

NDC 70183-125-03
Rx Only

XERMELO™
(telotristat ethyl) tablets

250 mg
per tablet

This daily dose pack contains
3 TABLETS

TE05001R1

Lexicon®
pharmaceuticals

For oral use only. As with all medications,
keep out of reach of children.

1. PRESS
2. PULL

Principal Display Panel - 28 Day Case Label

NDC 70183-125-84

28 DAY CASE

Rx Only

84 Tablets

XERMELO™
(telotristat ethyl) tablets

250 mg
per tablet

Lexicon®
pharmaceuticals

For oral use only.

As with all medications, keep out of reach of children.



XERMELO

telotristat ethyl tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70183-125
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
telotristat ethyl (UNII: 8G388563M7) (telotristat - UNII:381V4FCV2Z)	telotristat ethyl	250 mg

Inactive Ingredients

Ingredient Name	Strength
anhydrous lactose (UNII: 3SY5LH9PMK)	
hydroxypropyl cellulose (90000 wamw) (UNII: UKE75GEA7F)	
croscarmellose sodium (UNII: M28OL1HH48)	
magnesium stearate (UNII: 70097M6I30)	
silicon dioxide (UNII: ETJ7Z6XBU4)	
polyvinyl alcohol, unspecified (UNII: 532B59J990)	

titanium dioxide (UNII: 15FIX9V2JP)	
polyethylene glycol, unspecified (UNII: 3WJQ0SDW1A)	
talc (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	white (white to off-white)	Score	no score
Shape	OVAL (OVAL)	Size	17mm
Flavor		Imprint Code	T;E;250
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70183-125-84	4 in 1 CASE	03/01/2017	
1	NDC:70183-125-21	7 in 1 BOX		
1	NDC:70183-125-03	3 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA208794	03/01/2017	

Labeler - Lexicon Pharmaceuticals, Inc. (838179638)

Revised: 10/2022

Lexicon Pharmaceuticals, Inc.